

DIRECTIONS

of federal law to use this product inconsistent with its labeling

PRACTICAL TREATMENT—In case of with soap and water; for eyes flush minutes and get medical attention.

CAUTIONARY STATEMENTS

Humans & Domestic Animals Cautioned. Avoid contact with skin, eyes or breathing spray mist. Wash thoroughly.

Environmental Hazards

streams or ponds. Do not apply when favor drift from target areas. Avoid edible plants, as this product may injure, grapes, roses, tomatoes, and flowers and ornamental plants. Coarse low pressure are less likely to drift. Do not use on grass seed crop until seedlings less than two months old. Do not use equipment used to apply insecticides or fungicides to turf may be damaged. Do not use for irrigation or domestic purposes.

Storage and Disposal

Do not store near water, food or feed by storage, disposal of equipment. Do not store near other equipment. Do not reuse empty container. Wrap in trash collection. Assume all risk and liability for use. Labeling not in accordance with the terms



DANDELION and CHICKWEED KILLER

ACTIVE INGREDIENTS
Dimethylamine salt of 2,4-dichlorophenoxy acetic acid 25.00%
Dimethylamine salt of Dichloro (3,4-Dichloro) vanilic acid 7.28%
INERT INGREDIENTS 67.72%
Total 100.00%
2,4-dichlorophenoxyacetic acid equivalent 18.3% or 1.83 lbs per gallon
3,4-dichloro-vanilic acid equivalent 1.83% or .183 lbs per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

See Side Panel for Additional Precaution Statements and Statement of Ecological Treatment.
EPA Reg. No. 21693-3-1 EPA Reg. No. 21693-3-1
NET CONTENTS: 1 QUART

GENERAL INFORMATION

Dandelion and Chickweed Killer may be used for the control of broadleaved weeds in lawns, ornamental turf, vacant lots and fence rows. It is most effective when weeds are small and actively growing and should be applied when the temperatures range from 65 degrees to 90 degrees F. Do not apply on a windy day. If applied during extreme temperatures weed control will be erratic. Do not apply if rain is expected in 2 hours. Application may be made with power, knapsack or hose-attached sprayers. Do not treat newly seeded lawns until first cutting. Make sure to add Patterson's Spreader Sticker to the spray for best weed control.

LAWNS AND ORNAMENTAL TURF: How to Mix. Mix 3 tablespoons (1 1/2 oz.) per 2-3 gallons water to treat 1,000 sq. ft. using a pressure sprayer or a hose-end applicator. Add spreader sticker to the spray solution for best weed kill. Controls: Beggar-Tick, Bedstraw, Buckhorn, Chickweed, Clover, Dandelion, Dock, Ground Ivy, Heal-all, Henbit, Knotweed, Lambsquarter, Lespedeza, Morning Glory, Peppergrass, Plantains, Poison Ivy, Purslane, Ragweed, Sheep Sorrel, Shepards Purse, Speedwell, Spurge, Wild Garlic, Wild Onion, Yarrow.

VACANT LOTS AND SIMILAR NON-CROP AREAS: Mix 3 tablespoons per 5-10 gallons water to cover 1,000 sq. ft. Make sure to add spreader sticker to spray. Wet foliage thoroughly. Repeat treatment where heavy weeds appear to resist spray. Will control weeds as listed above.

SPOT SPRAYING: Spray anytime of the year. Add 3 tablespoons per gallon water. Thoroughly wet weeds.

APPLICATION CHART

Amount of Spray	Amount of Water	To Cover
3 tbsps (1 1/2 oz.)	2 Gallons	1,000 sq. ft.
15 tbsps (7 1/2 oz.)	10-15 Gallons	5,000 sq. ft.
1 Gallon	80-100 Gallons	85,000 sq. ft.