

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.
STATEMENT OF PRACTICAL TREATMENT—In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get medical attention.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals Caution

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

Environmental Hazards


Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from target areas. Avoid spray drift to susceptible plants, as this product may injure beans, cucumbers, grapes, roses, tomatoes, and other vegetables, flowers and ornamental plants. Coarse sprays applied at low pressure are less likely to drift. Do not exceed prescribed dosage rates. Do not use equipment used for this product to apply insecticides or fungicides to susceptible plants or seeds. Do not use on grass seed crop or on turf containing bent grasses. Do not apply to Dichondra, St. Augustine, Carpet grass or clover lawns. Do not use on new grass seedlings less than two months old. Clover present in turf may be damaged. Do not contaminate water used for irrigation or domestic purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides or seeds. Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Buyer assumes all risk and liability for use, storage, and/or handling not in accordance with the terms of this label.

REGISTERED
U.S. PAT. & TM. OFF.
EPA Reg. No. 2169-263
EPA Reg. No. 2169-KS1



**DANDELION, CHICKWEED
and
BROADLEAF WEED KILLER**

ACTIVE INGREDIENTS:
*Dimethylamine Salt of 2,4-Dichlorophenoxy Acetic Acid 14.22%
**Dimethylamine Salt of Dicamba (3,6-Dichloro-o-anisic Acid) 84.82%
INERT INGREDIENTS 100.96%
Total 100.00%
*2,4-D Acid Equivalent 11.8% or 1.03 lbs. 2,4-D acid per gallon
**3,6-Dichloro-o-anisic Acid Equivalent 79% or 0.7 lbs. per gallon

WARNING
KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Precautionary Statements
and Statement of Practical Treatment
EPA Reg. No. 2169-263 EPA Reg. No. 2169-KS1
NET CONTENTS: 1 QUART

GENERAL INFORMATION

Dandelion, Chickweed and Broadleaf Weed Killer is used for the control of broadleaf weeds in lawn, vacant lots and fence line. Apply when weeds are small and before they seed. Apply when temperature is between 50 degrees to 90 degrees F. Do not apply during extreme temperatures. Application may be made with attached sprayers. Do not treat immediately after first cutting. Make sure to apply the spray for best results.

LAWNS AND ORNAMENTAL PLANTS: 6 tablespoons (3 ozs.) per 2-3 sq. ft. using a pressure sprayer. Add spreader sticker to the spray. Controls: Beggar-Tick, Blackberry, Clover, Dandelion, Dock, Groundsage, Knotweed, Lambsquarters, Plantain, Poison Ivy, Sheep Sorrel, Shepherds Purse, Garlic, Wild Onion, Yarrow.

VACANT LOTS AND SIMILAR AREAS: 6 tablespoons per 5-10 gallons of water. Make sure to add spreader sticker to the spray. Repeat treatment as needed to resist spray. Will control weeds.


SPOT SPRAYING: Spray 6-10 tablespoons per gallon of water. Amount of Spray: 6 tbsp (3 oz.) 2.5 gal, 30 tbsp (15 oz.) 10.1 gal.

DIRECTIONS
 of federal law to use this product
 inconsistent with its labeling
PRACTICAL TREATMENT—In case of
 with soap and water; for eyes flush
 minutes and get medical attention.
CAUTIONARY STATEMENTS
Humans & Domestic Animals Caution
 ed. Avoid contact with skin, eyes or
 bathing spray mist. Wash thoroughly

Environmental Hazards
 streams or ponds. Do not apply when
 favor drift from target areas. Avoid
 edible plants, as this product may in-
 bers, grapes, roses, tomatoes, and
 lowers and ornamental plants. Coarse
 low pressure are less likely to drift. Do
 ed dosage rates. Do not use equipment
 to apply insecticides or fungicides to
 r seeds. Do not use on grass seed crop-
 ing bent grasses. Do not apply to Di-
 line, Carpet grass or clover lawns. Do
 ss seedlings less than two months old.
 urf may be damaged. Do not contamin-
 irrigation or domestic purposes.

Storage and Disposal
 e water, food or feed by storage, dis-
 of equipment. Do not store near other
 . Do not reuse empty container. Wrap
 in trash collection.

assumes all risk and liability for use.
 idling not in accordance with the terms



DANDELION, CHICKWEED and BROADLEAF WEED KILLER

ACTIVE INGREDIENTS:
 •Dimethylamine Salt of 2,4-Dichlorophenoxy Acetic Acid 14.22%
 •Dimethylamine Salt of Dicamba (3,6-Dichloro-o-anisic Acid) .95%
INERT INGREDIENTS 84.83%
Total 100.00%
 •2,4-D Acid Equivalent 11.8% or 1.03 lbs. 2,4-D acid per gallon
 •3,6-Dichloro-o-anisic Acid Equivalent .79% or .07 lbs. per gallon

WARNING
KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Precautionary Statements
 and Statement of Practical Treatment
 EPA Reg. No. 2109-263 EPA Reg. No. 2109-KS-1

NET CONTENTS: 1 QUART

GENERAL INFORMATION

Dandelion, Chickweed and Broadleaf Weed Killer may be used for the control of broadleaved weeds in lawns, ornamental turf, vacant lots and fence rows. It is most effective when weeds are small and actively growing and should be applied when the temperatures range from 65 degrees to 90 degrees F. Do not apply on a windy day. If applied during extreme temperatures weed control will be erratic. Do not apply if rain is expected in 2 hours. Application may be made with power, knapsack or hose-attached sprayers. Do not treat newly seeded lawns until first cutting. Make sure to add Patterson's Spreader Sticker to the spray for best weed control.

LAWNS AND ORNAMENTAL TURF: How to Mix: Mix 6 tablespoons (3 ozs.) per 2-3 gallons water to treat 1,000 sq. ft. using a pressure sprayer or a hose-end applicator. Add spreader sticker to the spray solution for best weed kill. Controls: Beggar-Tick, Bedstraw, Buckhorn, Chickweed, Clover, Dandelion, Dock, Ground Ivy, Heal-all, Henbit, Knotweed, Lambsquarter, Lespedeza, Morning Glory, Peppergrass, Plantains, Poison Ivy, Purslane, Ragweed, Sheep Sorrel, Shepards Purse, Speedwell, Spurge, Wild Garlic, Wild Onion, Yarrow.

VACANT LOTS AND SIMILAR NON-CROP AREAS: Mix 6 tablespoons per 5-10 gallons water to cover 1,000 sq. ft. Make sure to add spreader sticker to spray. Wet foliage thoroughly. Repeat treatment where heavy weeds appear to resist spray. Will control weeds as listed above.

SPOT SPRAYING: Spray anytime of the year. Add 6 tablespoons per gallon water. Thoroughly wet weeds.

APPLICATION CHART

Amount of Spray	Amount of Water	To Cover
6 tbsp (3 oz.)	2 gallons	1,000 sq. ft.
30 tbsp (15 oz.)	10-15 gallons	5,000 sq. ft.