

DIRECTIONS FOR USE

Applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, harrow, or other suitable means for greatest effectiveness.

FIELD AND FORAGE INSECTS

- BEANS (Pinto)**
Lesser Cornstalk Borers: Apply 7-14 lbs. per acre in a 6-inch band centered over the row when beans are 1/2 - 1 inch high.
- CORN**
Corn Rootworm Larvae Postemergence Basal Treatment: Apply 4 1/2 - 8 1/2 oz. per 1,000 linear feet of row in a band over the row at first sign of rootworms, but not later than June 24 in the Middle Corn Belt or July 1st in the Northern Corn Belt. Cover treated band with 2-3 inches of soil. Apply amount per acre recommended according to row spacing chart for basal treatment.

| Row Spacing | Linear feet of corn row/acre | Amount of Diazinon 14 G per acre — Postemergence basal treatment |
|-------------|------------------------------|--|
| 48" | 10,890 | 3-6 lbs. |
| 42" | 12,455 | 3 1/4-6 1/2 lbs. |
| 40" | 13,068 | 3 1/2-7 lbs. |
| 38" | 13,939 | 3 3/4-7 1/2 lbs. |
| 36" | 14,520 | 4-8 lbs. |
| 30" | 17,424 | 4 1/2-9 1/2 lbs. |
| 24" | 21,780 | 4 3/4-11 1/2 lbs. |
| 20" | 26,136 | 7-14 lbs. |
| 15" | 34,848 | 9 1/4-18 1/2 lbs. |

European Corn Borer: Apply 7-14 lbs. per acre over the corn so granules fall into whorls, when 75% of plants show first-generation borers feeding. Make a second application 7 days later if needed. To control second-generation corn borers, apply 7-14 lbs. per acre, as above, when field count shows 100 egg masses per 100 plants.

Diazinon 14G may be applied by aircraft for the above use. This material should be uniformly applied to the area to be treated at the recommended rates given above.

(Combining Postemergence Treatment for Control of Corn Rootworm Larvae and European Corn Borer - North Central United States Only): When infestations of these insects coincide, a single application may be made to replace the above basal postemergence treatment for corn rootworm larvae and one of the above applications for the control of first-generation European corn borer. Use the combination treatment where European corn borer egg masses or larvae are present at the same time corn rootworm larvae should be controlled. Treatment should be made at the peak of first-generation European corn borer egg laying, but not later than June 24 in the Middle Corn Belt or July 1 in the Northern Corn Belt for adequate rootworm control.

Apply 7-14 lbs. per acre over the corn plants. Cover granules falling to the ground by throwing 2-3 inches of soil into the corn row with a disk tiller or cultivator.

Repeat application to corn whorls 7 days later if needed, for control of European corn borer. To control second-generation corn borers apply 7-14 lbs. per acre as above, when field count shows 100 egg masses per 100 plants.

Fall Armyworms: Apply 7-14 lbs. per acre over the corn so granules fall into whorls when armyworms first appear. Repeat application if necessary.

Southwestern Corn Borers: Apply 7-14 lbs. per acre over the corn so granules fall into whorls when second-generation borers first appear. Repeat application one week later.

NOTE: Do not feed treated fodder to dairy or beef cattle or sheep for 10 days following application. Corn may be picked immediately. Temporary spotting of foliage sometimes occurs following application.

Cutworms: Broadcast 14-28 lbs. per acre. For surface cutworms apply just prior to planting or when larvae first appear and work into soil 1-2 inches. Do not apply after layby. For subterranean cutworms apply just prior to planting and work into soil 3-6 inches.

Lesser Cornstalk Borers: Apply 8 1/2-17 oz. per 1,000 linear feet of row in a 10-inch band centered over the row. Make application at planting time or when crop begins to emerge. Cover the treated band with 1-2 inches of soil.

Apply amount per acre recommended according to row spacing chart for basal treatment.

Seed Corn Maggots: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2 inches.

Garden Symphylans (Centipedes): Broadcast 70 lbs. per acre just prior to planting and work into soil 4-8 inches. This pre-plant treatment will also control other soil insects, cutworms (surface and subterranean) and wireworms.

Wireworms: Broadcast 21-28 lbs. per acre before planting. Work into soil 4-8 inches.

PEANUTS

Southern Corn Rootworms: Broadcast 28 lbs. per acre or the equivalent in a band over the fruiting zone, just prior to pegging. Work into top few inches of soil.

Cutworms: Apply 14 lbs. per acre in a 12-inch band over the row when cutworms first appear. Up to three additional applications at three-week intervals may be necessary. Do not apply within 21 days of peanut harvest.

Lesser Cornstalk Borers: Apply 14-21 lbs. per acre in a band over the fruiting zone just prior to pegging. Cover the band with 1-2 inches of soil.

NOTE: LIVESTOCK (1) may be fed peanut hulls (2) may be fed peanut forage 7 days following treatment and (3) may be fed peanut hay 21 days following treatment.

SORGHUM

Cutworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

Lesser Cornstalk Borers: Apply 7-14 lbs. per acre in a 10-inch band over the row at planting time or as crop emerges. Cover the band with 1-2 inches of soil.

NOTE: Do not apply directly to the seed furrow as reduction of sorghum seed germination may occur.

SOYBEANS

Cutworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-5 inches to control subterranean cutworms.

Lesser Cornstalk Borers: Apply 7-14 lbs. per acre in a 6-inch band over the row just prior to planting or when soybeans are 1-2 inches high. Cover the band with 1-2 inches of soil.

SMALL FRUIT INSECTS

STRAWBERRIES

Mole Crickets: Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to transplanting and work into soil 1-2 inches.

SUGAR BEETS

Sugar Beet Root Maggot: Apply 7-14 lbs. per acre in a 5-7 inch band (based on row or 4.7-9.4 oz. per 1,000 linear feet of row in a 5-7 inch band either incorporated planting or applying above the seed after the seed is covered. Apply ahead of the heavy maggot infestation is anticipated or a wet spring occurs, a second application early postemergence, when the adult maggots (flies) first start to appear. If a second application is necessary, apply 7-10 1/2 lbs. per acre in a 5-7 inch band (based on row width) of 22 oz. per 1,000 linear feet of row in a 5-7 inch band directly over the row.

NOTE: Diazinon should not come in direct contact with the seed or injury will result.

Wireworms: Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-8 inches.

SUGARCANE

Sugarcane Borers: Broadcast 10 1/2-17 1/2 lbs. per acre after joints have begun to show at least 5% of plants show borers feeding in or under the leaf sheath but which have not stalked. Additional applications may be necessary to assure continuing control.

NOTE: Do not make more than four applications per growing season. Allow 28 days between application and harvest. At that time bagasse and forage may be fed to livestock.

Wireworms: Apply 42 lbs. per acre in a 12-inch band over the open furrow, directly over the row.

TOBACCO

Cutworms: Broadcast 14-21 lbs. per acre before transplanting. Work into soil 1-2 inches.

Mole Crickets: Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

Seed Corn Maggots: Broadcast 10 1/2 lbs. per acre 3-4 weeks before transplanting.

Wireworms: Broadcast 14-21 lbs. per acre 3-4 weeks before transplanting. Work into soil 4-8 inches.

FORAGE CROPS (ALFALFA, CLOVER, COWPEAS, AND LESPEDEZA)

Cutworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER

Root Maggots: Broadcast 14-21 lbs. per acre before planting or transplanting. Work into soil 4-8 inches.

ONIONS

Onion Maggots: Apply 14-28 lbs. per acre as broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing, or other means.

RADISHES

Root Maggots: Apply 10 1/2-14 lbs. per acre (2-1/2- 2-4/5 oz. per 1,000 linear feet of row) at planting time.

SWEET CORN

See Corn under Field and Forage Crops.

TOMATOES

Vinegar Flies (Drosophila spp.): Broadcast 7 lbs. per acre when flies first appear. Repeat application at 7-day intervals or as required.

OTHER VEGETABLE INSECTS

CUTWORMS (SURFACE AND SUBTERRANEAN)

| | | | |
|------------------|-------------------|------------|---------------|
| Broccoli | Collards | Muskmelons | Spinach |
| Brussels Sprouts | Cucumbers | Parsley | Summer Squash |
| Cabbage | Endive (escarole) | Peas | Swiss Chard |
| Cantaloupes | Kale | Peppers | Tomatoes |
| Cauliflower | Lettuce | Pole Beans | Watermelon |
| Celery | Lima Beans | Snap Beans | Winter Squash |

Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

Potatoes: Broadcast 14-28 lbs. per acre before planting. Work into soil 4-6 inches.

GARDEN SYMPHYLANS (CENTIPEDES)

| | | | |
|---------|------------|------------|----------|
| Cabbage | Peas | Red Beets | Tomatoes |
| Carrots | Pole Beans | Snap Beans | Turnips |
| Lettuce | Radishes | | |

Broadcast 70 lbs. per acre just prior to planting. Work into soil 4-8 inches. This pre-plant treatment will also control other soil insects, cutworms (surface and subterranean), mole crickets, and wireworms.

MOLE CRICKETS

| | | | |
|------------------|-------------------|---------|----------|
| Broccoli | Collards | Lettuce | Radishes |
| Brussels Sprouts | Endive (escarole) | Muscard | Tomatoes |
| Cabbage | Kale | Peppers | Turnips |
| Cauliflower | | | |

Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

LESSER CORNSTALK BORERS

Beans (Lima, Pole, Snap): Apply 7-14 lbs. per acre in a 6-inch band centered over the row when plants are 1/2-1 inch tall.

WIREWORMS

| | | | |
|------------------|-------------------|------------|---------------|
| Broccoli | Collards | Onions | Summer Squash |
| Brussels Sprouts | Cucumbers | Parsley | Swiss Chard |
| Cabbage | Endive (escarole) | Peas | Tomatoes |
| Cantaloupes | Kale | Peppers | Turnips |
| Carrots | Lettuce | Pole Beans | Watermelon |
| Cauliflower | Lima Beans | Snap Beans | Winter Squash |
| Celery | Muskmelons | Spinach | |


Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-8 inches.

Potatoes: Broadcast 21-42 lbs. per acre before planting. Work into soil 4-8 inches.

per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of the row.

Sweet Potatoes (Southeastern U.S.): Broadcast 21 lbs. per acre over the top of the row when plants are 1-2 inches high.

BEST DOCUMENT AVAILABLE



NET WEIGHT 50 POUNDS
EPA Reg. No. 2169-265 E.P.A. Est. 2169-KS-1

Patterson GREEN-UP Company
Div. Curry Cartwright, Inc.
1400 Union Ave., Kansas City, Mo. 64101

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

ould be worked into the top few inches immediately after application with a
k. harrow, or other suitable means for greatest effectiveness.

FIELD AND FORAGE INSECTS

Apply 7-14 lbs. per acre in a 6-inch band centered over the row when
n.

Postemergence Basal Treatment: Apply 4½ - 8½ oz. per 1,000 linear
feet of row at first sign of rootworms, but not later than June 24 in the Middle
and Northern Corn Belt. Cover treated band with 2-3 inches of soil.
Recommended according to row spacing chart for basal treatment.

| Linear feet of corn row/acre | Amount of Diazinon 14 G per acre — Postemergence basal treatment |
|---------------------------------|---|
| 10,890 | 3-6 lbs. |
| 12,455 | 3¼-6½ lbs. |
| 13,068 | 3½-7 lbs. |
| 13,939 | 3¾-7½ lbs. |
| 14,520 | 4-8 lbs. |
| 17,424 | 4¾-9½ lbs. |
| 21,780 | 4¾-11½ lbs. |
| 26,136 | 7-14 lbs. |
| 34,848 | 9½-18½ lbs. |

Apply 7-14 lbs. per acre over the corn so granules fall into whorls, when
first-generation borers feeding. Make a second application 7 days later if
second-generation corn borers, apply 7-14 lbs. per acre, as above, when field
count shows 100 plants.

Applied by aircraft for the above use. This material should be uniformly applied
at the recommended rates given above.

**Pre-plant Treatment for Control of Corn Rootworm Larvae and European Corn
Borer (United States Only):** When infestations of these insects coincide, a single
application to replace the above basal postemergence treatment for corn rootworm
larvae applications for the control of first-generation European corn borer. Use
only where European corn borer egg masses or larvae are present at the
time larvae should be controlled. Treatment should be applied at the peak of
European corn borer egg laying, but not later than June 24 in the Middle Corn Belt or
Northern Corn Belt for adequate rootworm control.

Apply over the corn plants. Cover granules falling to the ground by throwing 2-3
inches of soil over the corn row with a disk harrow or cultivator.

Apply over corn whorls 7 days later if needed, for control of European corn borer. To
control corn borers apply 7-14 lbs. per acre as above, when field count shows
100 plants.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
first-generation borers first appear. Repeat application if necessary.

Borers: Apply 7-14 lbs. per acre over the corn so granules fall into whorls
when borers first appear. Repeat application one week later.

Apply over fodder to dairy or beef cattle or sheep for 10 days following application.
Temporary spotting of foliage sometimes occurs following

Apply 14-28 lbs. per acre. For surface cutworms apply just prior to planting or when
cutworms first appear. Work into soil 1-2 inches. Do not apply after planting. For subterranean
cutworms apply just prior to planting and work into soil 3-6 inches.

Borers: Apply 8½-17 oz. per 1,000 linear feet of row in a 10-inch band centered
over the row at planting time or when crop begins to emerge. Cover the treated
band with 2-3 inches of soil.

Recommended according to row spacing chart for basal treatment.

Broccoli: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2 inches.

(Centipedes): Broadcast 70 lbs. per acre just prior to planting and work into
soil 4-6 inches. This pre-plant treatment will also control other soil insects, cutworms (surface and
subterranean), mole crickets, root maggots, and wireworms.

Wireworms: Broadcast 21-28 lbs. per acre before planting. Work into soil 4-6 inches.

Wireworms: Broadcast 28 lbs. per acre or the equivalent in a band over the fruiting
zone. Work into top few inches of soil.

Wireworms: Apply 7-14 lbs. per acre in a 12-inch band over the row when cutworms first appear. Up to
three applications at three-week intervals may be necessary. Do not apply within 21 days
of planting.

Borers: Apply 14-21 lbs. per acre in a band over the fruiting zone just prior to
planting. Work into soil 1-2 inches.

(1) may be fed peanut hulls; (2) may be fed peanut forage 7 days following
application; (3) may be fed peanut hay 21 days following treatment.

Wireworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control
subterranean cutworms.

Borers: Apply 7-14 lbs. per acre in a 10-inch band over the row at planting time or
when crop begins to emerge. Cover the band with 1-2 inches of soil.

Apply directly to the seed furrow as reduction of sorghum seed germination may
occur.

Wireworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control
subterranean cutworms.

Borers: Apply 7-14 lbs. per acre in a 6-inch band over the row just prior to
planting. Work into soil 1-2 inches.

Wireworms: Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Wireworms: Broadcast 21-42 lbs. per acre before planting. Work into soil 4-8 inches or apply 7-21 lbs.
per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of row.

Sweet Potatoes (Southeastern U.S.): Broadcast 21 lbs. per acre over the tops of vines when roots
begin to enlarge.

SUGAR BEETS

Sugar Beet Root Maggot: Apply 7-14 lbs. per acre in a 5-7 inch band (based on row width of 22 ins.)
or 4.7-9.4 oz. per 1,000 linear feet of row in a 5-7 inch band either incorporating 2-3 inches at
planting or applying above the seed after the seed is covered. Apply ahead of the press wheel. If a
heavy maggot infestation is anticipated or a wet spring occurs, a second application may be made,
early postemergence, when the adult maggots (flies) first start to appear. If a second application is
necessary, apply 7-10½ lbs. per acre in a 5-7 inch band (based on row width of 22 inches) or 4.7-7.1
oz. per 1,000 linear feet of row in a 5-7 inch band directly over the row.

NOTE: Diazinon should not come in direct contact with the seed or injury will occur.

Wireworms: Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-8 inches.

SUGARCANE

Sugarcane Borers: Broadcast 10½-17½ lbs. per acre after joints have begun to form and when at
least 5% of plants show borers feeding in or under the leaf sheath but which have not bored into the
stalks. Additional applications may be necessary to assure continuing control.

NOTE: Do not make more than four applications per growing season. Allow 28 days between last
application and harvest. At that time bagasse and forage may be fed to livestock.

Wireworms: Apply 42 lbs. per acre in a 12-inch band over the open furrow, directly to seed pieces.

TOBACCO

Cutworms: Broadcast 14-21 lbs. per acre before transplanting. Work into soil 2-4 inches.

Mole Crickets: Apply 7 lbs. per acre when insects are present. Make application to plant beds or
fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

Seed Corn Maggots: Broadcast 10½ lbs. per acre 3-4 weeks before transplanting. Work into soil 2
inches.

Wireworms: Broadcast 14-21 lbs. per acre 3-4 weeks before transplanting. Work into soil 6-9
inches.

FORAGE CROPS (ALFALFA, CLOVER, COWPEAS, AND LESPEDEZA)

Cutworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control
surface cutworms; 3-6 inches to control subterranean cutworms.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER

Root Maggots: Broadcast 14-21 lbs. per acre before planting or transplanting. Work into soil 3-4
inches.

ONIONS

Onion Maggots: Apply 14-28 lbs. per acre as broadcast soil treatment just prior to planting.
Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing or other suitable
means.

RADISHES

Root Maggots: Apply 10½-14 lbs. per acre (2-1/2 - 2-4/5 oz. per 1,000 linear feet of row) in the
seed furrow at planting time.

SWEET CORN

See Corn under Field and Forage Crops.

TOMATOES

Vinegar Flies (Drosophila spp): Broadcast 7 lbs. per acre when flies first appear. Repeat in 7-10
day intervals or as required.

OTHER VEGETABLE INSECTS

CUTWORMS (SURFACE AND SUBTERRANEAN)

| | | | |
|------------------|-------------------|------------|---------------|
| Broccoli | Collards | Muskmelons | Spinach |
| Brussels Sprouts | Cucumbers | Parsley | Summer Squash |
| Cabbage | Endive (escarole) | Peas | Swiss Chard |
| Cantaloupes | Kale | Peppers | Tomatoes |
| Cauliflower | Lettuce | Pole Beans | Watermelons |
| Celery | Lima Beans | Snap Beans | Winter Squash |

Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface
cutworms; 3-6 inches to control subterranean cutworms.

Potatoes: Broadcast 14-28 lbs. per acre before planting. Work into soil 4-6 inches.

GARDEN SYMPHYLANS (CENTIPEDES)

| | | | |
|---------|------------|------------|----------|
| Cabbage | Peas | Red Beets | Tomatoes |
| Carrots | Pole Beans | Snap Beans | Turnips |
| Lettuce | Radishes | | |

Broadcast 70 lbs. per acre just prior to planting. Work into soil 4-8 inches. This pre-plant treatment will
also control other soil insects, cutworms (surface and subterranean), mole crickets, root maggots,
and wireworms.

MOLE CRICKETS

| | | | |
|------------------|-------------------|---------|----------|
| Broccoli | Collards | Lettuce | Radishes |
| Brussels Sprouts | Endive (escarole) | Mustard | Tomatoes |
| Cabbage | Kale | Peppers | Turnips |
| Cauliflower | | | |

Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior
to sowing of seed or transplanting and work into soil 1-2 inches.

LESSER CORNSTALK BORERS

Beans (Lima, Pole, Snap): Apply 7-14 lbs. per acre in a 6-inch band centered over the row. Apply
when plants are ½-1 inch tall.

WIREWORMS

| | | | |
|------------------|-------------------|------------|----------------|
| Broccoli | Collards | Onions | Summer Squash |
| Brussels Sprouts | Cucumbers | Parsley | Sweet Potatoes |
| Cabbage | Endive (escarole) | Peas | Swiss Chard |
| Cantaloupes | Kale | Peppers | Tomatoes |
| Carrots | Lettuce | Pole Beans | Watermelons |
| Cauliflower | Lima Beans | Snap Beans | Winter Squash |
| Celery | Muskmelons | Spinach | |

Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Potatoes: Broadcast 21-42 lbs. per acre before planting. Work into soil 4-8 inches or apply 7-21 lbs.
per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of row.

Sweet Potatoes (Southeastern U.S.): Broadcast 21 lbs. per acre over the tops of vines when roots
begin to enlarge.

BEST DOCUMENT AVAILABLE

23A

23

BEST DOCUMENT AVAILABLE

PTED

NET WEIGHT 50 POUNDS

EPA Reg. No. 2169-265 E.P.A. Est. 2169-KS-1

Patterson GREEN-UP Company

Div. Curry Cartwright, Inc.

1400 Union Ave., Kansas City, Mo. 64101

23A

FIELD AND FORAGE INSECTS

- BEANS (Pinto)**
Lesser Cornstalk Borers: Apply 7-14 lbs. per acre in a 6-inch band centered over the row when beans are 1/2 - 1 inch high.
- CORN**
Corn Rootworm Larvae Postemergence Basal Treatment: Apply 4 1/2 - 8 1/2 oz. per 1,000 linear feet of row in a band over the row at first sign of rootworms, but not later than June 24 in the Middle Corn Belt or July 1st in the Northern Corn Belt. Cover treated band with 2-3 inches of soil. Apply amount per acre recommended according to row spacing chart for basal treatment.

| Row Spacing | Linear feet of corn row/acre | Amount of Diazinon 14 G per acre — Postemergence basal treatment |
|-------------|------------------------------|--|
| 48" | 10,890 | 3-6 lbs. |
| 42" | 12,455 | 3 1/4 - 6 1/2 lbs. |
| 40" | 13,068 | 3 1/2 - 7 lbs. |
| 38" | 13,939 | 3 3/4 - 7 1/2 lbs. |
| 36" | 14,520 | 4-8 lbs. |
| 30" | 17,424 | 4 3/4 - 9 1/2 lbs. |
| 24" | 21,780 | 4 1/2 - 11 1/2 lbs. |
| 20" | 26,136 | 7-14 lbs. |
| 15" | 34,848 | 9 1/2 - 18 1/2 lbs. |

BEST DOCUMENT AVAILABLE

European Corn Borer: Apply 7-14 lbs. per acre over the corn so granules fall into whorls, when 75% of plants show first-generation borers feeding. Make a second application 7 days later if needed. To control second-generation corn borers, apply 7-14 lbs. per acre, as above, when field count shows 100 egg masses per 100 plants. Diazinon 14G may be applied by aircraft for the above use. This material should be uniformly applied to the area to be treated at the recommended rates given above. (Combining Postemergence Treatment for Control of Corn Rootworm Larvae and European Corn Borer - North Central United States Only): When infestations of these insects coincide, a single application may be made to replace the above basal postemergence treatment for corn rootworm larvae and one of the above applications for the control of first-generation European corn borer. Use the combination treatment where European corn borer egg masses or larvae are present at the same time corn rootworm larvae should be controlled. Treatment should be made at the peak of first-generation European corn borer egg laying, but not later than June 24 in the Middle Corn Belt or July 1 in the Northern Corn Belt for adequate rootworm control.

Apply 7-14 lbs. per acre over the corn plants. Cover granules falling to the ground by throwing 2-3 inches of soil into the corn row with a disk tiller or cultivator. Repeat application to corn whorls 7 days later if needed, for control of European corn borer. To control second-generation corn borers apply 7-14 lbs. per acre as above, when field count shows 100 egg masses per 100 plants.

Fall Armyworms: Apply 7-14 lbs. per acre over the corn so granules fall into whorls when armyworms first appear. Repeat application if necessary.

Southwestern Corn Borers: Apply 7-14 lbs. per acre over the corn so granules fall into whorls when second-generation borers first appear. Repeat application one week later.

NOTE: Do not feed treated fodder to dairy or beef cattle or sheep for 10 days following application. Corn may be picked immediately. Temporary spotting of foliage sometimes occurs following application.

Cutworms: Broadcast 14-28 lbs. per acre. For surface cutworms apply just prior to planting or when larvae first appear and work into soil 1-2 inches. Do not apply after layby. For subterranean cutworms apply just prior to planting and work into soil 3-6 inches.

Lesser Cornstalk Borers: Apply 8 1/2 - 17 oz. per 1,000 linear feet of row in a 10-inch band centered over the row. Make application at planting time or when crop begins to emerge. Cover the treated band with 1-2 inches of soil.

Apply amount per acre recommended according to row spacing chart for basal treatment.

Seed Corn Maggots: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2 inches.

Garden Symphylans (Centipedes): Broadcast 70 lbs. per acre just prior to planting and work into soil 4-8 inches. This pre-plant treatment will also control other soil insects, cutworms (surface and subterranean) and wireworms.

Wireworms: Broadcast 21-28 lbs. per acre before planting. Work into soil 4-6 inches.

PEANUTS
Southern Corn Rootworms: Broadcast 28 lbs. per acre or the equivalent in a band over the fruiting zone, just prior to pegging. Work into top few inches of soil.

Cutworms: Apply 14 lbs. per acre in a 12-inch band over the row when cutworms first appear. Up to three additional applications at three-week intervals may be necessary. Do not apply within 21 days of peanut harvest.

Lesser Cornstalk Borers: Apply 14-21 lbs. per acre in a band over the fruiting zone just prior to pegging. Cover the band with 1-2 inches of soil.

NOTE: LIVESTOCK (1) may be fed peanut hulls (2) may be fed peanut forage 7 days following treatment and (3) may be fed peanut hay 21 days following treatment.

SORGHUM
Cutworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

Lesser Cornstalk Borers: Apply 7-14 lbs. per acre in a 10-inch band over the row at planting time or as crop emerges. Cover the band with 1-2 inches of soil.

NOTE: Do not apply directly to the seed furrow as reduction of sorghum seed germination may occur.

SOYBEANS
Cutworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

Lesser Cornstalk Borers: Apply 7-14 lbs. per acre in a 6-inch band over the row just prior to planting or when soybeans are 1-2 inches high. Cover the band with 1-2 inches of soil.

SMALL FRUIT INSECTS

STRAWBERRIES
Mole Crickets: Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to transplanting and work into soil 1-2 inches.

or 4 7-9 4 oz. per 1,000 linear feet of row in a 5-7 inch band directly over the row. Apply ahead of the planting or applying above the seed after the seed is covered. Apply ahead of the planting or applying above the seed after the seed is covered. Apply ahead of the planting or applying above the seed after the seed is covered. Apply ahead of the planting or applying above the seed after the seed is covered. Apply ahead of the planting or applying above the seed after the seed is covered.

NOTE: Diazinon should not come in direct contact with the seed or injury will occur. **Wireworms:** Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

SUGARCANE
Sugarcane Borers: Broadcast 10 1/2 - 17 1/2 lbs. per acre after joints have begun to form and at least 5% of plants show borers feeding in or under the leaf sheath but which have not yet killed the stalks. Additional applications may be necessary to assure continuing control.

NOTE: Do not make more than four applications per growing season. Allow 28 days between applications and harvest. At that time bagasse and forage may be fed to livestock.

Wireworms: Apply 42 lbs. per acre in a 12-inch band over the open furrow, directly over the seed.

TOBACCO
Cutworms: Broadcast 14-21 lbs. per acre before transplanting. Work into soil 2-4 inches.

Mole Crickets: Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

Seed Corn Maggots: Broadcast 10 1/2 lbs. per acre 3-4 weeks before transplanting and work into soil 2 inches.

Wireworms: Broadcast 14-21 lbs. per acre 3-4 weeks before transplanting. Work into soil 4-6 inches.

FORAGE CROPS (ALFALFA, CLOVER, COWPEAS, AND LESPEDEZA)
Cutworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER
Root Maggots: Broadcast 14-21 lbs. per acre before planting or transplanting. Work into soil 2-3 inches.

ONIONS
Onion Maggots: Apply 14-28 lbs. per acre as broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing or similar means.

RADISHES
Root Maggots: Apply 10 1/2 - 14 lbs. per acre (2-1/2 - 2-4/5 oz. per 1,000 linear feet of row) at seed furrow at planting time.

SWEET CORN
See Corn under Field and Forage Crops.

TOMATOES
Vinegar Flies (Drosophila spp): Broadcast 7 lbs. per acre when flies first appear. Repeat application at 7-day intervals or as required.

OTHER VEGETABLE INSECTS

| CUTWORMS (SURFACE AND SUBTERRANEAN) | | | |
|--|-------------------|------------|---------------|
| Broccoli | Collards | Muskmelons | Spinach |
| Brussels Sprouts | Cucumbers | Parsley | Summer Squash |
| Cabbage | Endive (escarole) | Peas | Swiss Chard |
| Cantaloupes | Kale | Peppers | Tomatoes |
| Cauliflower | Lettuce | Pole Beans | Watermelon |
| Celery | Lima Beans | Snap Beans | Winter Squash |

Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

Potatoes: Broadcast 14-28 lbs. per acre before planting. Work into soil 4-6 inches.

GARDEN SYMPHYLANS (CENTIPEDES)
Cabbage Peas Red Beets Tomatoes
Carrots Pole Beans Snap Beans Turnips
Lettuce Radishes

Broadcast 70 lbs. per acre just prior to planting. Work into soil 4-8 inches. This pre-plant treatment will also control other soil insects, cutworms (surface and subterranean), mole crickets and wireworms.

| MOLE CRICKETS | | | |
|----------------------|-------------------|---------|----------|
| Broccoli | Collards | Lettuce | Radishes |
| Brussels Sprouts | Endive (escarole) | Mustard | Tomatoes |
| Cabbage | Kale | Peppers | Turnips |
| Cauliflower | | | |

Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.


LESSER CORNSTALK BORERS
Beans (Lima, Pole, Snap): Apply 7-14 lbs. per acre in a 6-inch band centered over the row when plants are 1/2-1 inch tall.

| WIREWORMS | | | |
|------------------|-------------------|------------|----------------|
| Broccoli | Collards | Onions | Summer Squash |
| Brussels Sprouts | Cucumbers | Parsley | Sweet Potatoes |
| Cabbage | Endive (escarole) | Peas | Swiss Chard |
| Cantaloupes | Kale | Peppers | Tomatoes |
| Carrots | Lettuce | Pole Beans | Watermelon |
| Cauliflower | Lima Beans | Snap Beans | Winter Squash |
| Celery | Muskmelons | Spinach | |

Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-8 inches.

Potatoes: Broadcast 21-42 lbs. per acre before planting. Work into soil 4-8 inches per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of the row.

Sweet Potatoes (Southeastern U.S.): Broadcast 21 lbs. per acre over the tops of the plants when they begin to enlarge.



NET WEIGHT 50 POUNDS
EPA Reg. No. 2169-265 E.P.A. Est. 2169-KS-1
Patterson GREEN-UP Company
Div. Curry Cartwright, Inc.
1400 Union Ave., Kansas City, Mo. 64101

BEST DOCUMENT AVAILABLE

ould be worked into the top few inches immediately after application with a
 disk, harrow, or other suitable means for greatest effectiveness.

FIELD AND FORAGE INSECTS

Apply 7-14 lbs. per acre in a 6-inch band centered over the row when
 ph.

Postemergence Basal Treatment: Apply 4½ - 8½ oz. per 1,000 linear
 feet of row at first sign of rootworms, but not later than June 24 in the Middle
 the Northern Corn Belt. Cover treated band with 2-3 inches of soil.
 recommended according to row spacing chart for basal treatment.

| Linear feet of corn row/acre | Amount of Diazinon 14 G per acre — Postemergence basal treatment |
|---------------------------------|---|
| 10,890 | 3-6 lbs. |
| 12,455 | 3¼-6½ lbs. |
| 13,068 | 3½-7 lbs. |
| 13,939 | 3¾-7½ lbs. |
| 14,520 | 4-8 lbs. |
| 17,424 | 4¾-9½ lbs. |
| 21,780 | 4¾-11½ lbs. |
| 25,136 | 7-14 lbs. |
| 34,848 | 9¾-18½ lbs. |

Apply 7-14 lbs. per acre over the corn so granules fall into whorls, when
 first-generation borers feeding. Make 2. second application 7 days later if
 second-generation corn borers, apply 7-14 lbs. per acre, as above, when field
 counts per 100 plants.

Applied by aircraft for the above use. This material should be uniformly applied
 at the recommended rates given above.

**Pre-planting Treatment for Control of Corn Rootworm Larvae and European Corn
 Borer (United States Only):** When infestations of these insects coincide, a single
 application to replace the above basal postemergence treatment for corn rootworm
 larvae and European corn borer. Use applications for the control of first-generation European corn borer. Use
 only where European corn borer egg masses or larvae are present at the
 time larvae should be controlled. Treatment should be made at the peak of
 European corn borer egg laying, but not later than June 24 in the Middle Corn Belt or
 Corn Belt for adequate rootworm control.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borer first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

Apply 7-14 lbs. per acre over the corn so granules fall into whorls when
 European corn borers first appear. Repeat application one week later.

SUGAR BEETS

Sugar Beet Root Maggot: Apply 7-14 lbs. per acre in a 5-7 inch band (based on row width of 22 ins.)
 or 4.7-9.4 oz. per 1,000 linear feet of row in a 5-7 inch band either incorporating 2-3 inches of
 planting or applying above the seed after the seed is covered. Apply ahead of the press wheel. If a
 heavy maggot infestation is anticipated or a wet spring occurs, a second application may be made,
 early postemergence, when the adult maggots (flies) first start to appear. If a second application is
 necessary, apply 7-10½ lbs. per acre in a 5-7 inch band (based on row width of 22 inches) or 4.7-7.1
 oz. per 1,000 linear feet of row in a 5-7 inch band directly over the row.

NOTE: Diazinon should not come in direct contact with the seed or injury will occur.

Wireworms: Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-6 inches

SUGARCANE

Sugarcane Borers: Broadcast 10½-17½ lbs. per acre after joints have begun to form and when at
 least 5% of plants show borers feeding in or under the leaf sheath but which have not bored into the
 stalks. Additional applications may be necessary to assure continuing control.

NOTE: Do not make more than four applications per growing season. Allow 28 days between last
 application and harvest. At that time bagasse and forage may be fed to livestock.

Wireworms: Apply 42 lbs. per acre in a 12-inch band over the open furrow, directly to seed pieces.

TOBACCO

Cutworms: Broadcast 14-21 lbs. per acre before transplanting. Work into soil 2-4 inches.

Mole Crickets: Apply 7 lbs. per acre when insects are present. Make application to plant beds or
 fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

Seed Corn Maggots: Broadcast 10½ lbs. per acre 3-4 weeks before transplanting. Work into soil 2
 inches.

Wireworms: Broadcast 14-21 lbs. per acre 3-4 weeks before transplanting. Work into soil 4-6
 inches.

FORAGE CROPS (ALFALFA, CLOVER, COWPEAS, AND LESPEDEZA)

Cutworms: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control
 surface cutworms; 3-6 inches to control subterranean cutworms.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER

Root Maggots: Broadcast 14-21 lbs. per acre before planting or transplanting. Work into soil 3-4
 inches

ONIONS

Onion Maggots: Apply 14-28 lbs. per acre as broadcast soil treatment just prior to planting.
 Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing or other suitable
 means.

RADISHES

Root Maggots: Apply 10½-14 lbs. per acre (2-1/2- 2-4.5 oz. per 1,000 linear feet of row) in the
 seed furrow at planting time.

SWEET CORN

See Corn under Field and Forage Crops.

TOMATOES

Vinegar Flies (Drosophila spp): Broadcast 7 lbs. per acre when flies first appear. Repeat in 7-10
 day intervals or as required.

OTHER VEGETABLE INSECTS

CUTWORMS (SURFACE AND SUBTERRANEAN)

| | | | |
|------------------|-------------------|------------|---------------|
| Broccoli | Collards | Muskmelons | Spinach |
| Brussels Sprouts | Cucumbers | Parsley | Summer Squash |
| Cabbage | Endive (escarole) | Peas | Swiss Chard |
| Cantaloupes | Kale | Peppers | Tomatoes |
| Cauliflower | Lettuce | Pole Beans | Watermelons |
| Celery | Lima Beans | Snap Beans | Winter Squash |

Broadcast 14-26 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface
 cutworms; 3-6 inches to control subterranean cut worms.

Potatoes: Broadcast 14-26 lbs. per acre before planting. Work into soil 4-6 inches.

GARDEN SYMPHYLANS (CENTIPEDES)

| | | | |
|---------|------------|------------|----------|
| Cabbage | Peas | Red Beets | Tomatoes |
| Carrots | Pole Beans | Snap Beans | Turnips |
| Lettuce | Radishes | | |

Broadcast 70 lbs. per acre just prior to planting. Work into soil 4-8 inches. This preplant treatment will
 also control other soil insects, cutworms (surface and subterranean), mole crickets, root maggots,
 and wireworms.

MOLE CRICKETS

| | | | |
|------------------|-------------------|---------|----------|
| Broccoli | Collards | Lettuce | Radishes |
| Brussels Sprouts | Endive (escarole) | Mustard | Tomatoes |
| Cabbage | Kale | Peppers | Turnips |
| Cauliflower | | | |

Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior
 to sowing of seed or transplanting and work into soil 1-2 inches.

LESSER CORNSTALK BORERS

Beans (Lima, Pole, Snap): Apply 7-14 lbs. per acre in a 6-inch band centered over the row. Apply
 when plants are ½-1 inch tall.

WIREWORMS

| | | | |
|------------------|-------------------|------------|----------------|
| Broccoli | Collards | Onions | Summer Squash |
| Brussels Sprouts | Cucumbers | Parsley | Sweet Potatoes |
| Cabbage | Endive (escarole) | Peas | Swiss Chard |
| Cantaloupes | Kale | Peppers | Tomatoes |
| Carrots | Lettuce | Pole Beans | Watermelons |
| Cauliflower | Lima Beans | Snap Beans | Winter Squash |
| Celery | Muskmelons | Spinach | |

Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-8 inches.

Potatoes: Broadcast 21-42 lbs. per acre before planting, work into soil 4-8 inches or apply 7-21 lbs.
 per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of row.

Sweet Potatoes (Southeastern U.S.): Broadcast 21 lbs. per acre over the tops of vines when roots
 begin to enlarge.

BEST DOCUMENT AVAILABLE

23A

BEST DOCUMENT AVAILABLE

24A


NET
 69-265

NET WEIGHT 50 POUNDS

EPA Reg. No. 2169-265 E.P.A. Est. 2169-KS-1

Patterson GREEN-UP Company
 Div. Curry Cartwright, Inc.
 1400 Union Ave., Kansas City, Mo. 64101

24



DIAZINON® 14G

Granular Insecticide

For control of soil insects, corn rootworms, wireworms, cutworms, and other pests as listed.

ACTIVE INGREDIENT:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothionate

14.0%

INERT INGREDIENTS:

86.0%

TOTAL 100.0%

*Diazinon Trademark of Ciba-Geigy
EPA Reg. No. 2169-265

EPA Est. No. 2169-KS-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person. **If inhaled** — Remove victim to fresh air. Apply respiration if indicated. **If on skin** — Wash contaminated skin with soap and water. **If in eyes** — Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

BEST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. May be absorbed through skin.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp, crabs, birds, and other wildlife. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when...

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

LAWN INSECTS

Sod Webworms — Remove thatch. Apply 1/2-1 lb. Diazinon per 1,000 sq. ft. Apply evenly with a drop type or rotary sprayer. Treat when insects first appear. Water grass thoroughly after application. Keep pets and children off treated area until...

Lawn Chinch Bugs — Apply 1/2-1 lb. per 1,000 sq. ft. (2 1/2 lbs. per 5,000 sq. ft.) and brush lawn to work granules down to surface. Water lawn thoroughly after application. Repeat application in 7-10 days if necessary. For longer residual control on dense growing turf such as St. Augustine grass, use 1 1/2 lbs. per 1,000 sq. ft. (5-7 1/2 lbs. per 5,000 sq. ft.).

Billbugs and White Grubs, Japanese Beetle Larvae, European Chafer and Southern Chafer — Apply 1 lb. per 1,000 sq. ft. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass is present, indicating damage from insect.

Hyperodes Weevil (New York Only) — Apply 3/4 lb. per 1,000 sq. ft. of grass. Water grass thoroughly after application.

09A
314

BEST DOCUMENT AVAILABLE

21

ACCEPTED
2169-265



000000
000000

greenUP

DIAZINON® 14G

Granular Insecticide

BEST DOCUMENT AVAILABLE

For control of soil insects, corn rootworms, wireworms, cutworms, and other pests as listed.

| | |
|--|--------|
| ACTIVE INGREDIENT: | |
| O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate | 14.0% |
| INERT INGREDIENTS: | 86.0% |
| TOTAL | 100.0% |

*Diazinon Trademark of Ciba-Geigy
EPA Reg. No. 2169-265

EPA Est. No. 2169-KS-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person. **If inhaled** — Remove victim to fresh air. Apply respiration if indicated. **If on skin** — Wash contaminated skin with soap and water. **If in eyes** — Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

PRECAUTIONARY STATEMENTS TO HUMANS AND DOMESTIC ANIMALS

21

CAUTION

Allowed. Avoid contamination of food and feed. Do not eat food or feed products. Avoid inhalation of dust with skin, eyes, and clothing. Wash thoroughly after use to prevent absorption through skin.

ENVIRONMENTAL HAZARDS

Is toxic to fish, shrimp, crabs, birds, and other wildlife. Avoid application to lakes, streams, ponds, tidal marshes, and estuaries. Do not apply when conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Crab may be killed at application rates recommended. Do not apply where these are important resources. Use only as specified on this label.

DISPOSAL

Use empty container. Wrap container and put in trash.

NOTICE OF WARRANTY

No warranty, expressed or implied, concerning the product other than indicated on the label. Buyer assumes risk of use and handling of this material when such is required.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. 2/A

LAWN INSECTS

Sod Webworms — Remove thatch. Apply 1/2-1 lb. Diazinon 14-G per 1,000 sq. ft. Apply evenly with a drop type or rotary spreader. Treat when insects first appear. Water grass thoroughly after application. Keep pets and children off treated area until dry.

Lawn Chinch Bugs — Apply 1/2-1 lb. per 1,000 sq. ft. (2 1/2-5 lbs. per 5,000 sq. ft.) and brush lawn to work granules down to the soil surface. Water lawn thoroughly after application. Repeat application in 7-10 days if necessary. For longer residual control and on dense growing turf such as St. Augustine grass, use 1 1/2 lbs. per 1,000 sq. ft. (5-7 1/2 lbs. per 5,000 sq. ft.).

Billbugs and White Grubs, Japanese Beetle Larvae, European Chafer and Southern Chafer — Apply 1 lb. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.

Hyperodes Weevil (New York Only) — Apply 3/4 lb. per 1,000 sq. ft. of grass. Water grass thoroughly after application. Treat when weevils are first observed.

Granular Insecticide

For control of soil insects, corn rootworms, wireworms, cutworms, and other pests as listed.

BEST DOCUMENT AVAILABLE

ACTIVE INGREDIENT:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate

14.0%

INERT INGREDIENTS:

86.0%

TOTAL 100.0%

*Diazinon Trademark of Ciba-Geigy
EPA Reg No. 2169-265

EPA Est. No. 2169-KS-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person. If inhaled — Remove victim to fresh air. Apply respiration if indicated. If on skin — Wash contaminated skin with soap and water. If in eyes — Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. May be absorbed through skin.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp, crabs, birds, and other wildlife. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product inconsistent with its labeling

LAWN INSECTS

Sod Webworms — Remove thatch. Apply 1/2-1 lb. D per 1,000 sq. ft. Apply evenly with a drop type or rotary. Treat when insects first appear. Water grass thoroughly after application. Keep pets and children off treated area.

Lawn Chinch Bugs — Apply 1/2-1 lb. per 1,000 sq. per 5,000 sq. ft.) and brush lawn to work granules down surface. Water lawn thoroughly after application. Reapplication in 7-10 days if necessary. For longer residual control on dense growing turf such as St. Augustine grass, use 1,000 sq. ft. (5-7 1/2 lbs. per 5,000 sq. ft.).

Billbugs and White Grubs, Japanese Beetle Larva, Chafer and Southern Chafer — Apply 1 lb. per 1,000 sq. ft. lawn. Water grass thoroughly after application. For application make application any time between late July and early August. For billbug control apply when activity is first observed or when brown grass is present, indicating damage by insect.

Hyperodes Weevil (New York Only) — Apply 3/4 lb. per 1,000 sq. ft. of grass. Water grass thoroughly after application. Apply to problem areas in mid-April and again in mid-May.

SEE ADDITIONAL DIRECTIONS ON BACK PANEL


NET WEIGHT 50 POUNDS

Patterson GREEN-UP Company

Div. Curry Cartwright, Inc.
1400 Union Ave., Kansas City, Mo. 64101

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE



Granular Insecticide

For control of soil insects, corn rootworms, wireworms, cutworms, and other pests as listed.

| | | |
|---|------------------------|---------------|
| ACTIVE INGREDIENT: | | |
| O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) | phosphorothioate | 14.0% |
| INERT INGREDIENTS: | | <u>86.0%</u> |
| | TOTAL | 100.0% |

*Diazinon Trademark of Ciba-Geigy
 EPA Reg. No. 2169-265

EPA Est. No. 2169-KS-1

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person. **If inhaled** — Remove victim to fresh air. Apply respiration if indicated. **If on skin** — Wash contaminated skin with soap and water. **If in eyes** — Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

PRECAUTIONARY STATEMENTS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Not allowed. Avoid contamination of food and feed. Do not eat or drink from treated areas. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after contact. Do not allow material to be absorbed through skin.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp, crabs, birds, and other wildlife. It may also be toxic to lakes, streams, ponds, tidal marshes, and estuaries. Fish and wildlife on treated areas may be killed. Do not apply when conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply where crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Use this product only as specified on this label.

DISPOSAL

Use empty container. Wrap container and put in trash.

NOTICE OF WARRANTY

We make no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use is contrary to label instructions.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling *2/A*

LAWN INSECTS

Sod Webworms — Remove thatch. Apply 1/2-1 lb. Diazinon 14-G per 1,000 sq. ft. Apply evenly with a drop type or rotary spreader. Treat when insects first appear. Water grass thoroughly after application. Keep pets and children off treated area until dry.

Lawn Chinch Bugs — Apply 1/2-1 lb. per 1,000 sq. ft. (2 1/2-5 lbs. per 5,000 sq. ft.) and brush lawn to work granules down to the soil surface. Water lawn thoroughly after application. Repeat application in 7-10 days if necessary. For longer residual control and on dense growing turf such as St. Augustine grass, use 1 1/2 lbs. per 1,000 sq. ft. (5-7 1/2 lbs. per 5,000 sq. ft.).

Billbugs and White Grubs, Japanese Beetle Larvae, European Chafer and Southern Chafer — Apply 1 lb. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.

Hyperodes Weevil (New York Only) — Apply 3/4 lb. per 1,000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

**SEE ADDITIONAL DIRECTIONS ON BACK PANEL
 NET WEIGHT 50 POUNDS**

Patterson GREEN-UP Company
 Div. Curry Cartwright, Inc.
 1400 Union Ave., Kansas City, Mo. 64101

DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

DIAZINON
 14-G
2/A