



PROTECK-PLUS

CONTAINS BALAN[™]

Pre-emergent Herbicide + 18-3-5 Fertilizer

For use on ornamental lawns and turf. Will control crabgrass, foxtail, watergrass, annual blue grass, (poa annua), and other grasses as listed.

FOR USE ON THE FOLLOWING TURF

- | | |
|------------------------|---------------------|
| Blue grass (perennial) | Bermuda Grass |
| Ryegrass (perennial) | Centipede Grass |
| Fescue | St. Augustine Grass |
| Bahia Grass | Zoysia Grass |

ACTIVE INGREDIENT:

N-Butyl-N-Ethyl-a-a-a-Trifluoro-2,6-Dinitro-p-Toluidine	.78%
INERT INGREDIENTS	99.22%
	100.0%

BALAN[™] The Trademark of BENEFIN Elanco Product Company, a division of Eli Lilly & Company

Covers Up to 5,000 Sq. Ft. of Turf

Statement of Practical Treatment

In case of Contact: **For eyes**, flush with water for 15 minutes; **For skin**, wash with plenty of soap and water. Get medical attention if irritation persists.

See Side Panel for Additional Precautionary Statements.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Net Weight 29 lbs.

E. P. A. Reg. No. 2169-269

E. P. A. Est. 2169-KS-1

Patterson GREEN-UP Company

Div. Curry-Cartwright, Inc.

1400 Union Avenue

Kansas City, Missouri 64101

PROTECK PLUS
DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Proteck Plus is a pre-emergence herbicide and lawn food which should be applied directly to established turf for the control of smooth crabgrass, hairy crabgrass, goosegrass (silver crabgrass), watergrass (barnyard grass), yellow and green foxtail, and annual blue grass (poa annua).

It controls annual grass weeds by killing their seeds as they germinate. It does not control established weeds. When it is applied as directed the following established grasses are tolerant to Balan: perennial bluegrasses, perennial ryegrass, centipede grass, fescue, zoysia grass, bermuda grass, St. Augustine grass, and bahia grass.

Apply to turf before germination of annual grass weeds. Do not reseed within 42 days after application.

GUARANTEED FERTILIZER ANALYSIS

Nitrogen (N)	18%
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	5%

Derived from Urea, Diammonium Phosphate and Muriate of Potash. When applied to 5,000 sq. ft.; supplies 1.04 lbs. Nitrogen per 1,000 sq. ft. Supplies 1.58 lbs. "N" per 1,000 sq. ft. on 3,300 sq. ft. application.

For Use In Permanent Turf

CRABGRASS, GOOSEGRASS, FOXTAIL:

Northern Cool Season Grass — Apply bag contents to 5,000 Sq. Ft. Area.

Southern Warm Season Grass — Apply bag contents to 3,300 Sq. Ft. Area.

Note: Apply in the spring before crabgrass germinates (mid-April, south to early May, north) Apply evenly with a drop or rotary spreader.

ANNUAL BLUEGRASS (Poa-Annua)

Northern Area — Apply contents to 5,000 sq. ft. of turf. This provides 2 pounds of active Balan per acre. Treat in late summer or early fall before seed germinates.

Southern Area — On grasses such as Bermuda, St. Augustine and Bahia, apply contents to 3,300 sq. ft. This provides 3 pounds of active Balan per acre. Treat before Poa Annua germinates in early fall.

SPREADER SETTINGS

Make sure of your spreader's calibration. Use a drop or cyclone type.

	<u>Northern</u> <u>5,000 Sq. Ft.</u>	<u>Southern</u> <u>3,300 Sq. Ft.</u>
Scott	5	7
Cyclone	3½	5

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, ponds, and streams. Do not apply when weather conditions favor run off or drift from target area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open burning and dumping is prohibited. Do not reuse empty container. Pesticide Disposal — Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or local disposal authorities. Consult them for approved procedures.

USE PRECAUTIONS

This product may thin fine-leaved fescues at rates above 60 pounds per acre (1.5 pounds active ingredient). It should not be applied in the spring to bluegrass planted the previous fall. Do not apply to bentgrass putting greens, or dichondra turf. Do not apply to newly sprigged areas of bermuda grass, St. Augustine grass, or centipede grass until these turfs are well established.

NOTICE: Buyer assumes all risk and liability of use, storage, and/or handling of this material not in accordance with the terms of this label.