



# 5% DIAZINON

## CRAWLING INSECT CONTROL GRANULES

For use in lawns, gardens and around the outside of the home.

Ready To Use

**Kills:** Ants • Fleas • Brown Dog Ticks • Cutworms  
Earwigs • Millipedes • Onion Maggots  
Sowbugs • Sod Webworms • White Grubs  
and other insects as listed.

EPA Reg. 2169-271 EPA Est. 2169-KS1

<b>ACTIVE INGREDIENT:</b>	
O, O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate .....	5%
<b>INERT INGREDIENTS:</b> .....	95%
<b>TOTAL</b> .....	100%

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED** — Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.  
**IF INHALED** — Remove victim to fresh air. Apply artificial respiration if indicated.  
**IF ON SKIN** — Remove contaminated clothing and wash affected areas with soap and water.  
**IF IN EYES** — Flush eyes with plenty of water. Call a physician if irritation persists.

**Keep out of reach of children**

## CAUTION

See additional precautionary statements on back panel

**NET WEIGHT 5 POUNDS**

**patterson GREEN-UP company.**  
Div. Curry-Cartwright Inc.  
1400 Union Ave., Kansas City, Mo. 64101

**ACCEPTED**  
APR 24 1981  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is amended, for the pesticide registered under EPA Reg. No. 2169-271

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.  
**Garden Insect Control:**  
5% Diazinon Crawling Insect Control Granules controls certain garden insects on crops listed below. Apply just prior to planting and immediately mix thoroughly into soil to the specified depth, except where applied in seed furrow.

<b>Cutworms (surface and subterranean) on —</b>			
Broccoli	Cucumbers	Muskmelons	Spinach
Cabbage	Endive	Parsley	Summer Squash
Cantaloupes	(Escarole)	Peas	Swiss Chard
Cauliflower	Kale	Peppers	Tomatoes
Celery	Lettuce	Potatoes	Watermelons
Collards	Lima Beans	Snap Beans	Winter Squash

Apply 7 to 14 oz. per 500 sq. ft. and mix into soil to a depth of 2-3 inches for surface cutworms and 3-6 inches for subterranean cutworms. For potatoes, mix into soil 4-6 inches deep.

**Broccoli, Cabbage, Cauliflower**  
Root Maggots: Apply 7 to 11 oz. per 500 sq. ft. over entire seedbed before planting or transplanting and mix into soil to a depth of 3-4 inches.

**Tomatoes**  
Vinegar Flies (*Drosophila* spp.): Apply 4 oz. per 500 sq. ft. over top of plants when flies first appear. Repeat at 7-10 day intervals, or as required. Tomatoes may be picked the same day.

**Turnips**  
Root Maggots: Apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

**Carrots**  
Carrot Rust Flies: Apply 7 oz. per 500 sq. ft. in seed furrow at planting time.

**Onions**  
Onion Maggots: Apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

**Peaches**  
Root Maggots: Sprinkle lightly at the rate of 3-4 oz. per 500 linear feet of row in seed furrow at planting time.

**Ants and Wasps:**  
Spot Treatment:  
Ants: Apply 1 1/2 tsp. over and around each ant hill.  
Wasps: Apply 1 1/2 tsp. in and around each nest opening. Apply at dusk when wasps are not active.

**Foundation Treatment:**  
The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter the home. Ants, Brown Dog Ticks, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails.

**How to Apply:**  
Apply a 2-foot band of 5% Diazinon Crawling Insect Control Granules next to the house foundation. Distribute granules uniformly, using a rate of one pound per each 500 sq. ft. area (5' x 100') treated.

**When to Apply:**  
Apply when insects are present, especially in late spring or early fall. Repeat as necessary.  
**Lawn Treatment:** Controls the following lawn pests in both grass and dichondra lawns: Ants, Armyworms, Bermudagrass Mites, Billbug, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Leafhoppers, Millipedes, Sod Webworms, Sowbugs, Springtails, White Grubs of Japanese Beetle, European Chigger and Southern Chigger.

**How to Apply:**  
For most insects, sprinkle one pound over 500 sq. ft. area (10' x 50'). For heavy lawns such as St. Augustine grass, spread one pound evenly over 320 sq. ft. (10' x 32'). For billbug and white grub control, spread one pound over 100 sq. ft. (10' x 10').  
For best results, the lawn should be dry before application. To control surface feeding insects, armyworms and sod webworms (lawn moths), apply in late afternoon. Water lawn lightly after application. To control ants, earwigs, chinch bugs, cutworms, and white grubs, water lawn thoroughly after application.  
Spreader Settings: 1 lb./500 sq. ft. Scott-4 Cyclone-3 1/2; 1 lb./400 sq. ft. Scott-4 Cyclone-4 1 lb./320 sq. ft. Scott-5 Cyclone-4 1/2. Walking speed and spreader condition may require adjustments in settings.

**When to Apply:**  
Apply when insects are present. To control ants, armyworms, Bermudagrass mites, chinch bugs, cutworms, leafhoppers, and sod webworms (lawn moths), apply at 4-6 week intervals. To control white grubs, apply any time between July and early October. To control billbugs, apply when activity is first observed, or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

### Precautionary Statements Hazards to Humans & Domestic Animals CAUTION

Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling and before smoking or eating.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**Disposal:** Do not reuse empty container. Wrap container and put in trash collection.