DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Garden Insect Control

5% Diazinon Crawling Insect Control Granules controls certain garden insects. Use this product prior to planting and immediately mix thoroughly into soil to the specified depth except where applied in seed furrow.

Cabbageworm (surface and subterranean)

Broccoli	Cucumbers	Muskmelons	Spinach
Cabbage	Endive	Parsley	Summer Squash
Cantaloupe	Escarole	Peas	Swiss Chard
Cauliflower	Kale	Peppers	Tomatoes
Celery	Lettuce	Potatoes	Watermelons
Collards	Lima Beans	Snap Beans	Winter Squash

Apply 7 to 14 oz per 500 sq ft and mix into soil 1 to 3 inches for surface cutworms and 4 to 6 inches for subterranean cutworms. For potatoes mix into soil 4 to 6 inches deep.

Broccoli, Cabbage, Cauliflower

Root Maggots: Apply 7 to 11 oz per 500 sq ft of row or seed section before planting or transplanting and mix into soil to a depth of 3 to 4 inches.

Tomatoes

Vinegar Flies (Drosophila): Apply 1 oz per 500 sq ft of row or 1 tsp per plant while fruit is ripening. Repeat at 7-10 day intervals or as required. Treat the fruit the same day.

Turnips

Root Maggots: Apply 4 oz per 500 sq ft of row at seeding or planting time.

Carrots

Carrot Rust Flies: Apply 7 oz per 500 sq ft of row at seeding or planting time.

Onions

Onion Maggots: Apply 4 oz per 500 sq ft of row at seeding or planting time.

Radishes

Root Maggots: Spread evenly at the rate of 7 oz per 500 sq ft of row at seeding or planting time.

Ants and Wasps

Spot Treatment

Ants: Apply to soil over and around ant mounds.

Digger Wasps: Apply at dusk when wasps are most active. Apply at dusk when wasps are most active.

Band Treatment Around House Foundation

The following pests are commonly found around homes. Treat soil around foundation of house where these pests may enter the home: Ants, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails.

How to Apply

Apply to soil around foundation of the house to a distance of 3 feet. Do not apply to soil around shrubs or other plantings. Do not apply to soil around shrubs or other plantings. Do not apply to soil around shrubs or other plantings.

When to Apply

Apply when insects are present. Repeat as necessary.

Lawn Protection Controls the following pests in lawns: Bermudagrass, Bluegrass and Zoysiagrass.

Ants, Armyworms, Bermudagrass Mites, Billbugs, Chiggers, Chinch Bugs, Clover Mite, Crickets, Cutworms, Earwigs, Fleas, Leafhoppers, Millipedes, Sod Webworms, Sowbugs, Springtails, White Grubs (Japanese Beetle, European Chafer, Southern Chafer).

How to Apply

For most insects, spread one pound per 500 sq ft of area. For heavy infestations, spread one pound per 250 sq ft. For chiggers, spread one pound per 500 sq ft.

Augustine grass: Spread one pound per 500 sq ft every 2 months. For chiggers, spread one pound per 500 sq ft.

For best results, mow lawn before application. Water lightly after application. For ants, sod webworms, lawn moths, cutworms, and earwigs, water lightly after application. For ants, billbugs, chinch bugs, cutworms, and earwigs, water lightly after application.

Spreader Settings: 1 lb 500 sq ft: Scotts Cyclone 4; 1 lb 400 sq ft: Scotts 4; Cyclone 4. Scotts 5 Cyclone 4. Walking speed and spreader condition may require adjustments in setting.

When to Apply

Apply when insects are present. For chiggers, apply when chiggers are present. For chinch bugs, leafhoppers and sod webworms, lawn moths, apply at 4-6 week intervals.

To control white grubs, apply any time between June and early October. To control white grubs, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

Precautionary Statements

Hazards to Humans & Domestic Animals

CAUTION

May be absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Avoid contamination of food or feed products. Avoid inhalation of dust. Wash thoroughly with soap and water after handling and before smoking, eating.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to treated areas. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treated area. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treated area.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Pesticide Disposal:** Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

green UP

5% DIAZINON

CRAWLING INSECT CONTROL GRANULES

For use in lawns, gardens and around the outside of the home

Ready To Use

Kills: Ants • Fleas • Brown Dog Ticks • Cutworms
Earwigs • Millipedes • Onion Maggots
Sowbugs • Sod Webworms • White Grubs
and other insects as listed

EPA Reg. 2169-271 EPA Est. 2169-KS1

ACTIVE INGREDIENT	
O,O-Diethyl O-[2-isopropyl-6-methyl-4-pyrimidinyl] phosphorothioate	5%
INERT INGREDIENTS	95%
TOTAL	100%

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink one to two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if not breathing.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

Keep out of reach of children

CAUTION

For additional precautionary statements see back panel.

NET WEIGHT 3 POUNDS

patterson GREEN-UP company
Div. Curry-Cantwright Inc.
1400 Union Ave. Kansas City, Mo. 64101