



ACCEPTED  
2169-275  
FEB 17 1981  
Under Federal Insecticide,  
Fungicide and Rodenticide Act,  
as amended for the pesticide  
registered under  
Reg. No. 2169-275

# Proteck 3-Way

CONTAINS

**Pre-emergent Herbicide + 18-3-5 Fertilizer  
+ Broadleaf Weed Killer**

**For Lawns, Fairways and Ornamental Turf**

- Pre-Emergence Control (Balan™) — Crabgrass, Foxtail, Watergrass, Annual Blue Grass.
- Post-Emergence Control—Dandelions, Chickweed, Henbit, Knotweed, Spurge and other weeds as listed.
- Feeds Lawn with 18-3-5 Analysis.

**Use on Bluegrass, Perennial Rye, Fescue,  
Bermuda, and Zoysia Grass.**

**ACTIVE INGREDIENT:**

|   |               |
|---|---------------|
| N-Butyl-N-Ethyl-a-a-Trifluro-2,6-Dinitro-p-Toluidine .....  | 1.15%         |
| *Dimethylamine salt of 2,4-dichlorophenoxyacetic acid ..... | 1.50%         |
| **Dimethylamine salt of dicamba .....                       | .28%          |
| Dimethylamine salt of related acids .....                   | .04%          |
| <b>INERT INGREDIENTS .....</b>                              | <b>97.03%</b> |

|   |        |
|---|--------|
|   | 100.0% |
| *2,4-Dichlorophenoxyacetic acid equiv. .... | 1.24%  |
| **3,6-Dichloro-o-anisic acid equiv. ....    | .23%   |

BALAN™ The Trademark of BENEFIN Elanco Product Company,  
a division of Eli Lilly & Company

Covers 5,000 Sq. Ft. of Turf

Statement of Practical Treatment

In case of Contact: For eyes, flush with water for 15 minutes; For skin, wash with plenty of soap and water. Get medical attention if irritation persists.

See Back Panel for Additional Precautionary Statements

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

Net Weight 20 lbs.

E.P.A. Est. 2169-275  
E.P.A. Reg. No. 2169-KS-1

**patterson GREEN UP co.**

Division Curry-Cartwright Inc.  
1400 Union Avenue  
Kansas City, Missouri 64101

# SUPER PROTECK

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Super Proteck is a pre-emergence herbicide, lawn food and a broadleaf Weed Killer. It should be applied to established turf for the control of grasses (smooth crabgrass, hairy crabgrass, goosegrass, watergrass, yellow and green foxtail and annual blue grass) as well as broadleaf weeds (buckhorn, burdock, bur.clover, common chickweed, mouse ear chickweed, chicory, clover (white), creeping charley (ground ivy), curly dock, dandelion, dog fennel, frenchweed, healall, henbit, knawel, knotweed, lambsquarter, r.allow, morning glory, mustard, oxalis, peppergrass, pigweed, plantain, poison ivy, poison oak, purslane, red sorrel, sheep sorrel, sow thistle, spotted spurge, and yarrow.

**WHEN TO APPLY** — For pre-emergence action apply this product prior to the germination of weed seeds in spring or early summer (April 1st thru May 1st; or later if the pre-emergence is being applied in two separate applications, one early the other the last of May.) Broadleaf weeds present at the time of application will be controlled.

Apply when there is adequate soil moisture and the turf (weeds) are growing actively. Do not apply to newly seeded lawns.

For Annual Blue Grass treat in late summer or early fall before 'Poa' seed germinates.

**HOW MUCH TO APPLY** — Apply 20 lbs. to 5,000 sq. ft. of turf. This rate feeds the lawn with .72 lbs. nitrogen per 1,000 sq. ft.

**PRECAUTIONS IN USING** — Do not use on Centipede Grass, St. Augustine Grass, or Dichondra; nor on lawns or turf where desirable bentgrass or clovers are present. Do not allow dust to come in contact with shrubs, flowers, trees, vegetables or other valuable plants. Avoid application to areas under the branches or drip line of trees and shrubs. Do not reseed within 42 days after application.

## GUARANTEED FERTILIZER ANALYSIS

|  |     |
|--|-----|
| Nitrogen (N).....  | 18% |
| Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) ..... | 3%  |
| Soluble Potash (K <sub>2</sub> O).....                           | 5%  |

Derived from Urea, Diammonium Phosphate and Muriate of Potash.

## SPREADER SETTINGS

Make sure of your spreader's calibration. Use \_\_\_\_\_ or cyclone type.

|               |      |
|---------------|------|
| Scott.....    | 5/8" |
| Cyclone ..... | 3/4" |

## PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

### ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, ponds, and streams. Do not apply when weather conditions favor run off or drift from target area.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open burning and dumping is prohibited. Do not reuse empty container. Pesticide Disposal — Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or local disposal authorities. Consult them for approved procedures. Store in a dry area away from plants.

### USE PRECAUTIONS

This product may thin fine-leaved fescues at rates above 60 pounds per acre (1.5 pounds active ingredient).

Do not apply to bentgrass putting greens, or dichondra turf. Do not apply to newly sprigged areas of bermuda grass.

2169-275