


Treats 10,000 sq. ft.



**greenUP**

TURF-PRO



# 50% Diazinon® Pre-Weed Plus

**CONTAINS  
INSECTICIDE PLUS BALAN™ PRE-EMERGENT HERBICIDE  
PLUS 30-5-5 FERTILIZER**

For Lawns, Fairways and Ornamental Turf

• Pre-Emergence Control of Crabgrass, Foxtail, Watergrass and Annual Blue Grass

### LAWN INSECT CONTROL

For Control of Chinch Bugs, Sod Webworms (Lawn Moths), Ants, Armyworms, Billbugs, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Fleas, Sowbugs (Pillbugs), Springtails, Bermudagrass Mites, Chiggers, Clover Mites, Leafhoppers, & Millipedes infesting Home Lawns and Other Ornamental and Recreational Turf Grass Areas

Keep Out of Reach of Children

### WARNING

Read Complete Cautions on Back Panel

#### ACTIVE INGREDIENTS

O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) Phosphorothioate

N-Butyl-N-Ethyl-a-a-Trifluoro-2,6-Dinitro-p-Toluidine

#### INERT INGREDIENTS:

5.00%  
2.06%  
92.94%

**TOTAL 100.00%**

Diazinon is a registered trademark of the University of California, Davis, California.  
U.S. Patent No. 2,754,243 and other patents pending.  
BALAN™ The Herbicide of 10 NEENE, Inc., Product of the University of California, Davis, California.  
Covers 10,000 Sq. Ft. of Turf

#### — STATEMENT OF PRACTICAL TREATMENT —

**IF SWALLOWED** — Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. **IF INHALED** — Remove victim to fresh air. Apply artificial respiration if indicated. **IF ON SKIN** — Remove clothing immediately and wash affected areas with soap and water. **IF IN EYES** — Flush eyes with plenty of water. Call a physician if irritation persists.

See Back Panel for Additional Precautionary Statements

**NET WEIGHT 20 LBS.**

EPA Reg. No. 2169-291 EPA Est. No. 2169-KS-1

**patterson GREEN-UP co.**  
Division Curry-Cartwright Inc.  
1331 Union Avenue  
Kansas City, Missouri 64101

200170

## DIRECTIONS FOR USE

### IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

This product is designed for a commercial or home-owner annual Lawn Service Program which features 1 to 2 granule insecticide + pre-emergent herbicide applications per year. Insect control will be effective when applied anytime during insect activity periods. The weed control benefit will only be effective when applied before weed seeds germinate. Apply this product when the grass is dry and water lightly after application.

**INSECTS AND WEEDS** — For the control of Ants, Armyworms, Bermudagrass Mites, Billbugs, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Leafhoppers, Millipedes, and Webworms, Sowbugs, and Springtails. Pre-emergent weed control of Annual Bluegrass (Poa Annual), Smooth & Hair Crabgrass, Goosegrass (Silver Crabgrass or Crowsfoot), Watergrass (Barnyardgrass) and Yellow & Green Foxtail.


**FIRST APPLICATION** — Apply 20 lbs. per 10,000 sq. ft. The first application should be made 10 to 20 days before your local Frost Free date in the spring. This will control existing insects as listed on this label along with pre-emergence control of Crabgrass, goosegrass, foxtail, water grass and certain broadleaf weeds, namely carpetweed, chickweed, lambsquarter, purslane and pigweed.

**SECOND APPLICATION** — Apply 20 lbs. per 10,000 sq. ft. Turf that has been power raked, aerated, or verticut will benefit from this application since spring rains encourage additional weed seed germination. Apply in late May or early June after the first treatment. Insect control will be effective as in the first application, however only the ungerminated weed seeds will be controlled. Established weeds will not be effected.

**ANNUAL BLUEGRASS** (Poa Annual) — Apply 20 lbs. per 10,000 sq. ft. Treatment of heavy infestations will result in thin turf. Observe good cultural practices such as fertilization and watering to help in these areas. Repeat treatment the following year for best control.

**Northern Cool Season Grass** should be treated in the late summer to early fall (late July to late September) or in early spring prior to annual bluegrass seed germination. It is difficult to get 100% control the first year in heavy turf growth.

### WEED CONTROL FOR ANNUAL GRASSES



Approximate weed germination times (1) Crabgrass, Foxtail, Watergrass, Goosegrass; (2) Annual Bluegrass (Poa Annual)

**INSECT CONTROL THROUGH THE YEAR** — If subsequent generations of insects appear in the turf after these applications continue throughout the season with 5% Diazinon Crawling Insect Control Granules, EPA Reg. No. 2169-271 according to the directions on the label.

**RESEEDING** — Delay reseeding for at least 6 weeks after application of this product.

**WHERE TO USE** — When applied as directed the following grasses are tolerant: perennial bluegrass, perennial ryegrass, centipedegrass, fescue, zoysiagrass, bermudagrass, St. Augustinegrass and Bahia grass. Apply only to established stands of these grasses.

### GUARANTEED FERTILIZER ANALYSIS

Nitrogen (N)	30%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	5%
Soluble Potash (K <sub>2</sub> O)	5%
Nutrients derived from Urea, Diammonium Phosphate, Potassium Sulfate Supplies 6 lb. N per 1,000 sq. ft. of turf when applied as directed.	

**SPREADER SETTING** — Scott 4 Cyclone 34

Walking speed and spreader condition may require setting adjustments.

### PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

## WARNING

### KEEP OUT OF REACH OF CHILDREN

May be absorbed through the skin. Keep children and pets off treated areas until this material is washed into the soil and the grass is dry. Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling and before smoking or eating.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Dispose of Bags** — According to approved Federal, State or Local Procedures under the Resource Conservation and Recovery Act.

**Pesticide Disposal** — Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or Local Procedures under the Resource Conservation and Recovery Act.

### NOTICE

Buyer assumed all risk and liability of use, storage and handling of this material. This material is sold in accordance with the terms of the label.

## NET WEIGHT 20 LBS.

patterson GREEN-UP co.  
Division Curry-Cartwright Inc.  
1331 Union Ave. Kansas City, Missouri 64101