




Page 1 of 4



BETASAN 7-G

GRANULES - A Selective Herbicide for Pre-emergence Control of Crabgrass and Annual Bluegrass in Turf, Ornamentals and Ground Covers

Contains 7 pounds of active ingredient per 100 pounds.

ACTIVE INGREDIENTS:

S-(O-diisopropyl phosphorodithioate of	
N-(2-mercaptoethyl) benzenesulfonamide	7.0%
INERT INGREDIENTS:	93.0%
U.S. Patent No. 3,205,253	100.0%

SPA Reg. No. 2169-

EPA Est. No. 2169-KS-1

12 Pounds Treats 5,000 Sq.Ft.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Precautionary Statements

Betasan - Reg. T.M. Stauffer Chemical Co.

STATEMENTS OF PRACTICAL TREATMENTS.

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never give anything by mouth to an unconscious person.

IF IN EYES: Wash with plenty of water. Get medical attention if irritation persists.

Net Weight 12 Pounds

Patterson Green-Up Company
Div. Curry Cartwright Inc.
1331 Union Ave.
Kansas City, Mo. 64101

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RATES OF APPLICATION

CRABGRASS ONLY: Apply 6 pounds of BETASAN-7G granular uniformly over 2,500 square feet. (107 pounds per acre) Scott Spreader Setting - 2.8(once over) and 2.2(twice over) using a 20 inch width.

ANNUAL BLUEGRASS AND OTHER ANNUAL WEEDS: Apply 10 pounds of BETASAN-7G granular uniformly on 2,500 square feet. (180 pounds per acre). Scott Spreader Setting- 3.2(once over) and 3.1(twice over) with 20 inch width.

WEEDS CONTROLLED

Smooth Crabgrass (*Digitaria ischaemum*)
Hairy Crabgrass (*Digitaria sanguinalis*)
Redroot Pigweed (*Amaranthus retroflexus*)
Armyardgrass (Watergrass) (*Echinochloa* spp.)
Lambquarters (*Chenopodium album*)
Moosegrass (*Eleusine indica*)
Annual Bluegrass (*Poa annua*)
Shepherdspurse (*Capsella bursa-pastoris*)
Deadnettle (Henbit) (*Lamium* spp.)

GENERAL USE PRECAUTIONS

BETASAN should be used only for recommended uses and at recommended rates. Sweep up spills on walks or patios and hose area. Do not allow children or pets on treated lawns until BETASAN has been watered into the soil and the grass is completely dry. Toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

PRODUCT INFORMATION

BETASAN is recommended for use on home lawns, parks and golf courses, on ornamental plants and on ground covers for control of crabgrass, annual bluegrass and other annual weeds listed.

DIRECTIONS FOR APPLICATION

Application of BETASAN must precede emergence of weeds from the soil. Crabgrass will emerge from the soil from four to six weeks before it is visible above lawn sod. If weeds are visible above the soil, it is too late to apply BETASAN.

Remove leaves, dead or tall grass, and other debris before applying BETASAN to turf areas. Any weeds present around ornamentals must be removed or worked into the soil before applying BETASAN.

BETASAN is recommended for use on mineral soils only.

Do not apply peat moss to lawn, ground covers or established ornamentals before treating with BETASAN.

TIMING OF APPLICATION - GENERAL COMMENTS

GRASS LAWNS: BETASAN should be applied only to well established grass lawns. BETASAN can be used on the following established turf grasses.

BAHIA	CENTPEDE	POA TRIVIALIS
BENTGRASS SPP.	FESCUE SPP.	SAINT AUGUSTINE
BERMUDAGRASS SPP.	PENSACOLA BAHIA	RED TOP
PERENNIAL BLUEGRASS SPP.	PERENNIAL RYEGRASS	ZOYSIA

DIRECTIONS FOR USE (cont.)

DICHONDRA LAWNS: BETASAN can be applied to dichondra lawns at the time of seeding or any time thereafter.

ORNAMENTALS AND GROUND COVERS: BETASAN should be applied only to the following well established ornamentals and ground covers:

ALYSSUM	CORAL BELL	PACHYSANDRA
ASTER	DAISY	STOCK
BACHELOR'S-BUTTON	MARIGOLD	SWEET PEA
CALENDULA	PANSY	WALL FLOWER
CANDY-TUFT	PRIMROSE	ZINNIA
DAFFODIL	GLADIOLUS	RANUNCULUS
DAHLIA	NARCISSUS	TULIP
FREESIA	JUNIPER	PYRACANTHA
ABELIA	MONTEREY CYPRUS	SANDANKWA
AZALEA	MONTEREY PINE	STAR JASIME
AZARA	MYRTLE	TOBIRA
BOXWOOD	HYPERICUM	PERIWINKLE
AJUGA	ICE PLANT	SEDUM
CAMPANULA	IVY	WILD STRAWBERRY
GAZANIA		

For control of crabgrass, apply BETASAN at recommended rate (see RATES OF APPLICATION) anytime from fall - September or October - through early spring and summer control. With a late winter or an early spring application, the application must be made prior to crabgrass emergence (see DIRECTIONS FOR APPLICATION).

For control of annual bluegrass and other annual weeds, apply BETASAN at the recommended rate (see RATES OF APPLICATION) before the weeds emerge from the soil.

In southern and western states where germination of crabgrass, annual bluegrass and other weeds extends into the late summer and early fall, a second application of BETASAN at the recommended rates should be made four to five months after the initial application.

BETASAN should be applied with a lawn spreader or any other applicator that will give uniform coverage. A more uniform application can be made by spreading half the required amount of BETASAN over the area to be treated, then applying the remainder at right angles or crosswise. Avoid misses by overlapping slightly. Sprinkle the area with water for 10 to 15 minutes after application to wash the BETASAN down into the soil where the weed seeds are located.

Consult weed specialists of the State Agricultural Experiment Station or the State Agricultural Extension Service for additional information, as the timing and rate of application needed will vary with local conditions.

Notice - Buyer assumes all risk and liability for use, storage and/or handling of this material not in accordance with the terms of this label.

DIRECTIONS FOR USE (cont.)

RESEEDING

Any material that provides effective control of annual grasses may affect new lawn grass seedlings which germinate after treatment. Proper fertilization and watering will encourage the existing turf grass to fill in the thin or bare spots formerly occupied by crabgrass.

The following procedures are suggested in order to minimize any stunting of turf grasses which may occur when reseeding into BETASAN-treated areas:

1. Do not reseed within 4 months after application. Renovate the area to be reseeded by raking thoroughly to scarify the soil surface. Where practical, mix peat in with the soil to provide an optimum seed bed.
2. When turf diseases or physiological disorders necessitate reseeding sooner than 4 months after BETASAN application, apply powdered activated charcoal prior to reseeding to deactivate the BETASAN.

Apply the activated charcoal at a rate of 7 lbs. per 1,000 square feet in 14 gallons of water per 1,000 square feet (300 lbs. per acre in 600 gallons of water per acre).

Following application, irrigate the turf with sufficient water to wash the charcoal into the soil. Reseed no sooner than 7 days after applying the charcoal.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Wash thoroughly after use. Sweep up spills on walks or patios. Bury sweepings and hose area. Do not allow children or pets on treated lawns until BETASAN has been watered into the soil and grass is completely dry. Do not contaminate food, feed, or drinking water.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes. Keep out of lakes, streams and ponds. Apply this product only as specified on this label. Read all label directions before using.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE AND BAG DISPOSAL

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.