It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Mixing Instructions Mix 4 1/2 tempoonful: per gallon of water for all uses, applying thoroughly to both upper and under sides of foliage when mites first appear and as necessary thereafter. [Rate is equivalent to 4 2/3 pints per 100 gallons of water.]

FLOWERS, ORNAMENTALS, NURSERY STOCK, TREES: To control two-spotted mite (red spider). European red mite. Spruce mite and other mites common on both greenhouse and outdoor plants such as roses, evergreens, shade trees, flowers, nursery stock and ornamentals. Do not use on Canaerti juniper.

VEGETABLES: To control two-spotted mites (red spider), Pacific, Tropical and tomato russet mites on beans, cucumbers, cantaloupes, nielons, watermelons, pumpkins, winter and summer squash, tomaties, eggplants and peppers. Do not use on beans within 7 days of harvest or on other vegetables within 2 days of harvest. Do not feed vines to meat or dairy animals.

strawberry mite and six-spotted mites (red spider), strawberry mite and six-spotted mite, use as above. To control cyclamen mites used to Steaspoonfuls per gallon water and drench plants thoroughly at 10 to 20 day intervals in spring when new growth starts. Repeat as necessary. Do not use within 2 days of harvest.

CLOVER MITES OUTSIDE HOMES: To control clover mites, spray walls, foundations, window-sills; also plants and lawn at base of building.

BERRIES, GRAPES, NUTS (as listed): To control brown almond (clover), two spotted (red spider), European red, Pacific, McDaniel, Willamette and yellow mitter on grapes, respheries, dewberries, loganberries, boysenberries, walnuts, filberts, pecans, chestnuts and hickory nuts, apply just before bloom or whenever mittes appear. On nuts do not apply within 14 days of harvest. Do not use on grapes within 7 days of harvest or on berries within 2 days of harvest.

FRUITS: APPLES, PEARS, CRABAPPLES, QUINCES — For European red mite, clover mite, two-spotted (red spider), Schoene. McDaniel, yellow (carpini), Willamette, Pacific, and apple rust mites. Apply in second or later cover sprays or whenever mites appear but not later than within 7 days of harvest.

PEACHES, APRICOTS, NECTARINES, CHARRIES, PLUMS, PRUNES, FIGS: For European red, clover, (brown almond) two-spotted (red Spider), Schoene, McDaniel, Pacific, Willamette, peach silver, plum, nursery and yellow mites. Apply at petal fall to first cover spray or whenever mites appear. If repeat application is necessary, wait 30 days to avoid excessive residues. Do not apply to cherries, plums, prunes or figs within 7 days of harvest or un other fruits listed above within 14 days of harvest.

CITRUS FRUITS: Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines. For the following mites; Citrus Flat, Citrus Red (purple), Six Spotted and Yuma Mites.

Do not use within 7 days of harvest. Follow local timing recom-

mendations from state extension service Specialists. Do not apply to Avacados.

# PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

## Caution

Harmful if awallowed. Avaid inhalation of spray mist; skin contact, contamination of feed or foodstuffs. Wash thoroughly after handling.

### **Environmental Hazards**

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### Physical & Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse container. Wrap and put in trash collection.

NOTICE: Buyer assumes all risk and li ability of use, storage and/or handling of this material not in accordance with the terms of this label.

DIRECTIONS FOR USE ( Cont.)

Clover Mites on Buildings: Mix at same rate as for spraying plants. Thoroughly spray outside walls, foundations, windowsills, plants, & lawns at base of infested buildings. Lawn & Turf Grasses: For Bermuda, Brown, and Clover Mites mix at same rate as for plants. Apply 2 1/2 Gal. diluted spray per 1,000 sq.ft. Treat entire lawn thoroughly and repeat in 10 to 14 days if necessary.

ILC ENVIRONMENTAL PROTECTION ACENCY	EPA REGISTRATION NO.	DATE OF ISSUANCE
U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (MH-567) ASSMINGTON TICLS 46	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	
ROTICE OF PESTICIDE: 1 SHERES STRATUS		
(Und. r the Federal Insecticide). Furnicide and Rodentis (de Act, 31) anended).	Í	
AME AND ADDRESS OF REGISTRANT (Include ZIP . ede)		
г	П	
L		
NOTE: Changes in labeling formula differing in substance submitted to and accepted by the Registration Division proproduct always refer to the above U.S. EPA registration numbers.	or to use of the label in commer	n with this registration must be ce. In any correspondence on this
On the basis of information furnished by the registrant, the the Federal Insecticide, Fungicide, and Rodenticide Act.	above named pesticide is here	by Registered/Rere - ered under
A copy of the labeling accepted in connection with this Re	egistration/Reregistration is ret	urned herewith,
Registration is in no way to be construed as an indorsement health and the environment, the Administrator, on his moticized in accordance with the Act. The acceptance of any na Act is not to be construed as giving the registrant a right to by others.	on, may at any time suspend or ame in connection with the regi	runce) the registration of a pest- stration of a product under this
·		• •
	,	
		31
		• •
•		
	•	•
		•
ATTACHMENT IS APPLICABLE		ė,
SIGNATURE OF APPROVING OFFICIAL		DATE

# **ZOECON** RF-199 GOLDEN **FLY BELT**

WITH MUSCAMONER tly attractant

[Attracts and Kills More Flies]

[Fly Bait Belt]

[Contains Improved GOLDEN MALRINK Fly Bait.]

[Flies feed on (the bait)/(Improved GOLDEN mALRIN<sup>R</sup> Fly Bait) and die.]

[No Sticky Glue Mess.]

[(Fly Belt is) Not a sticky fly paper.]

[Flies ingest a lethal dose of (bait)/ (Improved GOLDEN MALRINK Flu Bair) and [BUZZ-OFF to] die.]

ACTIVE INGREDIENTS:

Methomy1 [S-methy1 N-(methy1-carbamoy1) oxy] thioacetimidate................ 1.000. (2)-9-tricosene...... 0.025t 100.066% Total

# KEEP OUT OF REACH OF CHILDREN CAUTION

See back Panel For Additional Precautions

# **CONTENTS:**

#### PRECAUTIONARY STATEMENTS

HAZAROS TO FIMANS AND COMESTIC ANIMA'S - CAUTION: Harmtol of smallowed, inhaled or absorbed through the skin. Avoid breatning dust. Avoid contact with skin, eyes or clothing.

STATEMENT OF FMACTICAL INSAIMPAT: If swallower + Orion 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Cet medical attention. It on skin - Remove contaminated clothing and wash affected areas with soap and water. If in eyes - Flush eyes with plenty of water. Call a physician immediately.

NOTE TO PRINTCHAR: Methomyl is a cholinesterase inhibitor.

On not contaminate feed and tronstuffs. No not apply to surfaces in reach of poultry or animals, especially young calves. Do not use in homes or where milk is processed or stored. Attack to surfaces out of reach of food producing animals. Do not use in edible product areas of food processing plants, restaurints or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

ENVIPONMENTAL PAPAREDS: This product is toric to fish, birds and other wildlife. Exposed surfaces treated with this product are hexardous to birds and other wildlife. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIMESTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Zoecom PF-199 Golden fly Belt contains Muscamone fly attractant which encourages both male and remale flies to remain on the belt, thus feeding on the bait. The helt may be (cut to any desired length and) attached to posts, walls, ceilings and other locations where files tend to feed, rest or congregate. Be sure not to locate where food may be contaminated, or where animals or children may have access to the bait. Place belt (Place cards at approximately 10-20 ft. intervals) in areas sheltered from precipitation. (For heavy infestations, auditional cards may be used.) Replace belt when effectiveness diminishes.

Use Zo-con FF-199 Golden Fly Belt in (stables,) (dairy barns,) (kennels,) (poultry and) (swine houses,) (brotler houses,) (livestock barns,) (call barns,) (annual sheds,) and (pet shelters). Use Toeron RF-199 Colden Fly Belt (annual sheds,) and (pet shelters). Use Toeron RF-199 Colden Fly Belt (annual sheds,) around the cutsine of (food processing plants,) (meat and poultry processing plants,) (feed loss,) (feed mills,) (fast-food establishments,) (canneries,) (beverage plants,) (behavios,) (supermarkets,) (restaurants,) and (refuse dumpsters).

STORAGE & DISPOSAL to rot contaminate mater, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Fontainer Pisposif - Discose of empty containers in a sanitary landfill or by immineration if allowed by state and local authorities.

Sellen makes no warranty, express or implied, concerning the use of this product other than indicated or the latel. Buyer assumes all misks of use and handling of this material when such use and handling are contrary to label instructions.

loecon Industries A frankler of one on Corporation
That Counten Drive, Pallas, Texas 75/24

EVA Peg. No. 1/44 46 MUSERMON IS a tradement of Zoncon Gorp.

EPA Est. Hade In U.A c 19-3 Zuecon lé∍-C