

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (WH 567)
WASHINGTON, D.C. 20460

EPA REGISTRATION NO.

DATE OF ISSUANCE

29 OCT 1994

TERM OF ISSUANCE

NAME OF PESTICIDE PRODUCT

NOTICE OF PESTICIDE: ☐ REGISTRATION
☒ REREГИSTRATION

(Under the Federal Insecticide, Fungicide
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

┌

└

186/129055
17/1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.



ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

89 273
N 2169
331

ACCEPTED
with COMMENTS
in EPA Letter Dated:

OCT 29 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this product is
registered and labeled for use as

2169-331

Homeowner Protective Clo
Homeowner applicators must wear
protective clothing during the a
process: long-sleeved shirt, lon
proof gloves, full foot covering
covering.

CAUTION: Do not allow children t
this product. Children and pets
allowed in treated areas until t
are dry.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING — Causes eye damage and skin irritation. Do not get in
eyes, on skin, or on clothing. May be fatal if swallowed or absorbed
through the skin. Do not breathe vapor or spray mist.

Wash hands, arms, and face thoroughly with soap and
water after handling and before eating or smoking. Do not use on
humans or household pets. Keep children and animals away from
treated areas until spray has dried. Food utensils such as a tea-
spoon or measuring cup should not be used for food purposes after
use with pesticides.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply
directly to lakes, streams or ponds. Do not apply when weather
conditions favor drift from areas treated. Do not contaminate water
by cleaning of equipment or disposal of wastes. Apply this product
only as specified on the label. This pesticide is highly toxic to bees
exposed to direct treatment or to residues remaining on the treated
area. Do not apply when bees are actively visiting the crop. Cover
crops or weeds blooming in the treatment area. Applications should
be timed to provide the maximum possible interval between treat-
ment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.


STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage — Store in a cool dry place.

Pesticide Disposal — Pesticide, empty mixture, or rinse water that
cannot be used according to label instructions must be disposed of
according to applicable Federal, state, or local procedures.

Container Disposal — Triple rinse (or equivalent). Then offer for
recycling or reconditioning, or puncture and dispose in a sanitary
landfill, or by other approved state and local procedures.



**LINDANE 20%
INSECTICIDE**

**INSECT CONTROL FOR ORNAMENTALS — FRUIT TREES — FLOWERS
BULBS & VEGETABLES**

ACTIVE INGREDIENTS

Lindane (Gamma isomer of Benzene Hexachloride) 20.00%

Xylene 88.89%

INERT INGREDIENTS 13.15%

TOTAL 100.00%

12 PT. WARNING: KEEP OUT OF REACH OF CHILDREN 4-8 PT.
See Side Panel For Additional Precautionary Statements

NET CONTENTS: 1 PINT

P.O. Box 4029

patterson GREEN UP company

Kansas City, Missouri 64101

Division Curry-Ca

STORAGE & DISPOSAL ← 8 PT.

Do not contaminate water, food, or feed by storage or disposal.

STORAGE — Store product in original container in a cool, dry,
locked place out of reach of children.

DISPOSAL — Do not re-use empty container. Wrap container and put
in trash.

INSECT CONTROL FOR CUMMINS, PINE TREES, FLOWERS, BULBS, VEGETABLES
(Do not use on root crops or in soil in which root crops are to be planted)

PLANTS	INSECTS	DOSAGE RATE IN FLUID MEASURE	COMMENTS
Elm, Birch, Locust, Holly	Leaf Miners	1 tsp./gal. water	Apply when adult miners appear in April & mid-May. Treat when smaller miners appear in July.
Pines	Pine Bark Beetles	3 Fluid Ounces per gal. water	Apply to trunks up as high as sprayer will reach. Spray to point of run-off in April or May.
Pine, Spruce	White Pine Weevil	1 tbsp./gal. water	Thoroughly spray entire tree in late April when buds swell.
Taxus, Hemlock	Taxus Weevil	1 tbsp./gal. water	Apply in May or early June to the trunks & around tree base. Soak the bark and soil surface.
Dogwood, Lilac, Rhododendron	Dogwood Borer, Lilac Borer, Rhodo. Borer	1 tbsp./gal. water	Make 3 applications. May 1st; June 1st; July 1st. Treat limbs, twigs and base of trunk.
Apple, Peach, Cherry	Bark Beetles Flat-headed & Round headed Borer	1 tbsp./gal. water	Apply to bark of tree and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to runoff. Do not apply to fruit or foliage. Do not apply within 60 days of harvest.
Peaches	Peach Tree Borer	1 tbsp./gal. water	Apply to bark of tree and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to runoff. Do not apply within 60 days of harvest.
Maple, Aspen, Poplar, Ash, Willow, Elm, Locust	Aphids, Bark Beetles, Borers	2 tbsp./gal. water	Apply in spring when buds begin to break. Repeat in 10-14 days. For Aphids, apply when present in summer & fall.
Birch	Bronze Birch Borer	1 tbsp./gal. water	Make 3 treatments. May 1st; June 1st; July 1st. Soak bark.
Bulbs (Tulip, Iris, Glads, Lilies)	Aphids, Bulb Fly, Iris Borer, Thrips	1 tbsp./gal. water	Treat when insects appear. For Iris borer soak soil around tuberous roots May & June.
Vegetable Sets & Cabbage	White Grubs, Cutworms, Wireworms, Soil Treatment	1 tbsp./2 gal. water. Use 2 gal. of mixture to treat 250 sq. ft. of soil area	Do not apply after thinning. Do not treat plant foliage. Do not use in soil to be used for root crops. Apply around base of plant within 24 hours after sets are placed in soil. Repeat in 14 days.
Melons, Peppers, Okra, Tomatoes	Same for insects as for Cabbage	Same as for cabbage	Do not apply after edible parts start to form.
Flowers, Roses, Petunias, Carnations, etc	Aphids, Thrips, White Flies, Leafhoppers	1 tbsp./2 1/2 gal	Apply as insect appear. Repeat sprays as needed for control. Thoroughly spray