

BEST AVAILABLE COPY



2169-330  
Page 1 of 3  
Pm25

F R E C U R F P L U S

FOR BERMUDA GRASS LAWNS


A Pre & Post Emergence Herbicide

CONTAINS A 13-0-6 FERTILIZER WITH A  
UREA FORMALDEHYDE SLOW RELEASE NITROGEN BASE

Use In The Dormant Or Growing Season

Kills Grassy & Broadleaf Weeds In Bermuda Lawns

COVERS 5,000 SQ. FT.



ACTIVE INGREDIENT:

3,5-dichloro-N-(1,1-dimethyl-2-propynyl)

terzamide . . . . . 0.57%

INERT INGREDIENTS . . . . . 99.43%

Total 100.00%

Contains "Kerb" Herbicide Rohm & Haas Company Reg. T.M.  
U.S. Patents 3,534,098 and 3,640,699.

KEEP OUT OF REACH OF CHILDREN --- --- --- 12 PT.

CAUTION --- --- --- --- --- 18 PT.

See Back Panel For Additional Precautionary Statements

STATEMENTS OF PRACTICAL TREATMENTS

If Swallowed - Call a physician or Poison Control Center immediately.

Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never give anything by mouth to an unconscious person. If In Eyes - Rinse eyes with plenty of water. Get medical attention if irritation persists. If On Skin - Wash with soap and water.

EPA Reg. Number 2169-

EPA Est. No. 2169-MO-1

NET WEIGHT 20 Pounds

PATTERSON GREEN-UP COMPANY  
Div. Curry Cartwright, Inc.  
1331 Union Avenue-Box 4029  
Kansas City, MO - 64101

(For convulsions)

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECURB is a selective herbicide for use on Dormant or Actively growing Bermudagrass lawns & ornamental turf. At recommended rates it is generally effective against the following list of Annual and Perennial Grasses and certain Broadleaf weeds as listed. This product also contains a Slow Release Nitrogen base fertilizer from Ureaformaldehyde (Nitroform). When applied as directed it will supply the Bermuda grass with .52 pound ( 1/2 lb.) of nitrogen per 1,000 Sq.Ft. of turf. The lack of Phosphate in this product will discourage the growth of Annual Bluegrass (Poa Annua) which will enhance the Herbicide effect on this grassy weed.

**GRASSES** - Annual Bluegrass(Poa Annua), Perennial Bluegrass, Barnyardgrass, Canarygrass, Cheatgrass, Crabgrass, Fall Panicum, Foxtail, Foxtail Barley, Goosegrass, Lovegrass, Orchardgrass, Quackgrass, Ryegrass, and Volunteer Oats, Barley, Rye and Wheat.

**BROADLEAF WEEDS** - Annual Morning Glory, Carpetweed, Chickweed, Henbit, Knotweeds, Lambsquarter, Nettle, Nightshade, Purslane, Smartweed, and Volunteer Tomato.

**WHERE TO USE** - For use only on Bermudagrass lawns since bermuda is resistant to the herbicide action of Precurb. It also is not effective against Nutgrass, Johnson Grass or any number of Composite Family plants such as Dandelion, Dogfennel, Groundsel, Sow Thistle & Wild Lettuce or leguminous species such as Clover or Black Medic. These must be controlled by other suitable EPA approved herbicides.

This product is especially effective for Pre & Post Emergence control of Poa Annua (Annual Bluegrass) growing in Bermudagrass. It will control "Poa" from pre-germination and seedling stages through tillering, heading and seed formation. It kills plants slowly. Target weeds may first become dark green & then gradually turn yellow & die over a 3 to 5 week period.

The following plants are sensitive to Precurb treatments and should not be treated unless extensive damage can be tolerated: Dichondra, Bluegrass, Annual and Perennial Ryegrass (includes all new patented strains of Ryegrass), Rescue Grasses, and Bentgrass.

Do not use on areas that are to be overseeded with susceptible cool season grasses within 90 days of treatment. Do not treat fairways or approaches that drain into Bentgrass greens.

**HOW IT WORKS** - This product works mainly thru the root absorption of the herbicide. If no rain occurs, the application should be followed with a light overhead irrigation to move the chemical into the root zone. Avoid heavy irrigation of more than 1 inch water which might wash or leach the herbicide from the area of application.

**HOW & WHEN TO USE** Apply contents of bag to 5,000 Sq.Ft. of lawn surface. The turf should be mowed first, catching the clippings.

May be applied to Bermudagrass during the dormant or growing season. A Fall-Winter application should be made where weeds have become established during the summer months. All susceptible weeds will die during the winter. Fall treatment should be made on winter annuals such as Chickweed. For Quackgrass and other perennial weedy grasses apply in the fall prior to winter freeze-up.

SPREADER SETTINGS : Scott 5 1/2 Cyclone 3 1/2

Walking speed and spreader condition may require adjustments in setting.

DIRECTIONS FOR USE(Cont.)

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION - Harmful if Swallowed. Take special care to avoid contact with eyes, skin or clothing. In case of contact wash with soap and water. In case of eye contact wash with plenty of water. Do not breath dust. Wash clothing & gloves after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original container in a cool, dry, locked, place out of reach of children. Do not re-use empty bag. Wrap bag and put in trash.

NOTICE - Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

GUARANTEED FERTILIZER ANALYSIS

Nitrogen (N) . . . . .	13.0%
Available Phosphoric Acid( $F_2O_5$ ) . . . . .	0.0%
Soluble Potash ( $K_2O$ ) . . . . .	6.0%

Nutrients derived from Urea Formaldehyde & Sulfate of Potash  
Supplies turf with 1/2 lb. UF Nitrogen & 0.25 lb. Potash/1,000 Sq.Ft.  
when applied according to instructions.

SPREADER SETTINGS - Scott - 1 1/2 Cyclone - 3 1/2

Walking Speed and Spreader Condition may require setting adjustments