

ACCEPTED
2175-7
APR 24 1979
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND FUNTICIDE ACT
EPA PESTICIDE REGISTRATION NO. 2175-7



MERICIDE B.C.P. #1

THREE-M-SUPPLY COMPANY

East Columbia - Seattle - Washington 98122 - 329-1200
Tacoma - 833-9888 - Spokane - 477-0597
Wenatchee - 651-7653

ACTIVE INGREDIENT

Poly(oxyethylene (dimethyliminio) ethylene

dimethyliminio ethylene 100 %

INERT INGREDIENT

0 %

This product contains 0.85 lb. of active ingredient per gallon and weighs 8.49 lb. per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed. Avoid breath vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MERICIDE B.C.P. #1 is used to control algae, bacteria and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits. An initial slug addition of 6.13 to 15.32 fluid ounces of MERICIDE B.C.P. #1 per 1000 gallons of water to provide a concentration of 48 to 120 parts per million of MERICIDE B.C.P. #1 based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 1.53 to 15.32 fluid ounces of MERICIDE B.C.P. #1 per 1000 gallons of water (12 to 120 parts per million of MERICIDE B.C.P. #1) should be employed every 2 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

MERICIDE B.C.P. #1 is used to control bacteria in industrial air washing systems that maintain effective mist eliminating components. Prior to its use systems should be cleaned to remove bacterial slime and other deposits. An initial slug dose of 24.7 to 41.2 fluid ounces of MERICIDE B.C.P. #1 per 1000 gallons of water is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 16.9 to 41.2 fluid ounces of MERICIDE B.C.P. #1 per 1000 gallons of water should be employed each 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and severity of the bacterial problem. Slug additions may be made to the sump or to the water collection trays of the airwash system.

STORAGE & DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill or bury in a safe place. Rinse that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured by
Three - M Supply Company
1215 East Columbia Street
Seattle, WA 98122

EPA REG NO 2175-7 EPA EST NO 2175 WA 1

BEST DOCUMENT AVAILABLE