DIRECTIONS FOR USE:

To clean and disinfect floors, walls and furniture, wash with solution of one (1) ounce Elimstaph #3 per gallon of water. For porous or heavily contaminated surfaces, use two (2) ounces Elimstaph #3 per gallon of water. Surfaces to be disinfected must be thoroughly wetted and gross filth removed.

Elimstaph #3 is for use on floors, walls and other hard surfaces in Hospitals, Nursing Homes, Hotels, Public Areas, Industrial Areas and in Institutions. It is excellent for Operating Rooms, Rest Rooms, Nurseries and Food Service Areas. Especially effective on Conductive floors.

When used as directed, Elimstaph #3 is effective against Staphylococcus aureus, Salmonella choleraesuis and Pseudomonas aeruginosa. Elimstaph #3 is also effective against pathogenic fungi, Trichophyton interdigitale.

ELIMSTAPH #3 GERMICIDAL CLEANER DETERGENT — DISINFECTANT DEODORANT — GERMICIDAL PATHOGENIC FUNGICIDE

FOR HOSPITAL USE

Phenol Coefficients
Guaranteed Minimum
S. aureus 78
S. typhosa 48

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% 5% C18) dimethyl benzyl	6 C12,
ammonium chlorides	4.5%
n-Alkyl (68% C12, 32% C14)	
dimethyl ethylbenzyl ammoni	
chlorides	4.5%
Tetrasodium ethylenediamine	
tetraacetate	2.0%
Sodium Carbonate	4.0%
INERT INGREDIENTS:	85.0%
DANGER:	100%

Keep out of Reach of Children
Rinse empty container thoroughly with water and discard.

E.P.A. Registration No. 2212-7
WALTER G. LEGGE COMPANY, INC.
101 Park Avenue, New York, N.Y. 10017

DANGER

Keep out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food. Protect eyes and skin when handling concentrate.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against shock, respiratory depression and convulsion may be needed.

ACCEPTED WITH COMMENTS

ACCEPTED

13