

When used as directed, will give control of many annual and biennial weeds in corn, wheat, sorghum, barley, rice, oats, rye, timothy, pastures, turf and lawn areas. It is more effective when used on soil actively growing weeds. Temperatures best for spraying range from 70° to 90°. Do not apply if rain is predicted within the hour. Use low pressure spray at 40-50 p.s.i. or less. Avoid drift. Border areas may be sprayed with spot spraying technique. Do not spray if wind is blowing from the direction of the water.

**Early Spraying** When corn is 2 to 4 inches high, apply as soon as possible after most weeds have emerged. Use 1 pt. per acre in sufficient water to give uniform coverage. Corn drop nozzles are not necessary at this time.

**High Clearance Spraying:** This is a useful spray, especially when you've just started after Lay-By. Use it for the first 2-3 days after spraying nicotine to hit the nicotine.

**HARD TO KILL WEEDS IN NON-CROP LANDS**—For perennial weeds such as Star Thistle and Russian Knapweed (see list in *California Weeds*), fence rows and farm buildings, use 2 to 3 pints in 20 gallons water with 2 to 3 oz. liquid detergent and thoroughly wet weed foliage. Do not let spray drift. Treat in spring and again at bloom time before weeds go to seed, against the following year.

814-473

|                                      |        |
|--------------------------------------|--------|
| 2,4-Dichlorophenoyl acid             | 49.3%  |
| 2,4-Dichlorophenoyl acid             | 89.3%  |
| 2,4-Dichlorophenoyl acid             | 100.0% |
| *2,4-Dichlorophenoyl acid equivalent | 41.9%  |

Contains 4 H<sub>2</sub>O, 2.4-D acid per carbon

**CAUTION:** Keep Out of Reach of Children  
See Side Panel for Additional Cautions

NET CONTENTS      GALLONS

**SMALL GRAIN** (Wheat, Barley, Rye, Oats, etc.)—Treat seeds in **Winter Wheat** and **Rye**, treat before planting in the spring. Use 1 to 1½ pt. in sufficient water to cover one acre. Apply from the tiller to the jointing stage. In **Spring Wheat, Oats,** and **Barley**, use 1 to 1½ pt. per acre. Use higher dosage for control of weeds. Treat when crop is fully tillered, 4 to 8 inches high, up until early boot stage. Do not apply during seedling stage or boot stage.

**PASTURES AND TURF:** Use 2.3 pts. in 10 gal. of water per acre. Apply early in summer when plants are actively growing. Treat again in bud stage before weeds go to seed. **CAUTION:** On dairy pastures, remove cows during spray application and for 24 hours allow them access to the treated area for forage. Do not apply to newly seeded areas until grass is well established.

For lawns use 1½ tablespoons per gallon for 1,000 sq. ft. Repeat as needed for control. Do not spray new lawns until after first cutting. Clover may be damaged with this spray. Do not use on bentgrass, St. Augustine grass, Dichondra or carpetgrass lawns.

**Avoid inhalation of spray mist.**

E.P.A. Reg. No. 2217-2-AA

814473