

**HOUSEHOLD PESTS:** For control of Roaches, Ants, Silverfish, Waterbugs, Crickets, Spiders, Centipedes, and Boxelder Bugs, add 3 to 4 tablespoons to 1 quart water. Thoroughly spray infested cracks and crevices, baseboards, behind shelving and bookcases for Silverfish, storage area and water drains in basements. For Ants, spray ant trails. For Ants and Boxelder Bugs, treat window frames, door sills, openings around water pipes, electrical outlets, and other places where they may enter. For Flies, Mosquitoes, and Wasps, mix at same rate and spray screens, doors, door sills, window frames, and areas frequented by these pests, such as porches and refuse disposal areas. Repeat application as needed for control of any insect listed.

**LAWN INSECTS:** For following insects add the designated number of tablespoons to 2 gallons water for 1,000 square feet. For Ants, Army worms, Crickets, Grasshoppers, and Spiders, use 1 1/2 tablespoons. Spray lawn area and ornamentals as needed. For Ants, treat hills and trails. For Chiggers, Earwigs, and Ticks, use 3 tablespoons and thoroughly treat turf and ornamentals. For Chinch Bugs and Sod Webworms, use 4 1/2 tablespoons and treat as for Chiggers. For White Grubs and Wireworms use 8 tablespoons. Wash into soil immediately after application. Birds feeding on treated areas may be harmed. Keep children and pets off treated areas until insecticide is washed into the soil and the grass is thoroughly dried.


**TERMITE CONTROL:** Add 1 quart to 12 gallons water. Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations dig a trench to, but not below, the footing. Then, as the trench is refilled, treat the soil at a rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult the State Extension Service for advice.

**NOTICE** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 2217-B4 -A

100 815 673



**45% CHLORDANE**  
**EMULSIFIABLE**

MANUFACTURED BY  
**GORDON CORPORATION**  
KANSAS CITY, KANSAS 66118

UNDER THE FEDERAL INSECTICIDE ACT PURCHASE ONLY IN STATES WHERE ACT IS IN FULL FORCE AND EFFECT

JUL 3 1973

**VEGETABLE GARDENS:** Soil treatment against maggots, sowbugs, and wireworms, use. Apply at time of seeding of corn and cabbage, tomatoes, and peppers. Do not apply to soil to be planted to root crops.

**FOR FOLIAGE SPRAYING ON THE FOLIAGE:** 2 TABLESPOONS PER GALLON WATER:

**BEANS:** To control flea beetles after pods begin to form.

**CABBAGE AND CAULIFLOWER:** To control cabbage worm, diamondback moth. Do not apply after head begins to form.

**DRY ONIONS:** For thrip control.

**POTATOES:** Flea beetles and blister beetles. Repeat as necessary.

**GRASSHOPPERS:** Use 2 tablespoons per gallon of water around the garden border. Open blossoms may be stunted.

**LEAF MINERS:** Use 4 tablespoons per gallon of water in late May or at first sign of infestation.

**WARNING:** Harmful if swallowed. Do not use on skin. Contact with skin can cause toxic symptoms. Wash with soap and water. Avoid contact with eyes. Keep out of reach of children. Do not use in areas until spray has been completely dried.

**DO NOT USE, POUR, SPILL, OR STORE**

This product is toxic to fish, birds, and mammals. Do not use in lakes, streams, and ponds. Birds fed with contaminated water by drinking from wastes. Do not apply where runoff is likely.

This product is highly toxic to bees and other beneficial insects. Do not use on crops. Protective information. Cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.



# 45% CHLORDANE EMULSIFIABLE



MANUFACTURED BY  
**GORDON CORPORATION**  
KANSAS CITY, KANSAS 64118

UNION PACIFIC CHEMICALS DIVISION  
KANSAS CITY, MISSOURI

JUL 13 1973

8

nts. Silverfish, Waterbugs, Bugs, add 3 to 4 table-  
spoons per gallon of water.  
For Silverfish, storage area  
spray ant trails. For Ants  
per sills, openings around  
doors where they may enter.  
Repeat application as

the designated number of  
tablespoons per gallon of  
water. For Ants, Army-  
worms, Fleas, Flea Beetles,  
and Flea Larvae, treat hills and  
yards with 3 tablespoons and thor-  
ax. For Flea Beetles and Sod Web-  
worms, use 3 tablespoons per gallon.  
Wash into soil immediately  
after application. Areas may be harmed.  
If insecticide is washed  
away.

ions water. Dig a trench  
in brick or hollow block  
at not below, the footing.  
at a rate of 4 gallons  
trench 3 feet deep would

carrier which will prevent  
require treatment. It is  
in Service for advice.

conforms to the ingredient  
list, such as weather, com-  
position of application equipment,  
these stated on this label.

100 815-673

**VEGETABLE GARDENS:** Soil treatment to control cutworms, seed corn  
cutworms, sowbugs, and wireworms, use 4 tablespoons per gallon water.  
Apply at time of seeding of corn and beans or around transplants of  
cabbage, tomatoes, and peppers. Do not apply to green or spring  
onions or soil to be planted to root crops.

**FOR FOLIAGE SPRAYING ON THE FOLLOWING CROPS USE 2 TABLE-  
SPOONS PER GALLON WATER.**

**BEANS:** To control flea beetles and green stink bugs. Do not apply  
after pods begin to form.

**CABBAGE AND CAULIFLOWER:** To control flea beetles, imported  
cabbageworm, diamondback moth larvae, and Harlequin bugs.  
Do not apply after head begins to form.

**DRY ONIONS:** For thrip control. Do not apply to green onions.

**POTATOES:** Flea beetles and blister beetles. Make application when  
insects appear. Repeat as necessary.

**GRASSHOPPERS:** Use 2 tablespoons per gallon. Spray grass and  
weeds around the garden border. Apply to ornamentals and flowers.  
Open blossoms may be stained.

**LEAF MINERS:** Use 4 tablespoons per gallon. Spray birches and holly  
in late May or at first sign of infestation. Repeat in 10 to 14 days.

**WARNING:** Harmful if swallowed. Do not breathe spray mist. Contact  
with skin can cause toxic symptoms. In case of contact with skin, wash  
with soap and water. Avoid contamination of feed and foodstuffs.  
Keep out of reach of children. Do not allow children or pets on treated  
areas until spray has been completely dried. Do not treat forage for  
animals.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME**

This product is toxic to fish, birds, and other wildlife. Keep out of  
lakes, streams, and ponds. Birds feeding on treated areas may be  
killed. Do not contaminate water by cleaning of equipment, or disposal  
of waste. Do not apply where runoff is likely to occur.

This product is highly toxic to bees exposed to direct treatment or  
residues on crops. Protective information may be obtained from your  
Cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.

815-673

118

119