

LAWN WEED KILLER

(Amine Type)

PM23
2217-41
pg. 2 of 3

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid . . .	26.6%
INERT INGREDIENTS	73.4%
TOTAL	100.0%

THIS PRODUCT CONTAINS:

*2 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon
or 22.1%.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with
soap and water; for eyes, flush
with water for 15 minutes

See Side Panel for Precautionary Statements

8/10/90


NET CONTENTS 1 PINT

865/590 AP071067*

EPA REG. NO. 2217-41
EPA EST. NO. 2217-KS-1

NOTIFICATION
LABEL NOT REVIEWED
PER FR NOTICE 88-6
DATE AUG 13 1990

Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, KANSAS 66118



STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

pg. 2 of 3

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. When handling this product, wear chemical resistant gloves. Avoid inhalation of spray mist. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children and pets.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.


CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS:

Avoid spray drift to susceptible plants as this product may injure cotton, beans, cucumbers, grapes, roses, tomatoes, other vegetables, flowers, ornamental plants. Coarse sprays applied at low pressures are less likely to drift. Do not overdose. Do not use equipment used in applying this material to apply insecticides or fungicides to susceptible plants or seeds. Do not use on grass seed crops or on turf containing creeping Bentgrasses. Do not use on new grass seedings less than 2 months old. Clovers present in turf may be damaged. Do not use on St. Augustine grass, Dichondra, or Carpetgrass lawns.

GENERAL INFORMATION:

This product may be used for the control of many broadleaf annual weeds and certain perennial weeds growing in lawns and other turf and in vacant lots and fence rows. In general, this product is most effective when weeds are small and growth is active. It may be applied when the temperatures range from 65°F to 90°F. Results obtained above and below these temperatures will be considerably slower and more erratic. Do not apply if rain is expected within 4 hours. Application may be made with either power or hand type sprayers. Use with water only. Will not mix with oil. Since local



2393

weather and growing conditions may vary the action of this product, it should always be used according to recommendations of State Agricultural Extension Service.

LAWNS AND TURF: For control of susceptible or easy to kill weeds in lawns and turf, use 1 tablespoon of this product per gallon of water (1 gallon will cover approximately 1,000 square feet) and apply either as a spot or all over spray. For hard to kill or more resistant weeds use 2 tablespoons per gallon of water and apply as necessary for control.

VACANT LOTS AND WASTE AREAS: For controlling broadleaf weed growth in these areas use 2 tablespoons of this product per gallon of water and apply to thoroughly wet all weeds to be killed. Repeat application every 2 to 3 weeks as necessary to control new growth and any regrowth.

WEEDS CONTROLLED

Susceptible or easy to kill weeds that are generally killed with one application.

Beggarticks	Morningglory, Annual	Ragweed
Cocklebur	Mustard	Shepherdspurse
Dandelion	Peppergrass	Sunflower
Horseweed	Plantain	Wild buckwheat

Slightly resistant weeds that require repeated applications and stronger dosages for control.

Bindweed (field)	Pigweed	Wild carrot
Curly dock	Red sorrel	Wild garlic
Lambsquarter	Sour dock	Wild onion
Mexican sandbur		

Repeat treatment as necessary for control.

LIMITED WARRANTY. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

EXCLUSION OF OTHER WARRANTIES AND REMEDIES. Except where such disclaimers and exclusions are specifically prohibited by applicable law. THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty. BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

