


GORDON'S NET CONTENT ONE GALLON

SUPER BRUSH KILLER

SUPER BRUSH KILLER IS A MIXTURE OF 2,4-D AND 2,4,5-T WHICH CAN BE DISSOLVED IN FUEL OIL OR EMULSIFIED WITH WATER

Directions

FOLIAGE TREATMENT

Application can be made any time after the full leaf stage until three weeks before frost. Best results are obtained when the brush to be sprayed is under five feet tall. Brush over five feet tall may require repeat application or cutting, and stump treatment. Use in such areas as roadsides, pastures, along fence rows and power lines for control of such plants as are listed on this label.

DILUTION TABLE

SUPER BRUSH KILLER	Water
2 to 3 Tablespoons	1 Gal.
1½ to 2 Pints	25 Gal.
1½ to 2 Quarts	50 Gal.
3 to 4 Quarts	100 Gal.

Get all stems and leaves thoroughly as complete coverage is necessary for uniform kill. Amount of material required will depend upon the height and denseness of the brush. If properly applied one thorough application should control 75% or more of the susceptible growth. Where under some conditions repeat applications may be necessary to control regrowth, usually a single treatment per year is sufficient. If brush is dense some plants may not be covered so a second application may be necessary to control those missed, or any regrowth.

NOTICE: Please remember that this product conforms to the minimum standard on the label. Since conditions vary, such as weather, compatibility with other chemicals, and condition of application equipment, will vary. Do not make any claim other than that made on the label.

SUPER BRUSH KILLER

Net Content One Gallon
BRUSH AND WOODY PLANTS

SUPER BRUSH KILLER

SINCE LOCAL WEATHER AND GROWING CONDITIONS MAY VARY AND AFFECT THE ACTION OF THIS PRODUCT, IT SHOULD ALWAYS BE USED ACCORDING TO RECOMMENDATIONS OF STATE AGRICULTURAL EXPERIMENTAL STATIONS OR EXTENSION SERVICE WEED SPECIALISTS.

Directions

STUMP AND TRUNK TREATMENT

Dilute at rate of ½ pint per gallon of fuel oil. Apply by spraying or brushing over the entire stump or from 18 inches to ground on trunks. Make application heavy until run off is evident on entire area and on all sides. Application may be made to stumps and trunks any time of year. **THOROUGH COVERAGE IS IMPORTANT.** Any new seedling or regrowth which may appear should be treated with Super Brush Killer diluted with water as shown in dilution table.

SOME SPECIES OF BRUSH CONTROLLED

Easy to Kill

- | | |
|---------------------------|--------------------------|
| ALDER, SPECKLED | HAZEL NUT |
| BLACKBERRY | HONEYSUCKLE |
| BUCKBRUSH | IVY (Poison) |
| CHERRY (Wild, Pin) | JUNEBERRY |
| CURRANT (Prickly, Sticky) | OSAGE ORANGE (Hedge) |
| DEWBERRY | PLUM (Sand) |
| ELM (American, Winged) | SUMAC (Smooth, Staghorn) |
| GOOSEBERRY | TRUMPET VINE |
| GRAPE | WAHOO (Burning Bush) |
| | WILLOW (Black, Bright) |

More Tolerant

- | | |
|----------------|------------------------|
| ASH (White) | HICKORY |
| DOGWOOD (Red) | MAPLE (Soft, Sugar) |
| PITTERSWEET | OAK (Black Jack, Burr) |
| ROSE (Prairie) | Post, White) |
| | PERSIMMON |

CAUTION:

Avoid contact with eyes, skin or clothing. In case of contact, wash off promptly with plenty of water. Harmful if swallowed.

Avoid spray drift to susceptible plants as this product may injure cotton, sugar beets, cucumbers, tomatoes, or certain ornamentals. Coarse sprays are less likely to drift. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not store near fertilizers, seeds, insecticides or fungicides. Do not use weed control equipment for applying insecticides or fungicides. Do not contaminate irrigation ditches or water used for domestic purposes. Rinse spray equipment and containers and dispose of liquid wastes in a pit or non-crop land located away from water supplies. **DON'T OVER DOSE. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

Do not graze dairy animals in treated areas within 6 weeks after application. Do not graze pastures in treated areas within 2 weeks of slaughter.

Do not use around the home or recreation areas. Do not reuse empty container. Destroy by crushing or crushing. Bury or discard in a safe place.

This product is toxic to fish. Keep out of water. Do not use near water. Do not use near water.

 **MANUFACTURED BY**
GORDON CORPORATION
 KANSAS CITY, KANSAS

**BUTYL ESTER 400
2,4,5-T
BRUSH KILLER**

This product is soluble in oil or emulsifiable in water. The 2,4,5-T Esters are generally more effective than 2,4-D or mixtures of 2,4-D and 2,4,5-T Esters where the stands of brush contain any considerable amounts of tolerant or harder to kill species.

DIRECTIONS

FOLIAGE TREATMENT

Apply to foliage of brush and woody plants.

Apply to foliage of brush and woody plants.

WATER

1 Gal
5 Gal
50 Gal
100 Gal

Wet all stems and leaves thoroughly as complete coverage is necessary for uniform kill. Amount of material required will depend upon the height and denseness of the brush. Usually an acre will require close to 100 gallons of diluted spray. If properly applied one thorough application should control 75% or more of the susceptible growth. While under some conditions repeat applications may be necessary to control regrowth, usually a single treatment per year is sufficient. If brush is dense some plants may not be covered so a second application may be necessary to control those missed or any regrowth.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 2217-100-AA
903 473

**BUTYL ESTER 400
2,4,5-T
BRUSH KILLER**

**A Concentrated Herbicide Especially Compounded
for Use on Both Brush and Woody Plants**

CONTAINS 4 POUNDS 2,4,5-T ACID PER GALLON

ACTIVE INGREDIENTS:
Butyl Ester of 2,4,5-Trichlorophenoxyacetic acid*..... 52.7%
INERT INGREDIENTS..... 47.3%
TOTAL.....100.0%
*2,4,5-Trichlorophenoxyacetic acid equivalent..... 43.2%

CONTROLS:

OAK WILD BLACKBERRY
HICKORY ELM
OSAGE ORANGE (Hedge)

CAUTION: Keep Out of Reach of Children
See Side Panels for Additional Cautions

See right panel for other Woody plants controlled by 2,4,5-T

NET CONTENTS GALLONS

903-473

**BUTYL ESTER 400 2,4,5-T
BRUSH KILLER**

Since local weather and growing conditions may vary and affect the action of this product, it should always be used according to recommendations of State Extension Service Weed Specialists.

DIRECTIONS

STUMP AND TRUNK TREATMENT

Dilute at rate of 1/2 pint per gallon of fuel oil. Apply by spraying or brushing over the entire stump or from 18 inches to ground on

FIGURE 1
SOME SPECIES OF LEAFY BRUSH CONTROLLED BY BUTYL ESTER 400 2,4,5-T BRUSH KILLER

label. Do not apply when weather conditions favor drift or spray from areas treated. Avoid spray drift to susceptible plants as this product may injure cotton, sugar beets, clover, tomatoes, or certain ornamentals. Coarse sprays are less likely to drift. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not store near fertilizers, seeds, insecticides or fungicides. Do not use weed control equipment for applying insecticides or fungicides. Avoid contamination of water used for irrigation purposes. **DON'T OVER DOSE.** Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter. This product will kill fish. Keep out of any body of water. Do not use around the home or recreation areas. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes. 903-473