



50% METHOXYCHLOR WETTABLE POWDER

Active Ingredient
Methoxychlor 50%
Inert Ingredients 50%

Net
Wt.
100%

Contains 50% Methoxychlor
50% Inert Ingredients
Net Wt. 100%

FOR USE ON
DAIRY CATTLE
BEEF CATTLE - DAIRY BARNS
VEGETABLE CROPS

CAUTION Keep Out of Reach of Children

539-750

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

NET WEIGHT 100 LBS

293

50% METHOXYCHLOR WETTABLE POWDER

DIRECTIONS FOR USE

HORN FLIES ON DAIRY ANIMALS: Dust directly on poll, back and tail approximately 1 heaping tablespoonful of this product. Rub into hair to give even coverage. Repeat every three weeks if needed. Do not spray dairy animals with this product.

BEEF CATTLE LICE, HOG LICE AND SHEEP TICKS: Use 1 pound in 12 gallons of water, covering animals thoroughly. Use 2 quarts of the mixture per hog or sheep and 4 quarts on cattle. Repeat application if necessary after 15 days.

DAIRY BARN, ANIMAL PENS: To control house flies and and stable flies use 1# in three gallons of water, applying to all surfaces where flies rest or crawl. Use 1 to 1½ gallons per 500 to 1,000 square feet, depending upon the surfaces. Repeat applications when flies reappear. Do not contaminate milk, feed, or drinking water. Exclude dairy animals while treating their areas.

VEGETABLES: For control of insects listed below on the respective crops use 2 tablespoons per gallon of water (2 to 3# per 100 gallons of water.) Thoroughly spray plants. For commercial gardens apply 100-150 gallons of spray mix per acre. On small areas treat with a hand pressure sprayer covering all leaves, stems and fruit. Start applications at first sign of insects and repeat at 7 to 10 day intervals as required for control. Do not apply to food crops later than 7 days before harvest unless other wise indicated below.

ASPARAGUS	— Asparagus Beetle
BEANS	— Bean Leaf Beetle 3 days Mexican Bean Beetle 3 days
CABBAGE	— Imported Cabbage Worm 3 days Flea Beetles 3 days
CUCUMBERS	— Cucumber Beetles
MELONS	— Melon Worms
PEPPERS	— Flea Beetles
POTATOES	— Blister Beetles Colorado Potato Beetle
TOMATOES	— Blister Beetles Tomato Fruit Worms Flea Beetles

CAUTION: Remove spray deposits from plant products consumed as food. Keep out of any body of water. Rinse spray equipment and dispose of waste and container by burying at least 18 inches deep. This product is toxic to fish.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

539-669

U.S.D.A. Reg. No. 2217-129

29

3 of 3

ACCEPTED
2217-129
JUN 14 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO *2217-129* SUBJECT
TO ATTACHED COMMENTS.