



# 50% METHOXYCHLOR WETTABLE POWDER

ACCEPTED

Jan 13, 1972

UNDER THE CODE  
FOR THE  
UNDER NO. 2217-129

MANUFACTURED BY  
**PBI - GORDON CORPORATION**  
KANSAS CITY, KANSAS

## 50% METHOXYCHLOR WETTABLE POWDER

### DIRECTIONS FOR USE

**HORN FLIES ON DAIRY ANIMALS:** Dust directly on poll, back and tail approximately 1 heaping tablespoonful of this product. Rub into hair to give even coverage. Repeat every three weeks if needed. Do not spray dairy animals with this product.

**BEEF CATTLE LICE, HOG LICE AND SHEEP TICKS:** Use 1 pound in 12 gallons of water, covering animals thoroughly. Use 2 quarts of the mixture per hog or sheep and 4 quarts on cattle. Repeat application if necessary after 15 days.

**DAIRY BARN, ANIMAL PENS:** To control house flies and and stable flies use 1# in three gallons of water, applying to all surfaces where flies rest or crawl. Use 1 to 1½ gallons per 500 to 1,000 square feet, depending upon the surfaces. Repeat applications when flies reappear. Do not contaminate milk, feed, or drinking water. Exclude dairy animals while treating their areas.

**VEGETABLES:** For control of insects listed below on the respective crops use 2 tablespoons per gallon of water (2 to 3# per 100 gallons of water.) Thoroughly spray plants. For commercial gardens apply 100-150 gallons of spray mix per acre. On small areas treat with a hand pressure sprayer covering all leaves, stems and fruit. Start applications at first sign of insects and repeat at 7 to 10 day intervals as required for control. Do not apply to food crops later than 7 days before harvest unless other wise indicated below.

|           |   |
|-----------|---|
| ASPARAGUS | — Asparagus Beetle                                      |
| BEANS     | — Bean Leaf Beetle 3 days<br>Mexican Bean Beetle 3 days |
| CABBAGE   | — Imported Cabbage Worm 3 days<br>Flea Beetles 3 days   |
| CUCUMBERS | — Cucumber Beetles                                      |
| MELONS    | — Melon Worms   |
| PEPPERS   | — Flea Beetles  |
| POTATOES  | — Blister Beetles<br>Colorado Potato Beetle             |
| TOMATOES  | — Blister Beetles<br>Tomato Fruit Worms<br>Flea Beetles |

**CAUTION:** Remove spray deposits from plant products consumed as food. Rinse spray equipment and dispose of waste and container by burying at least 18 inches deep.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

539-771

U.S.D.A. Reg. No. 2217-129

# 50% METHOXYCHLOR WETTABLE POWDER

## 50% METHOXYCHLOR WETTABLE POWDER

### DIRECTIONS FOR USE

**HORN FLIES ON DAIRY ANIMALS:** Dust directly on poll, back and tail approximately 1 heaping tablespoonful of this product. Rub into hair to give even coverage. Repeat every three weeks if needed. Do not spray dairy animals with this product.

**BEEF CATTLE LICE, HOG LICE AND SHEEP TICKS:** Use 1 pound in 12 gallons of water, covering animals thoroughly. Use 2 quarts of the mixture per hog or sheep and 4 quarts on cattle. Repeat application if necessary after 15 days.

**DAIRY BARNS, ANIMAL PENS:** To control house flies and and stable flies use 1# in three gallons of water, applying to all surfaces where flies rest or crawl. Use 1 to 1½ gallons per 500 to 1,000 square feet, depending upon the surfaces. Repeat applications when flies reappear. Do not contaminate milk, feed, or drinking water. Exclude dairy animals while treating their areas.

**VEGETABLES:** For control of insects listed below on the respective crops use 2 tablespoons per gallon of water (2 to 3# per 100 gallons of water). Thoroughly spray plants. For commercial gardens apply 100-150 gallons of spray mix per acre. On small areas treat with a hand pressure sprayer covering all leaves, stems and fruit. Start applications at first sign of insects and repeat at 7 to 10 day intervals as required for control. Do not apply to food crops later than 7 days before harvest unless otherwise indicated below.

|           |   |
|-----------|---|
| ASPARAGUS | Asparagus Beetle                                      |
| BEANS     | Bean Leaf Beetle 3 days<br>Mexican Bean Beetle 3 days |
| CABBAGE   | Imported Cabbage Worm 3 days<br>Flea Beetles 3 days   |
| CUCUMBERS | Cucumber Beetles                                      |
| MELONS    | Melon Worms   |
| PEPPERS   | Flea Beetles  |
| POTATOES  | Blister Beetles<br>Colorado Potato Beetle             |
| TOMATOES  | Blister Beetles<br>Tomato Fruit Worms<br>Flea Beetles |

**CAUTION:** Remove spray deposits from plant material as soon as food. Rinse spray equipment and dispose of waste and material by burying at least 18 inches deep.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, storage, weather, compatibility with other chemicals, and condition of application equipment will vary, seller assumes no liability for results not obtained on the above crops.

© 1964, FMC Corporation, P.O. Box 100, Philadelphia, Pa. 19106

