

11 22 1990 194

# 50% METHOXYCHLOR WETTABLE POWDER

FOR USE ON  
DAIRY CATTLE -- BEEF CATTLE -- DAIRY BARNS  
VEGETABLE CROPS

ACTIVE INGREDIENT:

*Methoxychlor Technical . . . . .	50%
INERT INGREDIENT . . . . .	50%
TOTAL	100%

\*Equivalent to 44% 1,1,1-trichloro-2,2-bis(p-methoxyphenyl) ethane and 6% other isomers and related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautionary Statements


NET WEIGHT FOUR POUNDS

539/1089 AP

EPA REG NO 2217-129

EPA EST. NO 2217-KS-1

Mfd by PBI/GORDON CORPORATION  
KANSAS CITY, KS 66118



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist.

### **Statement of Practical Treatment**

**IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

**IF INHALED:** Remove victim to fresh air and apply respiration as indicated.

**IF ON SKIN:** Wash promptly with soap and water. Rinse thoroughly.

**IF IN EYES:** Flush with plenty of water and get medical attention immediately.

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

### **Physical or Chemical Hazard**

Do not use or store near heat or open flame.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

## **STORAGE & DISPOSAL**

**STORAGE:** Store in original container in a locked storage area inaccessible to children or pets.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Species	Pest	Dosage Rate	Application Interval	Directions and Limitations
Nonlactating Dairy Animals	Horn Flies and Lice	1 tablespoon per adult animal	21 days	Animal dust treatment. Dilute with talc powder (1:3 dilution). Apply to the neck, back and upper sides of young dairy cattle. Do not treat animals less than 30 days of age. Do not apply during lactation. Observe 2 week pre-freshening interval.
Beef Cattle	Lice Ticks	1.5-2.0 lbs per 50 gal/water	15 days if needed	Animal spray or dip treatment. Apply as a spray or as a dip. When applied as a spray use 2 quarts of finished spray per head of large animals and proportionally less for smaller animals. When applied as a dip force animal to walk through a tank full of solution sufficiently deep to thoroughly wet animal. For longnosed cattle louse and short-nosed cattle louse, apply as a drenching spray. Do not spray, dip, or treat calves under 30 days of age.
Hogs	Lice (including hog louse)	1.5-2.0 lbs per 50 gal/water	15 days if needed	Animal spray or dip treatment. When applied as a spray, use 2 quarts of finished spray per head of large animals and proportionally less for smaller animals. When applied as a dip, force animal to walk through a tank full of solution sufficiently deep to thoroughly wet animals. For hog louse, apply as a drenching spray. Do not treat animals less than 1 month of age.
Sheep	Ticks	20 lbs per 50 gal/water	15 days if needed	Animal spray or dip treatment. When applied as a spray, use 2 quarts of finished spray, per head of large animals and proportionally less for smaller animals. When applied as a dip, force animal to walk through a tank full of solution sufficiently deep to thoroughly wet animals. Observe a 14 day freshening interval.

4/4/57

Site	Pest	Dosage Rate	Application Interval	Directions and Limitations
Livestock Buildings and Premises				
	Stable Fly	5% solution	—	Indoor premise treatment. Apply to wall and floor
	House Fly	applied at		surfaces inside the building, including cracks,
		1-2 gals. per		crevices, baseboards, and animal quarters such
		1000 sq. ft.		as stalls, pens and bedding. Spray formulations
Specific sites include: animal shelters,				may be applied as a residual contact or space
barns (including dairy barns), milk				spray. Apply spray formulations by mist blower,
rooms, pens, sheds, and stalls				hydraulic sprayer, hand sprayer or cylindrical
				pressure sprayer. Supplement treatment with
				proper sanitary measures, including removal of
				breeding sources, such as manure in and around
				buildings. Cover securely all milk-handling
				equipment, feedstuffs, troughs, bins, or other
				food and water receptacles while spraying. Dairy
				animals should not be present while spraying.

Vegetables	Pests	Dosage Rates		Pre-harvest Interval	Directions and Limitations
		lbs. product 100 gallon	tbsps 1 gallon		
Asparagus	Asparagus Beetle	1.0 - 1.5	2-3	7	Foliar application. Apply when pests first appear and repeat at 7 to 10 day intervals or as needed.
Beans	Beanleaf Beetle	1.0 - 1.5	2-3	3	Foliar application. Repeat at 7 to 14 day intervals or as needed. Do not use vines as food or feed.
	Mexican Bean Beetle				
Cabbage	Imported Cabbageworm	1.0 - 1.5	2-3	7	Foliar application.
	Flea Beetles				
Cucumbers	Cucumber Beetles*	1.0 - 1.5	2-3	7	Foliar application.
	*(Including striped cucumber beetle and twelve-spotted cucumber beetle)				
Melons	Melons*	1.0 - 1.5	2-3	7	Foliar application.
	*(Including cantaloupe, casaba melon, crenshaw melon, honeyball melon, honeydew melon, musk melon, Persian melon, and watermelon)				
Peppers	Flea Beetles	1.0 - 1.5	2-3	7	Foliar application.
Potatoes	Blister Beetles	1.0 - 2.5	2-3	None	Foliar application.
	Colorado Potato Beetle				
Tomatoes	Blister Beetles	1.0 - 1.5	2-3	7	Foliar application.
	Flea Beetles				
	Tomato Fruit Worm				

**LIMITED WARRANTY.** Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

**EXCLUSION OF OTHER WARRANTIES AND REMEDIES.** Except where such disclaimers and exclusions are specifically prohibited by applicable law, THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.