

**METHOXYCHLOR
EMULSION CONCENTRATE**
Directions for Use
LIVESTOCK

Beef Cattle:

To control horn fly, mix 1 cup of this concentrate with 3 gallons of water, or 2 gallons of this concentrate with 100 gallons of water. Use about 2 quarts of spray per animal, thoroughly covering the animal. Repeat treatment in 2 to 3 weeks, or when flies reappear.

Cattle Lice, Hog Lice:

Mix 1 cup of this concentrate with 3 gallons of water or 2 gallons of this concentrate per 100 gallons of water. Apply about 2 quarts per hog and 4 quarts per beef animal, thoroughly covering the animal including bush of tail and inside of ear. Repeat treatment in 15 days.

FIELD CROPS

Alfalfa and Clover:

To control blister beetle, clover leaf weevil, leaf hoppers, alfalfa weevil and alfalfa caterpillar, spray at the first sign of infestation and repeat as needed. Use 3 quarts of this concentrate in sufficient water to cover one acre. Do not spray during bloom. After spraying allow 7 days waiting period before harvest.

CAUTION: Harmful if swallowed. Avoid inhalation and prolonged contact with skin. Wash with soap and water.

This product will kill fish. Keep out of any body of water. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of spray from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes. Remove spray deposits from plant products consumed as food. Do not store near fire or open flame.

897-467

Methoxychlor
Emulsion Concentrate

(Contains 2 lbs. Methoxychlor per gallon)

ACTIVE INGREDIENTS

Methoxychlor, Technical 24.0%
Aromatic Petroleum Hydrocarbon 72.5%

INERT INGREDIENTS 3.5%
100%

*Consists of 21% 2,2-bis(p-methoxyphenyl)-1,1,1 trichloroethane and 3% other isomers and related compounds.

EFFECTIVE FOR USE ON

**BEEF CATTLE FIELD CROPS
GRAIN BINS**

WHEN USED AS DIRECTED

CAUTION: Keep Out of Reach of Children
See Side Panels for Additional Cautions

Contents 55 Gallon

MANUFACTURED BY

GORDON CORPORATION
Kansas City, Kansas

**METHOXYCHLOR
EMULSION CONCENTRATE**
Directions for Use
BUILDINGS

Grain Bins:

To aid in the control of the exposed stages of cadelle, sawtoothed beetle and certain other insects attacking stored grain, mix 1 quart of this concentrate with 2½ gallons of water or 10 gallons of this concentrate with 100 gallons of water. Thoroughly clean the bin and apply about 2 gallons of spray per 1000 square feet. Application should be made about two weeks before grain is placed in the bin.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

WARNING: This product will tend to settle out under certain temperature conditions. **Keep this material stored above 40° F.** If it settles, warm to 80°F. and agitate thoroughly before using. The properties are not affected.

897-467

U.S.D.A. Reg. No. 2217-131

2217-131

12/21/1971

1/2

EPA-PA

10-10-71

ACCEPTED

DEC 21 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

2217-131

2217-131