



2 quarts of spray per animal, thoroughly covering the animal. Repeat treatment in 2 to 3 weeks, or when flies reappear.

Cattle Lice, Hog Lice:

Mix 1 cup of this concentrate with 3 gallons of water or 2 gallons of this concentrate per 100 gallons of water. Apply about 2 quarts per hog and 4 quarts per beef animal, thoroughly covering the animal including bush of tail and inside of ear. Repeat treatment in 15 days.

FIELD CROPS

Alfalfa and Clover:

To control blister beetle, clover leaf weevil, leaf hoppers, alfalfa weevil and alfalfa caterpillar, spray at the first sign of infestation and repeat as needed. Use

METHOXYCHLOR

EMULSIFIABLE

2217-131

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

ENVIRONMENTAL WARNINGS:
This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Bees: This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not apply when weather conditions favor drift of spray from areas treated.

Apply this product only as specified on this label.

CONTAINER DISPOSAL: Rinse equipment and containers and dis-

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

Directions for Use **LIVESTOCK**

Beef Cattle:

To control lice, mix 1 cup of this concentrate with 3 gallons of water, or 2 gallons of this concentrate with 100 gallons of water. Use about 2 quarts of spray per animal, thoroughly covering the animal. Repeat treatment in 10 to 15 days or when flies appear.

Cattle Lice (Dog Lice)

Apply 1 quart of this concentrate to the animal's body. Apply 4 quarts per animal, covering the entire body and inside of the ears. Repeat in 10 to 15 days.

FIELD CROPS

Grain and Cereals

For control of insects on grain, beat the grain and apply this concentrate at the first sign of infestation. Use 3 quarts of this concentrate in sufficient water to cover one acre. Do not spray during bloom. After spraying allow 7 days waiting period before harvest.

BUILDINGS

Grain Bins:

To aid in the control of the exposed stages of cadelle, sawtoothed beetle and certain other insects attacking stored grain, mix 1 quart of this concentrate with 2½ gallons of water or 10 gallons of this concentrate with 100 gallons of water. Thoroughly clean the bin and apply about 2 gallons of spray per 1000 square feet. Application should be made about two weeks before grain is placed in the bin.

97-78

E.P.A. Reg. No. 221-131-AA

METHOXYCHLOR **EMULSIFIABLE**

TO ATTACHED COMMENTS.

(Contains 3 lbs. Methoxychlor* per gallon)

Active Ingredients	24.0%
Aromatic Petroleum Hydrocarbon	72.5%
INERT INGREDIENTS	3.5%
	100%

*Consists of 21% 2,2-bis(p-methoxyphenyl)-1,1,1 trichloroethane and 3% other isomers and related compounds.

EFFECTIVE FOR USE ON

BEEF CATTLE FIELD CROPS
GRAIN BINS

WHEN USED AS DIRECTED

WARNING: Keep out of Reach of Children

See Side Panel for Additional Warnings

CONTENTS: ONE GALLON

HUMAN WARNINGS: Harmful if swallowed. Avoid inhalation and contact with skin. Wash with soap and water.

Remove spray deposits from plant products consumed as food.

DO NOT USE FOR SPRAYING OF FOLIAGE OR FRUIT OF PLANTS.

ENVIRONMENTAL WARNINGS

CONTAINER DISPOSAL

pose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

WARNING: This product will tend to settle out under certain temperature conditions. Keep this material stored above 40° F. If it settles, warm to 80°F. and agitate thoroughly before using. The properties are not affected.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

897-878