

FOR USE ON	TO CONTROL	IN 100 GAL.	IN 1 GAL.	COMMENTS
Apples, Pears, Cherry, Blue, Tree Sprayed with	European Red Mite 1-2 pts. 2 tsp.			Make two or three applications at 7-day intervals. Do not use for mites, use or more for aphids. At pre-bloom or post-bloom of blossoms or when mites appear. Do not use within 3 days of harvest.
Wash Apple Aphid, Pear Aphid, Green Apple Aphid 1 1/2 pt. 1 1/2 tsp.				
Tomatoes	Spider, White, Aphid 1 1/2 pt. 1 1/2 tsp.			Make one or more applications as needed. Do not use within 3 days of harvest.
Beans	Black Bean Beetle, Spider Mite, Leafminer 2 pts. 2 tsp.			
Peanut	Pea Aphid 2 pts. 2 tsp.			Make one or more applications. Do not use within 3 days of harvest. Do not use on peas vines for storage.
Potatoes	Aphid 2 pts. 2 tsp.			
Cabbage, Broccoli, Kale, Mustard, Turnip	Aphid, Imported Cabbage Worm 2 pts. 2 tsp.			Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on Broccoli and Turnip. 7 days on others.
Alfalfa for Seed	Aphid 2 pts. per acre to at least 10 gal. water			Do not apply within 7 days of harvest, or when in bloom.
Flowers, Ornamentals	Spider Shell Scale, European Scale 1 pt. 1 tsp.			Apply in summer when infestation occurs.



57% MALATHION

Emulsifiable Concentrate

FOR CONTROL OF ANIMAL AND PLANT PESTS

ACTIVE INGREDIENTS:
 Malathion* 57%
 Xylene 34%
 INERT INGREDIENTS 9%
 100%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate. Contains 5 pounds Malathion per gallon.
WARNING: Keep Out of Reach of Children
 See Side Panels for Additional Warnings
 934-1067 NET CONTENTS ONE GALLON



MANUFACTURED BY
GORDON CORPORATION
 KANSAS CITY, KANSAS

DIRECTIONS FOR USE

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots, Poultry houses, Utility buildings and around food disposals mix 5 ounces with 5 ounces of corn syrup and add to 2 gallons of water. Paint or spray on desired surfaces.

POULTRY: To control Chicken Lice add 1 oz. to 1 gallon water and apply directly to the birds. To spray walls, nests and litter make a solution of 2 oz. in 1 gallon water. Do not apply this higher dosage to birds. As a roost paint for lice control mix 2 ounces of 57% Malathion in one quart of water and apply at the rate of one pint of finished spray per 100 feet of roost.

SHEEP CATTLE: For control of Lice, add 1 gallon to 100 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed. For Control of ticks, apply 1 to 2 gallons per 100 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed.

SWINE: For control of Lice, apply 1 gallon per 100 gallons of water. Apply a complete spray to animals, pens and litter. One treatment may be sufficient. Repeat application only if needed.

BACK RUBBING DEVICES: For the reduction of Lice on Beef Cattle apply a 2% mixture (1 pint in 4.5 gal. of fuel oil or kerosene) to the back rubbing device. There may also be a reduction in Horn Flies. Make rubbing device accessible one to 35 or 45 head of cattle. Retreat the oiler every 2 or 3 weeks. DO NOT MAKE ACCESSIBLE TO DAIRY ANIMALS.

CAUTION: Do not apply to dairy animals. Do not treat calves under one month of age. Do not contaminate food containers or watering devices. Do not use where milk is processed or stored.

Our Name Is Your
 Guarantee
 of
 Quality

2017-143

94-69

1042

2 of 2

A 10x10 grid of dots forming a stylized letter 'A'. The top row has 10 dots. The second row has 9 dots. The third row has 8 dots. The fourth row has 7 dots. The fifth row has 6 dots. The sixth row has 5 dots. The seventh row has 4 dots. The eighth row has 3 dots. The ninth row has 2 dots. The tenth row has 1 dot.

ACCEPTED
SEP 4 1969
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 8217-143