

FOR USE ON	TO CONTROL	IN 100 GAL.	IN 1 GAL.	COMMENTS
Apples, Pear,	European Red Sawfly 1-2 pt.	2 qt.		Make tea of three applications at end of the season, one or more for apples. At pre-bloom, or just after of flower or when adults appear. Do not use within 3 days of harvest.
	Two Spotted Wasp			
	Warty Apple Aphid 1 pt.	1 qt.		
	Parasitic Scales			
	Green Apple Aphid 1 1/2 pt.	1 1/2 qt.		
	Red Apple Aphid			
Tomatoes	Spider Mites, Aphids	1 1/2 pt. 1 1/2 qt.	1 pt. 1 qt.	Make tea of three applications at end of the season, one or more for apples. At pre-bloom, or just after of flower or when adults appear. Do not use within 3 days of harvest.
Beans	Mexican Bean Beetle, Spider Mites, Leafhopper	2 pt. 2 qt.		
Pean	Pea Aphid	2 pt. 2 qt.		Make tea of three applications at end of the season, one or more for apples. At pre-bloom, or just after of flower or when adults appear. Do not use within 3 days of harvest.
Prunus	Aphid	2 pt. 2 qt.		
Cabbages, Broccoli, Kale, Mustard, Turnip	Aphids, Imported Cabbages Worm	2 pt. 2 qt.		Make tea of three applications at end of the season, one or more for apples. At pre-bloom, or just after of flower or when adults appear. Do not use within 3 days of harvest.
Alfalfa for Seed	Aphids	2 pt. per acre at least 10 gal. water.		Do not spray within 7 days of harvest, or when in bloom.
Flowers, Ornamentals	Orchard Small Scale, European Scale	1 pt. 1 qt.		Apply to the first when infestation is seen.



57% MALATHION

Emulsifiable Concentrate
FOR CONTROL OF
ANIMAL AND PLANT PESTS

ACTIVE INGREDIENTS:
Malathion* 57%
Xylene 34%
INERT INGREDIENTS 9%
100%

*O,O-dimethyl dithiophosphate of diethyl mercapto-succinate. Contains 5 pounds Malathion per gallon.
WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings

NET CONTENTS ONE GALLON

MANUFACTURED BY
GORDON CORPORATION
KANSAS CITY, KANSAS

DIRECTIONS FOR USE

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots, Poultry houses, Utility buildings and around food disposals mix 5 ounces with 5 ounces of corn syrup and add to 3 gallons of water. Paint or spray on desired surfaces.

POULTRY: To control Chicken Lice add 1 oz. to 1 gallon water and apply directly to the birds. To spray walls, nests and litter make a solution of 2 oz. in 1 gallon water. Do not apply this higher dosage to birds. As a roost bait for lice control stir 3 ounces of 57% Malathion in one quart of water and apply at the rate of one pint of finished spray per 150 feet of roost.

BEEF CATTLE: For control of Lice, add 1 gallon to 100 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed. For Control of ticks, apply 1 to 2 gallons per 100 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed.

SWINE: For control of Lice, apply 1 gallon per 100 gallons of water. Apply a complete spray to animals, pens and litter. One treatment may be sufficient. Repeat application only if needed.

BACK RUBBING DEVICES: For the reduction of Lice on Beef Cattle apply a 2% mixture (1 pint in 4.3 gals. of fuel oil or kerosene) to the back rubbing device. There may also be a reduction in Horn Flies. Make rubbing device accessible one to 35 or 45 head of cattle. Retreat the other every 2 or 3 weeks. DO NOT MAKE ACCESSIBLE TO DAIRY ANIMALS.

CAUTION: Do not apply to dairy animals. Do not treat calves under one month of age. Do not contaminate food containers or watering devices. Do not use where milk is

Our Name Is Your

Guarantee

of

Quality

2217-143

1-14-72

1042

Figure 1 shows a 10x10 grid of dots. The dots are arranged in a regular grid pattern, with some dots missing to form a shape resembling a large 'C' or a bracket on the right side. The dots are arranged in a regular grid pattern, with some dots missing to form a shape resembling a large 'C' or a bracket on the right side.

ACCEPTED
JAN 14 1972
THE FEDERAL INSECTICIDE ACT
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE FOR HILSON REGISTERED
FOR ECONOMIC HILSON REGISTERED
ED UNDER NO. 3807-15 SUBJECT
TO ATTACHED COMMENTS.