


FOR USE ON	TO CONTROL	IN 100 GAL.	IN 1 GAL.	COMMENTS
Apples, Pears,	European Red Mite	1-2 pts.	2 tsp.	Make two or more applications as needed for mites, one or more for aphids. At pre-pink or pink stage of bloom or when aphids appear. Do not use within 3 days of harvest.
	Clover Mite			
	Two Spotted Mite			
	Woolly Apple Aphid	1 pt.	1 tsp.	
	Forbes Scale			
	Green Apple Aphid	1½ pts.	1½ tsp.	
	Rosy Apple Aphid			
Tomatoes	Spider, Mites, Aphids	1½ pts.	1½ tsp.	Make one or more applications as needed. Do not use within 3 days of harvest.
Beans	Mexican Bean Beetle, Spider Mites, Leafhopper	2 pts.	2 tsp.	Do not use within 3 days of harvest.
Peas	Pea Aphid	2 pts.	2 tsp.	Make one or more applications. Do not use within 3 days of harvest. Do not use pea vines for forage.
Potatoes	Aphids	2 pts.	2 tsp.	Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on Broccoli and Turnips, 7 days on others.
Cabbage, Broccoli, Kale, Mustard, Turnip	Aphids Imported Cabbage Worm	2 pts.	2 tsp.	
Alfalfa for Seed	Aphids,	2 pts. per acre in at least 10 gal. water.		Do not apply within 7 days of harvest, or when in bloom.
Flowers, Ornamentals, Shrubs	Oyster Shell Scale	1 pt.	1 tsp.	Apply in summer when infestation occurs. Repeat in 10 to 12 days. For scales, apply when crawlers have settled on foliage.
	Euonymus Scale			
	Aphids,	1½ pts.	1½ tsp.	
	Spider Mites,			
	White Fly, Thrips,			
	Mealy Bugs,			
	Japanese Beetle Adult,			
	Four Lined Plant Bug,			
	Tarnished Plant Bug,			
	Rose Leafhopper			
	Potato Leafhopper			
	Bagworms, Juniper Scale	2 pts.	2 tsp.	
	Soft Brown Scale	2½ pts.	2½ tsp.	
	Pine Needle Scale	4 pts.	4 tsp.	

CAUTION: Do not use on certain ferns, including Boston maidenhair and Pteris, as well as some species of Crassula, because of possible injury to these ornamentals.



DIRECTIONS FOR USE

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots, Poultry houses, Utility buildings and around food disposals mix 5 ounces with 5 ounces of corn syrup and add to 2 gallons of water. Paint or spray on desired surfaces.

POULTRY: To control Chicken Lice add 1 oz. to 1 gallon water and apply directly to the birds. To spray walls, nests and litter make a solution of 2 oz. in 1 gallon water. Do not apply this higher dosage to birds. As a roost paint for lice control mix 2 ounces of 57% Malathion in one quart of water and apply at the rate of one pint of finished spray per 150 feet of roost.

BEEF CATTLE: For control of Lice, add 1 quart to 25 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed. For control of ticks 1 to 2 quarts per 25 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed.

SWINE: For control of Lice, apply 1 quart per 25 gallons of water. Apply a complete spray to animals, pens, and litter. One treatment may be sufficient. Repeat application only if needed.

BACK RUBBING DEVICES: For the reduction of Lice on Beef Cattle apply a 2% mixture (1 pint in 4.3 gals. of fuel oil or kerosene) to the back rubbing device. There may also be a reduction in Horn Flies. Make rubbing device accessible one to 35 or 45 head of cattle. Retreat the oiler every 2 or 3 weeks. **DO NOT MAKE ACCESSIBLE TO DAIRY ANIMALS.**

NOTE: Do not apply to dairy animals. Do not treat calves under one month of age. Do not contaminate food containers or watering devices. Do not use where milk is processed or stored.

HUMAN WARNINGS: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist, avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before reuse. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

Malathion may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL WARNINGS: This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. This product is highly toxic to bees exposed to direct treatment. Do not spray when bees are active in the area.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Do not reuse container. Destroy when empty.

Apply this product only as specified on this label.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

934-573

E.P.A. Reg. No. 2217-143-AA

2217-143

8-1-74

1081