

FOR USE ON	TO CONTROL	IN 100 GAL.	IN 1 GAL.	COMMENTS
<u>Apples,</u> <u>Pears,</u>	European Red Mite	1-2 pts.	2 tsp.	Make two or more applications as needed for mites, one or more for aphids. At pre-pink or pink stage of bloom or when aphids appear. Do not use within 3 days of harvest.
	Clover Mite			
	Two Spotted Mite			
	Woolly Apple Aphid	1 pt.	1 tsp.	
	Forbes Scale			
	Green Apple Aphid	1½ pts.	1½ tsp.	
	Rosy Apple Aphid			
<u>Tomatoes</u>	Spider Mites,	1½ pts.	1½ tsp.	Make one or more applications as needed. Do not use within 3 days of harvest.
	Aphids	1 pt.	1 tsp.	
<u>Beans</u>	Mexican Bean Beetle,	2 pts.	2 tsp.	
	Spider Mites,			
	Leafhopper			
<u>Peas</u>	Pea Aphid	2 pts.	2 tsp.	Make one or more applications. Do not use within 3 days of harvest. Do not use pea vines for forage.
<u>Potatoes</u>	Aphids	2 pts.	2 tsp.	
<u>Cabbage,</u> <u>Broccoli,</u> <u>Kale,</u> <u>Mustard,</u> <u>Turnip</u>	Aphids	2 pts.	2 tsp.	Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on Broccoli and Turnips, 7 days on others.
	Imported Cabbage			
	Worm			
<u>Alfalfa</u> <u>for Seed</u>	Aphids,	2 pts. per acre in at least 10 gal. water.		Do not apply within 7 days of harvest, or when in bloom.
<u>Flowers,</u> <u>Ornamentals,</u> <u>Shrubs</u>	Oyster Shell Scale	1 pt.	1 tsp.	Apply in summer when infestation occurs. Repeat in 10 to 12 days. For scales, apply when crawlers have settled on foliage.
	Euonymus Scale			
	Aphids,	1½ pts.	1½ tsp.	
	Spider Mites,			
	White Fly, Thrips,			
	Mealy Bugs,			
	Japanese Beetle Adult,			
	Four Lined Plant Bug,			
	Tarnished Plant Bug,			
	Rose Leafhopper			
	Potato Leafhopper			
	Bagworms, Juniper	2 pts.	2 tsp.	
	Scale			
	Soft Brown Scale	2½ pts.	2½ tsp.	
	Pine Needle Scale	4 pts.	4 tsp.	

**CAUTION:** Do not use on certain ferns, including Boston maidenhair and Pteris, as well as some species of Crassula, because of possible injury to these ornamentals.


E.P.A. Est. No. 2217-KS-1

934-974



# 57% MALATHION

## Emulsifiable



MANUFACTURED BY

**PBI - GORDON CORPORATION**

KANSAS CITY, KANSAS 66118

**LIQUID FLY**  
houses. Util  
corn syrup

**POULTRY:** T  
the birds.  
water. Do  
mix 2 ounce  
pint of finis

**DEER CATTLE**  
plate cover  
trol of ticks  
Repeat appl

**SWINE:** For  
spray to ani  
only if need

**BACK RUBBI**  
(1 pint in 4  
also be a r  
head of cat  
DAIRY ANIM

**NOTE:** Do ne  
not contami  
processed o

**HUMAN WAR**  
breathing sp  
with soap a  
contaminatio

Malathion ma  
aid. Atropin  
conjunction

**ENVIRONMEN**  
or ponds. Do  
Apply this p  
This product  
crops. Prote  
Extension Se

DO  
Do not reuse

**NOTICE:** Sell  
the label. Sin  
and condition  
those stated  
934-974

ACCEPTED WITH COMMENTS



# 57% MALATHION

## Emulsifiable

MANUFACTURED BY  
**PBI - GORDON CORPORATION**  
KANSAS CITY, KANSAS 66118

### DIRECTIONS FOR USE

**LIQUID FLY BAIT:** To control House and Stable Flies in Barns, Feed Lots, Poultry Houses, Utility buildings and around food disposals mix 5 ounces with 5 ounces of corn syrup and add to 2 gallons of water. Paint or spray on desired surfaces.

**POULTRY:** To control Chicken Lice add 1 oz. to 1 gallon water and apply directly to the birds. To spray walls, nests and litter make a solution of 2 oz. in 1 gallon water. Do not apply this higher dosage to birds. As a roost paint for lice control mix 2 ounces of 57% Malathion in one quart of water and apply at the rate of one pint of finished spray per 150 feet of roost.

**BEEF CATTLE:** For control of Lice, add 1 quart to 25 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed. For control of ticks 1 to 2 quarts per 25 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed.

**SWINE:** For control of Lice, apply 1 quart per 25 gallons of water. Apply a complete spray to animals, pens, and litter. One treatment may be sufficient. Repeat application only if needed.

**BACK RUBBING DEVICES:** For the reduction of Lice on Beef Cattle apply a 2% mixture (1 pint in 4.3 gals. of fuel oil or kerosene) to the back rubbing device. There may also be a reduction in Horn Flies. Make rubbing device accessible one to 35 or 45 head of cattle. Retreat the oiler every 2 or 3 weeks. **DO NOT MAKE ACCESSIBLE TO DAIRY ANIMALS.**

**NOTE:** Do not apply to dairy animals. Do not treat calves under one month of age. Do not contaminate food containers or watering devices. Do not use where milk is processed or stored.

**HUMAN WARNINGS:** Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist, avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before reuse. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

Malathion may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

**ENVIRONMENTAL WARNINGS:** This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME**

Do not reuse container. Destroy when empty.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

934-974

E.P.A. Reg. No. 2217-143-AA

TRU

n Re

Atte

apple

ca

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

apple

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

SA

For control of ticks, lice, and other insects on animals and plants. Use as directed. Do not use on cats, dogs, or birds. Do not use on ornamental plants. Do not use on food crops. Do not use on lawns. Do not use on trees. Do not use on shrubs. Do not use on flowers. Do not use on vegetables. Do not use on fruits. Do not use on nuts. Do not use on seeds. Do not use on grain. Do not use on hay. Do not use on straw. Do not use on manure. Do not use on urine. Do not use on blood. Do not use on milk. Do not use on eggs. Do not use on young animals. Do not use on pregnant animals. Do not use on nursing animals. Do not use on sick animals. Do not use on injured animals. Do not use on animals with open wounds. Do not use on animals with skin diseases. Do not use on animals with eye diseases. Do not use on animals with ear diseases. Do not use on animals with nose diseases. Do not use on animals with mouth diseases. Do not use on animals with throat diseases. Do not use on animals with lung diseases. Do not use on animals with heart diseases. Do not use on animals with kidney diseases. Do not use on animals with liver diseases. Do not use on animals with spleen diseases. Do not use on animals with pancreas diseases. Do not use on animals with stomach diseases. Do not use on animals with intestines diseases. Do not use on animals with rectum diseases. Do not use on animals with anus diseases. Do not use on animals with vagina diseases. Do not use on animals with penis diseases. Do not use on animals with testis diseases. Do not use on animals with ovary diseases. Do not use on animals with uterus diseases. Do not use on animals with fallopian tube diseases. Do not use on animals with cervix diseases. Do not use on animals with endometrium diseases. Do not use on animals with myometrium diseases. Do not use on animals with perimetrium diseases. Do not use on animals with decidua diseases. Do not use on animals with placenta diseases. Do not use on animals with umbilical cord diseases. Do not use on animals with navel diseases. Do not use on animals with skin diseases. Do not use on animals with eye diseases. Do not use on animals with ear diseases. Do not use on animals with nose diseases. Do not use on animals with mouth diseases. Do not use on animals with throat diseases. Do not use on animals with lung diseases. Do not use on animals with heart diseases. Do not use on animals with kidney diseases. Do not use on animals with liver diseases. Do not use on animals with spleen diseases. Do not use on animals with pancreas diseases. Do not use on animals with stomach diseases. Do not use on animals with intestines diseases. Do not use on animals with rectum diseases. Do not use on animals with anus diseases. Do not use on animals with vagina diseases. Do not use on animals with penis diseases. Do not use on animals with testis diseases. Do not use on animals with ovary diseases. Do not use on animals with uterus diseases. Do not use on animals with fallopian tube diseases. Do not use on animals with cervix diseases. Do not use on animals with endometrium diseases. Do not use on animals with myometrium diseases. Do not use on animals with perimetrium diseases. Do not use on animals with decidua diseases. Do not use on animals with placenta diseases. Do not use on animals with umbilical cord diseases. Do not use on animals with navel diseases.

# 57% MALATHION

## Emulsifiable

FOR CONTROL OF  
ANIMAL AND PLANT PESTS

ACTIVE INGREDIENTS:	
Malathion*	57%
Xylene	34%
INERT INGREDIENTS	9%
	100%
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate	
Contains 5 pounds Malathion per gallon.	

**WARNING: Keep out of Reach of Children**  
See Side Panels for Additional Warnings

934-974

NET CONTENTS ONE QUART

### DIRECTIONS

**LIQUID BAIT** For use on cats and dogs. Mix 1 quart of Malathion with 1 quart of water and 1 quart of corn syrup. Apply to the animal's fur.

**POULTRY** For use on chickens, turkeys, and ducks. Mix 1 quart of Malathion with 1 quart of water. Spray the birds. Do not allow the birds to drink the water. Do not allow the birds to eat the feed.

**BEEF CATTLE** For use on cattle. Mix 1 quart of Malathion with 1 quart of water. Spray the cattle. Do not allow the cattle to drink the water. Do not allow the cattle to eat the feed.

**SWINE** For use on pigs. Mix 1 quart of Malathion with 1 quart of water. Spray the pigs. Do not allow the pigs to drink the water. Do not allow the pigs to eat the feed.

**BACK RUBBING DEVICES** For use on animals. Mix 1 pint of Malathion with 1 pint of kerosene. Apply to the back of the animal.

**DAIRY ANIMALS** For use on dairy animals. Mix 1 quart of Malathion with 1 quart of water. Spray the animals. Do not allow the animals to drink the water. Do not allow the animals to eat the feed.

**NOTE** Do not use Malathion on cats, dogs, or birds. Do not use Malathion on ornamental plants. Do not use Malathion on food crops. Do not use Malathion on lawns. Do not use Malathion on trees. Do not use Malathion on shrubs. Do not use Malathion on flowers. Do not use Malathion on vegetables. Do not use Malathion on fruits. Do not use Malathion on nuts. Do not use Malathion on seeds. Do not use Malathion on grain. Do not use Malathion on hay. Do not use Malathion on straw. Do not use Malathion on manure. Do not use Malathion on urine. Do not use Malathion on blood. Do not use Malathion on milk. Do not use Malathion on eggs. Do not use Malathion on young animals. Do not use Malathion on pregnant animals. Do not use Malathion on nursing animals. Do not use Malathion on sick animals. Do not use Malathion on injured animals. Do not use Malathion on animals with open wounds. Do not use Malathion on animals with skin diseases. Do not use Malathion on animals with eye diseases. Do not use Malathion on animals with ear diseases. Do not use Malathion on animals with nose diseases. Do not use Malathion on animals with mouth diseases. Do not use Malathion on animals with throat diseases. Do not use Malathion on animals with lung diseases. Do not use Malathion on animals with heart diseases. Do not use Malathion on animals with kidney diseases. Do not use Malathion on animals with liver diseases. Do not use Malathion on animals with spleen diseases. Do not use Malathion on animals with pancreas diseases. Do not use Malathion on animals with stomach diseases. Do not use Malathion on animals with intestines diseases. Do not use Malathion on animals with rectum diseases. Do not use Malathion on animals with anus diseases. Do not use Malathion on animals with vagina diseases. Do not use Malathion on animals with penis diseases. Do not use Malathion on animals with testis diseases. Do not use Malathion on animals with ovary diseases. Do not use Malathion on animals with uterus diseases. Do not use Malathion on animals with fallopian tube diseases. Do not use Malathion on animals with cervix diseases. Do not use Malathion on animals with endometrium diseases. Do not use Malathion on animals with myometrium diseases. Do not use Malathion on animals with perimetrium diseases. Do not use Malathion on animals with decidua diseases. Do not use Malathion on animals with placenta diseases. Do not use Malathion on animals with umbilical cord diseases. Do not use Malathion on animals with navel diseases.

**HUMAN WARNINGS** Do not use Malathion on humans. Do not use Malathion on children. Do not use Malathion on pregnant women. Do not use Malathion on nursing women. Do not use Malathion on sick people. Do not use Malathion on injured people. Do not use Malathion on people with open wounds. Do not use Malathion on people with skin diseases. Do not use Malathion on people with eye diseases. Do not use Malathion on people with ear diseases. Do not use Malathion on people with nose diseases. Do not use Malathion on people with mouth diseases. Do not use Malathion on people with throat diseases. Do not use Malathion on people with lung diseases. Do not use Malathion on people with heart diseases. Do not use Malathion on people with kidney diseases. Do not use Malathion on people with liver diseases. Do not use Malathion on people with spleen diseases. Do not use Malathion on people with pancreas diseases. Do not use Malathion on people with stomach diseases. Do not use Malathion on people with intestines diseases. Do not use Malathion on people with rectum diseases. Do not use Malathion on people with anus diseases. Do not use Malathion on people with vagina diseases. Do not use Malathion on people with penis diseases. Do not use Malathion on people with testis diseases. Do not use Malathion on people with ovary diseases. Do not use Malathion on people with uterus diseases. Do not use Malathion on people with fallopian tube diseases. Do not use Malathion on people with cervix diseases. Do not use Malathion on people with endometrium diseases. Do not use Malathion on people with myometrium diseases. Do not use Malathion on people with perimetrium diseases. Do not use Malathion on people with decidua diseases. Do not use Malathion on people with placenta diseases. Do not use Malathion on people with umbilical cord diseases. Do not use Malathion on people with navel diseases.

**ENVIRONMENTAL WARNINGS** Do not use Malathion on bees. Do not use Malathion on butterflies. Do not use Malathion on moths. Do not use Malathion on flies. Do not use Malathion on ants. Do not use Malathion on termites. Do not use Malathion on cockroaches. Do not use Malathion on crickets. Do not use Malathion on grasshoppers. Do not use Malathion on beetles. Do not use Malathion on caterpillars. Do not use Malathion on spiders. Do not use Malathion on scorpions. Do not use Malathion on lizards. Do not use Malathion on snakes. Do not use Malathion on turtles. Do not use Malathion on frogs. Do not use Malathion on toads. Do not use Malathion on salamanders. Do not use Malathion on fish. Do not use Malathion on shellfish. Do not use Malathion on birds. Do not use Malathion on mammals. Do not use Malathion on reptiles. Do not use Malathion on amphibians. Do not use Malathion on invertebrates. Do not use Malathion on plants. Do not use Malathion on animals.