



1% LINDANE DUST

ACTIVE INGREDIENT

Gamma Isomer of Benzene Hexachloride from Lindane* 1%

INERT INGREDIENTS: 99%

TOTAL 100%

*Gamma Isomer of Benzene Hexachloride

CAUTION: KEEP OUT OF REACH OF CHILDREN

DIRECTIONS FOR USE:

Use 1% LINDANE DUST against the following pests on the listed plants. The usual rate of application is 25-30 lbs. per acre. However, when plants are small, as little as 10-15 lbs. per acre may be used and when plants are large, insects numerous, or when applied by airplane, up to 40 lbs. per acre may be required. Consult your State Agriculture Extension Service. Repeat applications as necessary. Note specific limitations under listed crops.

CELERY—Aphids, Thrips, Tarnished Plant Bugs and certain types of Lepidopterous Larvae. Apply before crop is half grown and before bunching.

COLE CROPS (CABBAGE, BROCCOLI)—Aphids, Thrips, Cutworms, Maggots. Apply only before heads or other edible parts begin to form.

CUCUMBERS, CANTALOUPE, WATERMELONS, SQUASH, PUMPKINS, MELONS—Cucumber Beetles, Aphids, Melon Worm, Leaf Miner, Thrips. Apply only before the blossoms open. Application to Melons, Pumpkins and Squash may result in reduction of quality at harvest time.

EGG PLANTS—Aphids, Leaf Miners, Lace Wing Bugs. Do not apply after edible parts start to form.

GLADIOLUS—Thrips.

ONIONS—Thrips. Make no applications to green or spring onions.

PEPPERS—Aphids, Leaf Miners. Apply only prior to blossom fall. Do not apply after edible parts start to form.

SPINACH—Aphids, Leaf Miners, Thrips, Loopers, Web Worms. Apply only to seedling Spinach.

STRAWBERRIES—Spittle Bugs, Strawberry Weevils. Apply only before blossoms open. Thoroughly treat the flower buds. Do not apply after fruit starts to form. Do not apply more than 15 pounds of 1% Lindane Dust per acre for foliage treatment.

TOMATOES—Colorado Potato Beetle, Flea Beetle, Aphids, Thrips. Apply only before blossoms fall. Do not apply after edible parts begin to form.

TOBACCO—Green Peach Aphids. Plant beds should be dusted at the rate of 20 lbs. per acre, repeating once a week.

1% LINDANE DUST is an insecticidal dust containing Gamma Isomer of Benzene Hexachloride. It is suitable for application by means of standard ground dusters or airplane dusting equipment at any time of the day that the weather permits.

CAUTION: Avoid breathing of vapor or dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and food stuffs. Rinse spray equipment and containers and dispose of liquid wastes in a pit in non-crop lands located away from water supplies. May be absorbed through the skin. Harmful if swallowed. Do not reuse container. Destroy when empty.

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift from areas treated. 1% LINDANE DUST should not be applied under conditions which might result in a toxic residue, or residual odor or taste remaining on foodstuffs or edible portions of treated crops or forage at harvest time. Do not use on forage crops to be fed to dairy animals or to animals being finished for slaughtering.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

517-372

E.P.A. Reg. No. 2217-179

CONTENTS 50 LBS.

MANUFACTURED BY

PBI - GORDON CORPORATION

KANSAS CITY, KANSAS