FOR FARM GRAIN STORAGE

For control of the insects as listed on the front panel, use a coarse, wet spray to cover walls, ceilings, floors and surrounding areas of the grain bin. Use approximately one gallon for 1,000 square feet of surface. 'reat empty bins and walls at least 24 hours before filling. Dead insects should be swept up and burned. Do not spray directly onto grain. Residual insect control will last up to three weeks.

OTHER USES

This product may be used in dairy and livestock barns for control of House Flies, Gnats, and Roaches. Apply the material as is, using one gallon for 500 to 1,000 square feet of walls, ceilings, beams, or other surfaces. Treat areas where Flies congregate such as beams, posts, and windows. For Roaches treat all cracks and crevices paying particular attention to baseboards and floor areas. Repeat application as needed for control.

NOTE: Remove animals from buildings during the spray operation. Do not contaminate milking utensils, food, or feed bunks.

CAUTION

Do not use directly on food, grain, feed, or animals. Cover all containers before spraying the area.

Harmful if swallowed. Avoid contact with eyes, skin and clothing. Wash hands and rinse equipment thoroughly after spraying. Do not inhale spray mist. Keep away from Children.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

324-1067

MILL AND FARM BIN SPRAY

Ready to Use

A New Duick hilling Residual Spray for Farm Branaries, Mills and Grain Elevators

WEEVE, RICE WEEVIL, BRAN BUGS, SLOUS
BEETLES, AND ENDIAN MEAL MOTH

CAUTION: Keep Out of Reach of Children

ACTIVE INOPEDIENTS 103

the second second

the state of the s

CONTENTS ONE GALLON

GORDON CORPORATION KANSAS CITY, KANSAS

FOR GRAIN ELEVATOR STORAGE

For control of the insects listed on the front panel use one gallon for approximately 1,000 square feet of bin area, conveyors and dump bins. Apply as a coarse, wet spray to cracks and crevices of infested areas. Allow 24 hours before refilling bin. Do not spray directly onto grain. Plan on this treatment at any time when bins are empty and infestation are present are anticipated.

FOR CEREAL PROCESSING MILLS

For control of insects as listed as well as Roaches and Indian Meal Moth apply to surfaces using one gallon for 1,000 square feet. Treat storage warehouses before refilling. Repeat treatment as needed spraying cracks and crevices and other hiding places for Roaches and insects as listed.

FOR USE IN FOGGING

Apply only in closed empty storage bins, elevators or warehouses. Employ a fog or mist generator to produce a gallon of Mill and Farm Bin Spray per 40,000 cubic feet. This application will control exposed stages of Indian Meal Moth, Angoumois, Grain Moths, Grain Beetles, Granary and Rice Weevils as well as those House Flies, Gnats, and Roaches that may be present at the time of fogging. Keep the area closed over night or at least one to two hours. Sweep the dead insects up and burn. Do not apply in any way that would contaminate food utensils, food, feed, or other food handling equipment.

U.S.D.A. Reg. No. 2217-251 924-1067



Ready to Use

A New Quick Killing Residual Spray for Farm Granaries, Mills and Grain Elevators

KILLS THE EXPOSED STAGES OF GRANARY WEEVIL, RICE WEEVIL, BRAN BUGS, FLOUR BEETLES, AND INDIAN MEAL MOTH

CAUTION: Keep Out of Reach of Children
See Side Panels for Additional Cautions

ACTIVE INGREDIENTS 100%

*Technica!	Piperonyl Buto	oxide	·	.250%
Pyrethrins				.025%
**Technical	Metho cychlor		2	2.500%
Petroleum	Hydrocarbons		97	7.225%
			100	000%

^{**}Consists of 2.2% 2,2-bis (p-methoxyphanyl) -1,1,1-trichloroethane and .3% other isomers and related compounds.

CONTENTS ONE GALLON

^{*}Equivalent to .20% of (butylcarbityl) (6-propylpiperonyl) ether and 0.5% of related compounds.