

MILL AND FARM BIN SPRAY

Ready to Use

A Quick Killing Residual Spray for
Farm Granaries, Mills and Grain Elevators

KILLS THE EXPOSED STAGES OF GRANARY
WEEVIL, RICE WEEVIL, BRAN BUGS, FLOUR
BEETLES, AND INDIAN MEAL MOTH

ACTIVE INGREDIENTS:

*Technical Piperonyl Butoxide	0.250%
Pyrethrins	0.025%
**Technical Methoxychlor	2.500%
INERT INGREDIENTS	97.225%
TOTAL	100.000%

* Equivalent to 0.20% of butylcarbitol 6-propylpiperonyl ether and 0.05% of related compounds.
** Consists of 2.2% 1,1,1-trichloro-2,2-bis(p-methoxyphenyl)ethane and 0.3% other isomers and related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION


See Side Panels for Precautionary Statements and Statement of Practical Treatment

NET CONTENTS ONE GALLON

924/1089 AP

EPA REG. NO. 2217-251
EPA EST. NO. 2217-KS-1

Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, KS 66118



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin and clothing. Wash hands and rinse equipment thoroughly after spraying. Do not inhale spray mist. Keep away from children. Do not use directly on food, grain, feed, or animals. Cover all containers before spraying the area.

Statement of Practical Treatment

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

IF INHALED: Remove victim to fresh air and apply respiration as indicated.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES: Flush with plenty of water and get medical attention immediately.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes.) Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazard

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

FOR FARM GRAIN STORAGE

For control of the insects as listed on the front panel, use a coarse spray to cover walls, ceilings, floors and surrounding areas of the grain bin. Use approximately one gallon for 1,000 square feet of surface. Treat empty bins and walls at least 24 hours before filling. Dead insects should be swept up and burned. Do not spray directly onto grain. Residual insect control will last up to three weeks.

FOR GRAIN ELEVATOR STORAGE

Indoor residual spray treatment: For control of the insects listed on the front panel use one gallon for approximately 1,000 square feet of bin area, conveyors and dump bins. Apply with adequate force to drive the spray into cracks and crevices of bins. Spray when bins are empty or before products are stored. Clean area thoroughly and remove old grain and debris before spraying. On elevator tunnels, gallery floors and head houses, make 3 applications. Allow at least 24 hours for sprayed surfaces to dry before introducing grain into bins.

FOR CEREAL PROCESSING MILLS

Specific sites include: cereal processing mills, cereal storage areas, and flour mills.

Indoor residual spray treatment: For control of insects as listed as well as Roaches and Indian Meal Moth apply to surfaces using one gallon for 1,000 square feet. Treat storage warehouses before refilling. Apply in small amounts into cracks and crevices between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs, and bases. Care should be taken to avoid introducing the material into the air. Repeat treatment as needed spraying cracks and crevices and other hiding places for Roaches and insects as listed.

INDOOR FOGGING

Apply only in closed empty storage bins, elevators or warehouses. Apply only to those areas where grain and other food products have been removed. Employ a fog or mist generator to produce a gallon of Mill and Farm Bin Spray per 40,000 cubic feet. This application will control exposed stages of Indian Meal Moth, Angoumois, Grain Moths, Grain Beetles, Granary and Rice Weevils as well as those of House Flies, Gnats and Roaches that may be present at the time of fogging. Keep the area closed overnight or at least one to two hours. Sweep the dead insects up and burn. Do not apply in any way that would contaminate food utensils, food, feed, or other food handling equipment.

LIVESTOCK BUILDINGS AND PREMISES

Specific sites include: animal shelters, barns (including dairy barns), milk rooms, pens, sheds, and stalls

Indoor premise treatment for the control of House Flies, Gnats and Roaches. Apply the material as is, using 1 gallon of product for 500-1000 square feet. Apply to wall and floor surfaces inside the building, including cracks, crevices, base boards, and animal quarters such as stalls, pens and bedding. Treat areas where flies congregate such as beams, posts, and windows. For roaches treat all cracks and crevices paying particular attention to baseboards and floor areas. Repeat application as needed for control. Cover securely all milk-handling equipment, feedstuffs, troughs, bins, or other food and water receptacles while spraying. Dairy animals should not be present while spraying.

LIMITED WARRANTY. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

EXCLUSION OF OTHER WARRANTIES AND REMEDIES. Except where such disclaimers and exclusions are specifically prohibited by applicable law. THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty. BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.