

DIRECTIONS FOR USE

SURFACE TREATMENTS FOR INDIAN MEAL MOTH

As a Spray: To protect stored grain from attack by Indian meal moth, apply Grain Gard Emulsifiable to the surface of clean or uninfested grain at the rate of $\frac{1}{2}$ pint in one to two gallons of water for each 1,000 square feet of grain surface area. Apply the spray evenly over the surface of the grain. Apply immediately after grain is loaded into storage and repeat if necessary.

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist, avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before reuse. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

Malathion may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

923-1174

DIRECTIONS FOR USE

APPLIED TO GRAIN AT TIME OF STORAGE

For the protection of stored grain such as wheat, oats, rice, corn, rye, barley, and grain sorghum, apply one pint of Grain Gard Emulsifiable in 2 to 5 gallons of water to each 1,000 bushels of grain.

Apply the spray as the grain is being loaded or turned into final storage. Any of the standard applicators on the market which can be calibrated to deliver a known volume of liquid are suitable for applying malathion.

For small amounts of grain in farm storage, where special application equipment is not available, any type of low-pressure sprayer (holding a gallon or more) can be used. The spray can be applied to the grain stream as the grain is being elevated into storage. The first step would be to test-spray a tank of water to determine the rate at which the sprayer is discharging—then regulate the flow of grain to get on the proper amount of malathion.

AS A CLEAN-UP SPRAY

Before new grain goes in, a residual malathion spray should be applied to walls, floors, and machinery in grain elevators or farm storage. This applies also to truck beds, boxcars, and ships' holds. Mix this spray at the rate of one gallon of Grain Gard Emulsifiable in 25 gallons of water. Spray all surfaces to the run-off point—making particularly sure that the spray is forced into cracks and crevices. This same spray should be used around the outside of bins and elevators to help prevent re-infestation.

USE

INDIAN MEAL MOTH

from attack by Indian
sifiable to the surface
rate of 1/2 pint in one
00 square feet of grain
ly over the surface of
r grain is loaded into

inhaled, or absorbed
spray mist, avoid pet-
Wash thoroughly with
immediately and wash
of feed or foodstuffs
estic animals

e inhibition. If person
Atropine is antidota-
e administered in con

TORE NEAR HEAT OF

out of lakes, streams,
water by cleaning up
apply the product


her empty

product conforms to the
l. Since conditions
with other chemical
ment, will vary. Seller
e stated on this label



GRAIN GARD EMULSIFIABLE

PREMIUM 57% MALATHION



APP

for the pro
the corn,
f Grain G
each 1,000

Apply the
into final
the market
volume of

For small a
application
pressure sp
The spray o
is being ele
test-spray a
the sprayer
to get on t

Before new
should be a
elevators or
boxcars, an
one gallon
water. Spra
particularly
previews. Th
side of the

GRA