

DIRECTIONS FOR USE

ANIMAL BUILDINGS - Apply Vapona Insecticide 25-E as a 0.5% spray to control flies, mosquitoes, ticks, ear mites, fleas, ants, sowbugs, and brown bugs.

To prepare a 0.5% spray dilute one pint of Vapona Insecticide 25-E with 39 pints of water. Apply as a coarse wet spray at the rate of 1 quart per 1,000 square feet.

Before spraying, close doors to reduce air movement as much as possible. Close windows, awnings, and other openings. Do not treat areas where animals drink. Do not contaminate water, feed or foodstuffs, milk or milk products. Repeat as necessary.

Use Vapona Insecticide 25-E in livestock buildings, loading sheds, pig pens, and cattle kraals. Spray around doorways, feed and water troughs, and animal resting areas. Do not contaminate milk or milk products.

Use Vapona Insecticide 25-E in poultry houses, brooder houses, and chicken coops. Spray around doorways, feed and water troughs, and animal resting areas. Do not contaminate feed or feedstuffs.

Use Vapona Insecticide 25-E in horse stalls, feed racks, and horse trailers. Spray around doorways, feed racks, and animal resting areas. Do not contaminate feed or feedstuffs.

Use Vapona Insecticide 25-E in feed racks, windows, stalls, and horse trailers. Spray around doorways, feed racks, and animal resting areas. Do not contaminate feed or feedstuffs.

EGGING ANIMAL BUILDINGS - Apply Vapona Insecticide 25-E as a 0.5% spray to control flies, mosquitoes, ticks, ear mites, fleas, ants, sowbugs, and brown bugs.

To prepare a 0.5% spray dilute one pint of Vapona Insecticide 25-E with 39 pints of water. Apply as a coarse wet spray at the rate of 1 quart per 1,000 square feet.

Before spraying, close doors to reduce air movement as much as possible. Close windows, awnings, and other openings. Do not treat areas where animals drink. Do not contaminate water, feed or foodstuffs, milk or milk products. Repeat as necessary.



VAPONA[®] INSECTICIDE 25-E

(Contains 2.0 lbs. VAPONA Insecticide per gallon)

ACTIVE INGREDIENTS:	
2,2-Dichlorovinyl dimethyl phosphorotriester*	22.9%
Related Compounds	1.7%
Xylene	68.9%
INERT INGREDIENTS	
	5.5%
	100.00%

© A Shell Company
*Equivalent to DDT insecticide

NET CONTENTS ONE GALLON

Keep Out of Reach of Children.

See side panels for Antidote and additional precautions.

CATTLE FEED LOTS, STOCKYARDS, HOLDING PENS - Apply Vapona Insecticide 25-E as a 0.5% spray to control flies, mosquitoes, ticks, ear mites, fleas, ants, sowbugs, and brown bugs.

To prepare a 0.5% spray dilute one pint of Vapona Insecticide 25-E with 39 pints of water. Apply as a coarse wet spray at the rate of 1 quart per 1,000 square feet. Par be given to areas where flies congregate such as around feed troughs and walls of buildings. Animal treatment. Avoid direct application to exposed feed.

STOP! READ CAREFULLY

DANGER: POISONING - Vapona Insecticide 25-E and its contents. Insecticide preparations must be stored in cool and dark places.

Use the concentrate in a safe place. Children should be kept away from the concentrate. Do not apply immediately to water. Do not apply to feed.

Use the concentrate in water with care. Do not apply to the outside of the bottle of concentrate.

Use the concentrate in water with care. Do not apply to the outside of the bottle of concentrate.

Use the concentrate in water with care. Do not apply to the outside of the bottle of concentrate.

DANGER: POISONING - Vapona Insecticide 25-E and its contents. Insecticide preparations must be stored in cool and dark places.

Use the concentrate in a safe place. Children should be kept away from the concentrate. Do not apply immediately to water. Do not apply to feed.

Use the concentrate in water with care. Do not apply to the outside of the bottle of concentrate.

Use the concentrate in water with care. Do not apply to the outside of the bottle of concentrate.

DIRECTIONS FOR USE

ANIMAL BUILDINGS—Apply Vapona Insecticide 25-E as a 0.5% spray to control flies, gnats, mosquitoes, roaches, fleas, ants, sowbugs, and brown dog ticks.

To prepare a 0.5% spray, dilute one pint of Vapona Insecticide 25-E with 6 gallons of water. Apply as a coarse, wet spray at the rate of quart per 1000 square feet.

Before applying in enclosed areas, reduce air movement as much as possible by closing doors, windows, and other openings. Do not treat animals or humans. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Repeat as necessary.

Dairy Barns, Stables, Livestock Buildings, Loafing Sheds, Pig Pens, and Outdoor Areas: Spray around doorways, feed storage rooms, alleyways, and window sills. Do not contaminate milk or milking utensils.

Poultry Houses: Apply to manure, window sills, outside of penned enclosures, and on floor of feed storage rooms.

Manure Piles and Garbage Dumps: For the control of fly maggots in maggot breeding areas apply at the rate of one to two quarts of 0.5% spray per 100 square feet.

Dog Kennels: Apply to outside dog runways, window sills, and ledges. For the control of fleas and brown dog ticks increase dosage to two quarts of 0.5% spray per 1000 square feet.

FOGGING ANIMAL BUILDINGS—Apply Vapona Insecticide 25-E as a 0.5%-1.0% to control flies, gnats, and mosquitoes.

To prepare a 0.5% fogging solution, dilute one pint of Vapona Insecticide 25-E with 7 gallons diesel oil (for a 1.0% solution, dilute one quart per 7 gallons oil).

Apply at the rate of one quart of a 0.5% solution or one pint of a 1% solution per 8,000 cubic feet. Before applying, reduce air movement as much as possible by closing doors, windows, and other openings. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Do not use this solution for fogging purposes in areas where animals have received direct application within 8 hours.


834-1273



VAPONA[®] INSECTICIDE 25-E

DANGER: Keep Out of Reach of Children.

POISON



POISON

MANUFACTURED BY

PBI - GORDON CORPORATION

KANSAS CITY, KANSAS 66118

CATTLE FEED LOTS, STOCKYARDS, HOLDING PENS: Apply Vapona Insecticide 25-E as a 0.5% spray to control flies, gnats, and mosquitoes.

To prepare a 0.5% spray dilute one pint of Vapona Insecticide 25-E with 6 gallons of water. Apply with a mist blower or other equipment for an over-all application at the rate of 0.2 pound per acre (approximately 5 gallons of 0.5% spray). Do not apply to areas where flies congregate such as around feed bunks, and walls of buildings. Animal treatment. Avoid direct application to exposed feed. For baited solutions dissolve one pound of sugar per quart of water.

STOP! READ CAREFULLY

DANGER IN USING — Vapona Insecticide 25-E and its concentrate are highly poisonous. Therefore, precautions must be observed (see first aid, antidote and warnings below).

Store the concentrate in a safe place. Children should not be allowed to reach or handle the concentrate or dilute solution. Only the amount of diluted emulsion recommended that only the amount of diluted emulsion should immediately be prepared from the concentrate.

Mix the concentrate into water outdoors or in a well-ventilated area. Wipe off the outside of the bottle of concentrate in running water.

This product is highly toxic to fish and wildlife. Do not apply to any body of water nor apply to any area not specified on the label. Fish and wildlife on treated areas may be killed. Do not apply to areas where drift from areas treated. Do not apply where it may contaminate water. Do not contaminate water by cleaning of equipment. Do not contaminate feed, foodstuffs, water, milk or other animal products. **NOT FOR USE OR STORAGE IN OR AROUND THE HOME.**

DANGER: Poisonous if swallowed, inhaled, or absorbed. Do not get on skin, in eyes, and do not breathe vapors or fumes. Wear clean rubber gloves, waterproof clothing, goggles when handling the concentrate to prepare dilute solution or respirator of a type passed by the U.S. Bureau of Health (for insecticide) protection. Keep all unprotected persons away. In case of contact, wash immediately with soap and water. Wash with soap and water before eating or smoking. Wash clothing with soap and hot water before reuse. **DO NOT USE, POUR, SPILL, OR STORE NEAR HEADS.**

CATTLE FEED LOTS, STOCKYARDS, HOLDING PENS, AND CORRALS — Apply Vapona Insecticide 25-E as a 0.5% spray to control flies, gnats, and mosquitoes.

To prepare a 0.5% spray dilute one pint of Vapona Insecticide 25-E with 6 gallons of water. Apply with a mist blower or other suitable equipment as an over-all application at the rate of 0.2 pound of Technical Vapona per acre (approximately 5 gallons of 0.5% spray). Particular attention should be given to areas where flies congregate such as fences, spillage areas, around feed bunks, and walls of buildings. Animals may be present during treatment. Avoid direct application to exposed feed and water.

For baited solutions dissolve one pound of sugar per 5-10 gallons of water.

STOP! READ CAREFULLY

DANGER IN USING — Vapona Insecticide 25-E and the diluted emulsion are poisonous. Therefore, precautions must be observed in handling them. (See antidote and warnings below).

Store the concentrate in a safe place. Children or animals must not be allowed to reach or handle the concentrate or diluted emulsion. It is recommended that only the amount of diluted emulsion that will be used immediately be prepared from the concentrate.

Mix the concentrate into water outdoors or in a well ventilated room. Wash off the outside of the bottle of concentrate in running water before storing away.

This product is highly toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on this label. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not contaminate feed, foodstuffs, water, milk or milking utensils.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

DANGER: Poisonous if swallowed, inhaled, or absorbed through the skin or eyes. Do not get on skin, in eyes, and do not breathe fumes. Avoid exposure to fumes. Wear clean rubber gloves, waterproof protective clothing, and goggles when handling the concentrate to prepare emulsions. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for DDVP (Vapona insecticide) protection. Keep all unprotected persons out of operational area.

In case of contact, wash immediately with soap and water. Wash thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

ANTIDOTE AND FIRST AID TREATMENT: Atropine is the emergency antidote for Vapona Insecticide poisoning. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL—If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (one tablespoon of salt to a glass of water.) NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL—If the material has been spilled on the skin, immediately remove the patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least 10 minutes.

IF WARNING SYMPTOMS APPEAR: (See WARNING SYMPTOMS below). Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals in doses of one to two milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage.

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

2-PAM is also antidotal and may be used in conjunction with atropine.

CONTAINER DISPOSAL: Do not reuse. Destroy when empty. Glass Jugs—After rinsing, break and bury with wastes. Metal Cans—Destroy by perforating or crushing, and bury with wastes.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

E.P.A. Reg. No 2217-291-AA