

**BEEF CATTLE:** For control of Horn Flies and Ticks; mix one quart of Hi-Tox to 38 gallons of water. This treatment will give two to four week control. Treat when Horn Flies appear in spring or, if Ticks are present. Treat animals to thoroughly wet the hide. For Lice use the same rate and single treatment is usually sufficient.

**CALVES:** If calves must be treated for Lice, dilute 1 qt. to 75 gallons water (2 oz. per 4½ gallons). Do not treat those under 3 months of age. Do not apply more than 1 pint of diluted spray per animal.

**Backrubber Use:** For control of Hornflies and to aid in control of Face Flies and Lice. Dilute one quart with four gallons of fuel oil. Apply the diluted material to saturate the Backrubber mechanism. Recharge the Backrubber every two weeks or as necessary for control. Locate Backrubber to give Cattle free access to it. Do not allow calves under 12 weeks of age, or Dairy Animals, or Animals within four weeks of slaughter to treat themselves.

**Face Flies:** For use against Face Flies, one end of the Backrubber must be low enough to allow Cattle to rub their heads. Not more than 15 to 20 inches above the ground. The product is only effective for these cattle that rub the face and head. Dilute as for Backrubber use.

**HOGS:** For Lice control; use one quart to 75 gallons of water. Use high pressure spray to thoroughly wet animals. Treat when Lice are present, or about November 1 and again, if necessary for complete control in about 15 days. Treat sows about 30 to 45 days before farrowing.

**HORSES:** For control of Horn Flies and Lice as needed; mix one ounce per gallon of water. Apply to animals as a sponge application, treating areas where insects are present, or are troublesome.

**Sarcoptic Mange Mites:** Accurately mix one quart to 38 gallons of water. Thoroughly wet animals. Repeat applications in 15 days. 835-867



## HI-TOX LIVESTOCK SPRAY EMULSIFIABLE CONCENTRATE

Contains 6 pounds Technical Toxaphene per gallon

**WARNING: Keep Out of Reach of Children**

See Side Panels for Additional Warnings

**ACTIVE INGREDIENTS:**

Toxaphene (Technical Chlorinated Camphene containing 67% to 69% Chlorine).....	59.0%
Petroleum Hydrocarbons .....	27.9%
Pine Oil .....	0.7%

<b>INERT INGREDIENTS:</b> .....	12.4%
---------------------------------	-------

Total.....	100.0%
------------	--------

**FOR USE ON BEEF CATTLE — HOGS**

**SHEEP — HORSES — GOATS**

**CONTROLS**

**FACEFLIES — HORNFLIES — SARCOPTIC MANGE —**

**LICE AND TICKS — PSOROPTIC SHEEP SCAB**

**NET CONTENTS 1 GALLON**

835-867



MANUFACTURED BY

**GORDON CHEMICAL CO., INC.**  
KANSAS CITY, KANSAS



**HI-TOX  
LIVESTOCK SPRAY  
EMULSIFIABLE CONCENTRATE**

Contains 6 pounds Technical Toxaphene per gallon  
**WARNING: Keep Out of Reach of Children**  
See Side Panels for Additional Warnings

**ACTIVE INGREDIENTS:**

Toxaphene (Technical Chlorinated Camphene containing 67% to 69% Chlorine).....	59.0%
Petroleum Hydrocarbons .....	27.9%
Pine Oil .....	0.7%
<b>INERT INGREDIENTS:</b> .....	12.4%
Total.....	100.0%

**FOR USE ON BEEF CATTLE — HOGS  
SHEEP — HORSES — GOATS**

**CONTROLS  
FACEFLIES — HORNFLIES — SARCOPTIC MANGE —  
LICE AND TICKS — PSOROPTIC SHEEP SCAB  
NET CONTENTS 1 GALLON**

835-867

MANUFACTURED BY  
**GORDON CHEMICAL CO., INC.**  
KANSAS CITY, KANSAS

**SHEEP AND GOATS:** For Lice and Sheep Tick control; mix one quart in 300 gallons of water (.06% as applied). If necessary, repeat this treatment 15 days later.

**Fleeca Worms:** Use one quart to 38 gallons of water and thoroughly spray to wet animals.

**Psoroptic Sheep Scab:** For control and cure, dip animals in a vat containing a solution of one quart Hi-Tox per every 152 gallons of water. Make sure each animal remains in dipping solution at least one-half minute. Duck heads for thorough treatment.

**WARNING:** Do not take internally. May be fatal if swallowed. Avoid contamination of feed and foodstuffs. Do not breathe mist. Do not get into eyes, on skin or on clothing. In case of spillage on skin wash with soap and water.

Do not use on household pets, dairy animals, or animals within 28 days of slaughter, or humans. Do not use in Dairy Barns, Milk Houses or Poultry Houses. Do not mix or store near fire or flame.

To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material or washings from spray tank.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

**ANTIDOTE:** Give Tablespoon of table salt in glass of warm water and repeat until vomit is clear. Call Physician.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

835-867

USDA Reg. No. 2217-301

3-27-68

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTRATION  
ED UNDER NO. 2217-301

*Our Name Is Your*

*Guarantee*

*of*

*Quality*