

tie with water to 12 or 15 gallon level.
Apply to the pre-measured area of 1,000 square feet. Spray until bottle is empty. Repeat for each 1,000 square feet of lawn.

Pressure Sprayer: Mix 4 ounces with each 2 gallons of water and spray on 1,000 square teet. Sprinkle area to get chemical to soil level.

When to Use: Make two applications, the first near April 15th, and second June 1. Spray in early evening, as worms feed at this time. In late summer, if lawn has been attacked and taribegins to die, rake the thatch from in

Superkill sod webworm spray

appear around the first of June and then again in late August. Spray ground around trees to kill larvae.

WARNING: May be fatal if swallowed! Keep out of reach of children. Do not breathe spray mist. Do not get on skin, clothing or in eyes. Wash thoroughly after using. Do not contaminate food or cooking utensils. May be absorbed through the skin. Do not use on household pets or humans. In case of contact, wash immediately with soap and water.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

This product will kill fish and wildlife. Keep out of takes, streams, and ponds. Birds feeding on treated areas may



DIRECTIONS FOR USE

Sod Webworm Spray is offered especially for the cantrol of the Lawn Sod Webworms found in Biue Grass Turf. Read directions carefully before using. Also controls certain other pests found on ornamental plants.

PESTS

LAWN AND TURF RATE OF DILUTION AND COMMENTS HOW TO USE:

SOD WEBWORM (Blue Grass Lawns) mix as follows:

for each 1.000 square feet of lawn.

Hose-end Lawn Sprayers: Add 4 ounces concentrate to sprayer bottle. Fill bot-The matter well to the action of the graph of the control of the c

surface of the lawn and use 6 ounces per 1,000 square feet. Thoroughly water lawn 3 to 4 days after treatment Plan on using a good lawn fertilizer to rejuvenate turf.

CHIGGERS and

Mix 4 tbsp. to 4½ pints water (2 qt. in 10 gallons). Spray to thoroughly cover all grass and weed foliage where insects hide. If possible, make application 6 to 12 hours before rain. Repeat as needed.

BOXELDER BUGS

696-372

Mix 2 qts. to 5 gallons water (4 tbsp. per 142 pint). Spray or paint on foundations, porches, steps, window and door frames and places where insects may enter house. Treat boxelder trees and also spray insects directly. Repeat as needed for control.

Ø



superkill

sod webworm spray

CONTROLS: Chiggers, Ticks, Boxelder Bugs

ACTIVE INGREDIENTS: Toxaphene (Technical chlorinated camphene, (From Lindane) Petroleum Hydrocarbons INERT INGREDIENTS 8.00% 100.00%

WARNING: Keep Out of Reach of Children See Side Panels for Additional Warnings 696-372

CONTENTS 1 QUART

DIRECTIONS FOR USE

APPLY WHEN INSECTS APPEAR OR PLANT DAMAGE IS FIRST SEEN

CANKERWORMS ON ELMS

ORNAMENTALS

Mix 2 qts. in 50 gailons water (4 tbsp. in 11/2 gallons). Apply in spring when insects are first seen. Cover all new growth and leaves. Use high pressure

CATALPA WORMS on Catalpa

Mix 2 qts. in 50 gallons of water (4 tbsp. in 11/2 gallons) and thoroughly spray foliage and insects. Worms may appear around the first then also not

and the second be killed. Inis product is not to be used on animals of any kind.

Do not apply whore runoff is likely to occur. Do not contaminate water by cleaning of equipment.

Do not use on Redbud trees.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Apply this product only as specified on this label. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

*T.M. Registered, P.B.I.

E.P.A. Reg. No. 2217-324

696-372