

STUPA OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals
WARNING: May be fatal if swallowed, inhaled, or absorbed through skin. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating or smoking. If swallowed, induce vomiting immediately (See Statement of Practical Treatment). Do not allow spray mist to contaminate milk or milking utensils. Cover feed and feed troughs or remove before spraying.

Wear clean, natural rubber gloves, protective clothing, and goggles, face shield or equivalent. Wear a pesticide respirator jointly approved by the Mine Safety and Health Administration and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11.

This product is an organic phosphate which is a cholinesterase inhibitor, and if poisoning should occur, CALL A PHYSICIAN IMMEDIATELY.

Statement of Practical Treatment
If Swallowed: Call a physician or Poison Control immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching back of throat with a finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If Inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If On Skin: Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes: Flush eyes with plenty of water. Call a physician immediately.

Note to Physician: Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally. If symptoms of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

ENVIRONMENTAL HAZARDS:
This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS:
DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

STORAGE & DISPOSAL


STORAGE: Store in original container in a locked storage area inaccessible to children or pets. To prevent cross-contamination, keep separate from other pesticides, fertilizers, seeds, food and feed.

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.



VAPONA® INSECTICIDE DAIRY CATTLE SPRAY



Ready-To-Use for FACE FLIES

Stable Fly — House Fly — Horn Fly

ACTIVE INGREDIENTS:	
2, 2-Dichlorovinyl dimethyl phosphate*	0.93%
Related Compounds*	0.07%
Petroleum distillate	98.93%
INERT INGREDIENTS	0.07%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN

WARNING

See side panels for additional precautionary statements and Statement of Practical Treatment.

NET CONTENTS ONE U.S. GALLON (3.785 Liters)

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

HOUSE FLIES in Dog Kennels, Hog Pens, Stables, Compost areas, manure, Poultry Houses, and Barns. Application of this product should be as a coarse wet spray in these areas. Apply to thoroughly cover walls, flooring, etc. in kennels, barns, and stables. Use no more than 1 pint per 1,000 square feet. This is not a residual spray, therefore treat weekly or as needed for control of flies and fleas. Do not contaminate food or food utensils.

Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

DAIRY AND BEEF CATTLE FACE FLY — STABLE FLY — HORN FLY — HOUSE FLY — MOSQUITOES:

Use a hand mist sprayer and treat entire animal to wet hair with 1 ounce in morning and evening. For Horn Flies, especially treat the back line, withers, and belly. For Stable Flies, make sure to spray leg areas and particularly ankles. Repeat daily as needed for control. Do not apply more than 2 ounces per animal daily. For walk-thru mist sprayers, set to treat animals at rate not to exceed 2 ounces daily.

IMPORTANT: Avoid wetting the hides of animals. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

OVERHEAD MIST DISPENSERS — For control of flies, gnats, and mosquitoes in milkhouses and barns, use 1 pint for each 8,000 cubic feet. Do not contaminate feed or milk utensils. If a program of spraying animals directly is employed daily, do not fog within 8 hours of direct applications.

WARRANTY: The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

965985
EPA REG. NO. 2217-332
EPA EST. NO. 2217-KS-1

