

ACCEPTED
MAY 15 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 271-30 SUBJECT
TO ATTACHED COMMENTS.

VAPONA*
FOGGING SPRAY
CONTAINS DDVP

~~Ready-To-Use for~~

FLIES-MOSQUITOES-GNATS

To be applied by or under the supervision of trained personnel, or by Pest Control Operators only.

ACTIVE INGREDIENTS 100%

2,2-Dichlorovinyl dimethyl phosphate*93%
Related Compounds*07%
Insecticide Base Oil (Petroleum Derivative)	<u>99.00%</u>
TOTAL.....	100.00%

*Equivalent to 1% Vapona Insecticide
*VAPONA_® Insecticide is a registered trademark

CAUTION: Keep Out of Reach of Children
See Side Panel for Additional Cautions

CONTENTS ONE GALLON

Manufactured by
Private Brands, Inc. for Various
Customers as Subregistered

DIRECTIONS FOR USE

FOR FOGGING MACHINES:

MOSQUITOES-FLIES-GNATS-OUTDOORS: Use a fogging machine and apply at rate of 5 to 10 pints of spray per acre in such areas as golf courses, parks, open-air drive-in theaters, disposal areas. Repeat as necessary. Do not treat areas where food or feed crops are growing. **INDOORS:** Fog at rate of one pint of fogging spray per 8,000 cubic feet for control of flies, mosquitoes and gnats in warehouses, industrial plants, theaters, and horse stables. Use this product as a spacespray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. Occupants should vacate before indoor treatments and should not reoccupy the premises until it has been ventilated.

CAUTION: Extinguish all pilot lights before fogging.

FOR MIST TYPE SPRAYERS:


FACE FLIES-HORN FLIES-MOSQUITOES-HOUSE FLIES ON BEEF AND DAIRY CATTLE: Use a mist type hand sprayer or automatic unit that will insure a fine spray mist for desired coverage of the animal. For Horn Flies, direct the spray mist to the back, withers, and belly of the cattle. Make 2 applications of one ounce each day. Treat in the Morning and Evening.

STABLE FLIES--Apply at same rate and timing as for Horn Flies; however, direct the spray to the leg areas making sure to directly spray the ankles

NOTE--Do not exceed the maximum dose of 2 ounces per animal daily. It is highly toxic to flies but loses its strength within the day as most dairy cattle sprays. For this reason, no insecticidal residue is built up. A well planned Fly Control Program, that includes the two spray applications per day as directed, will give results.

HOUSE FLIES: For porches, patios, and Drive-In's, spray refuse area and trash. Treat walks, steps and adjacent walls with a light spray mist. Repeat as needed. Do not contaminate food or food utensils. Do not use more than 1 pint per 1,000 sq. feet.

WARNING--To be applied by or under the supervision of trained personnel or by Pest Control operators only. Use a mask or respirator passed by the U. S. Department of Agriculture for Vapona protection during application. Do not allow the spray mist to contaminate milk or milking utensils. Cover feed and feed troughs or remove before spraying. May be fatal if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling and before eating or smoking. **KEEP OUT OF REACH OF CHILDREN.** This product is an organic phosphate which is a cholinesterase inhibitor and if poisoning should occur, CALL A PHYSICIAN IMMEDIATELY. **ATROPINE IS ANTIDOTAL.**



Keep out of lakes, streams and ponds. This product will kill fish.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be destroyed beyond reuse and buried with wastes.

NON-WARRANTY--The seller makes no warranty of any kind, expressed or implied, concerning the use of this product. The buyer assumes all risk of use or handling, whether in accordance with directions or not.

U.S.D.A. Reg. No. 2217-340

965-

