

FOR USE ON	TO CONTROL	IN 100 GAL.	IN 1 GAL.	COMMENTS
Apples, Pear	European Red Mite, 1-2 pts. Clover Mite Two Spotted Mite Woolly Apple Aphid 1 pt. Forsythia Mite Green Apple Aphid 1 1/2 pts. Woolly Apple Aphid	2 tsp. 1 pt. 1 1/2 tsp.	2 tsp. 1 tsp. 1 1/2 tsp.	Make two or more applications as needed for mites, one or more for aphids. At pre-pink or pink stage of bloom or when aphids appear. Do not use within 3 days of harvest.
Tomatoes	Spider Mites, Aphid	1 1/2 pts. 1 pt.	1 1/2 tsp. 1 tsp.	Make one or more applications as needed. Do not use within 3 days of harvest.
Beans	Mexican Bean Beetle, Spider Mites, Leafhopper	2 pts.	2 tsp.	
Peas	Pea Aphid	2 pts.	2 tsp.	Make one or more applications. Do not use within 3 days of harvest. Do not use pea vines for forage.
Potatoes	Aphids	2 pts.	2 tsp.	
Cabbage, Brussels, Kale, Mustard, Turnip	Aphids, Imported Cabbage Worm	2 pts.	2 tsp.	Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on Broccoli and Turnips, 7 days on others.
Alfalfa for Seed	Aphids	2 pts. per acre in at least 10 gal. water.		Do not apply within 7 days of harvest, or when in bloom.
Flowers, Ornamentals, Shrubs	Oyster Shell Scale Eucynus Scale Aphids Spider Mites White Fly, Thrips	1 pt. 1 1/2 pts.	1 tsp. 1 1/2 tsp.	Apply in summer when infestation occurs. Repeat in 10 to 12 days. For scales, apply when crawlers have



50% MALATHION

Emulsifiable Concentrate

FOR CONTROL OF ANIMAL AND PLANT PESTS

ACTIVE INGREDIENTS:

Malathion* 50%
Xylene 45%

INERT INGREDIENTS 5%

*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings

893-667

NET CONTENTS 1 QUART

MANUFACTURED BY

GORDON CORPORATION

KANSAS CITY, KANSAS

ACCEPTED

May 20, 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT

ACCEPTED WITH COMMENTS

DIRECTIONS

LIQUID FLY BAIT: To barns, feed lots, poultry around food disposals, in syrup and add to 2 gal. desired surfaces.

BEEF CATTLE: For control of lice. Apply 1 to 2 gal. in water. Apply applications at two week intervals. Apply 1 to 2 gal. a complete coverage spray intervals if needed.

SWINE: For control of lice. Apply 1 to 2 gal. in water. Apply a complete coverage spray. One treatment may be needed.

BACK RUBBING DEVICE: Cattle, apply a 2% mixt. oil or kerosene) to the back. There will be a reduction in Horn Fly one to 35 or 45 head of 3 weeks. Do not use on

CAUTION: Do not apply of freshening. Do not use when applying sprays, containers and watering processes or stored. Do not use on children.

WARNING: Harmful if through the skin. Avoid getting in eyes, clothing. Wash thoroughly with soap and water before re-use.

DIRECTIONS FOR USE

LIQUID FLY BAIT: To control House and Stable Flies, Barns, Feed Lots, Poultry houses, Utility buildings and around food disposals, mix 5 ounces with 5 ounces of corn syrup and add to 2 gallons of water. Paint or spray on desired surfaces.

BEEF CATTLE: For control of Lice, add 1 gallon to 100 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed. For control of Ticks, apply 1 to 2 gallons per 100 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed.

SWINE: For control of Lice, apply 1 gallon per 100 gallons of water. Apply a complete spray to animals, pens and litter. One treatment may be sufficient. Repeat application only if needed.

BACK RUBBING DEVICES: For the reduction of Lice on Beef Cattle, apply a 2% mixture (1 quart in 7½ gallons of fuel oil or kerosene) to the back rubbing device. There may also be a reduction in Horn Flies. Make rubbing device accessible one to 35 or 45 head of cattle. Retreat the oiler every 2 or 3 weeks. Do not use on dairy animals.

CAUTION: Do not apply during lactation or within two weeks of freshening. Do not treat calves under one month of age. When applying sprays, avoid contamination of feed, food containers and watering troughs. Do not use where milk is processed or stored. Do not use in dwelling or in reach of children.


WARNING: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist or dust. Avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before re-use. Destroy containers completely when

Our Name Is Your

Guarantee

of

Quality



50% MALATHION

Emulsifiable Concentrate

FOR CONTROL OF ANIMAL AND PLANT PESTS

ACTIVE INGREDIENTS:

Malathion* 50%
Xylene 45%
INERT INGREDIENTS 5%
*0,0-dimethyl dithiophosphate of diethyl
mercaptosuccinate.

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings

893-667

NET CONTENTS 1 QUART

GORDON CORPORATION

FOR USE ON	TO CONTROL	IN 100 GAL.	IN 1 GAL.	COMMENTS
Alfalfa for Seed	Aphids	2 pts. per acre in at least 10 gal. water.	1 tsp.	Do not apply within 7 days of harvest, or when in bloom.
Flowers, Ornamentals, Shrubs	Oyster Shell Scale Cherry Scale	1 pt.	1 tsp.	Apply in summer when infestation oc- curs. Repeat in 10 to 12 days.
	Aphids, Spider mites, White Fly, Thrips	1 1/2 pts.	1 1/2 tsp.	For scales, apply when mature.

ACCEPTED

May 20, 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.

ALATHION

le Concentrate

CONTROL OF D PLANT PESTS

DIENTS:

..... 50%
 45%
 DIENTS 5%
 dithiophosphate of diethyl
 ate.

Out of Reach of Children
 e Panels for Additional Warnings

TENTS 1 QUART

CORPORATION

DIRECTIONS FOR USE

HOUSE AND STABLE FLIES: To control House and Stable Flies
 mix 1 gallon of concentrate with 100 gallons of water. Paint or spray on
 the inside of walls, ceiling and floor.

HOUSE AND STABLE FLIES: To control House and Stable Flies
 mix 1 gallon of concentrate with 100 gallons of water. Paint or spray on
 the inside of walls, ceiling and floor.

HOUSE AND STABLE FLIES: To control House and Stable Flies
 mix 1 gallon of concentrate with 100 gallons of water. Paint or spray on
 the inside of walls, ceiling and floor.

HOUSE AND STABLE FLIES: To control House and Stable Flies
 mix 1 gallon of concentrate with 100 gallons of water. Paint or spray on
 the inside of walls, ceiling and floor.

CAUTION: Do not apply during lactation or within two weeks
 of freshening. Do not treat calves under one month of age.
 When applying sprays, avoid contamination of feed, food
 containers and watering troughs. Do not use where milk is
 processed or stored. Do not use in dwelling or in reach of
 children.

WARNING: Harmful if swallowed, inhaled or absorbed
 through the skin. Avoid breathing spray mist or dust.
 Avoid getting in eyes, on skin, or on clothing. Wash thor-
 oughly with soap and water. Change clothing immediately
 and wash before re-use. Destroy containers completely when

Our Name Is

Guarant

of

Quality

FOR USE ON	TO CONTROL	IN 100 GAL	IN 1 GAL	COMMENTS
Apples, Pears	European Red Mite Clover Mite Two Spotted Mite Woolly Apple Aphid Forbes Scale Green Apple Aphid Rosy Apple Aphid	1-2 pts 1 pt 1 pt 1 1/2 pts 1 1/2 pts	2 tsp 1 tsp 1 tsp 2 tsp 2 tsp	Make two or more applications as needed for mites, one or two for aphids. At the pink stage of the apple when aphids appear. Do not use within 2 days of harvest.
Tomatoes	Spider Mites, Aphids	1 1/2 pts 1 pt	1 1/2 tsp 1 tsp	Make one or more applications as needed. Do not use within 2 days of harvest.
Beans	Mexican Bean Beetle, Spider Mites, Leafhopper	2 pts	2 tsp	
Peas	Pea Aphid	2 pts	2 tsp	Make one or more applications. Do not use within 3 days of harvest. Do not use pea vines for forage.
Potatoes	Aphids	2 pts	2 tsp	Make full coverage applications. Repeat as necessary. Do not use within 2 days of harvest.
Cabbage, Broccoli, Kale, Mustard, Turnip	Aphids, Imported Cabbage Worm	2 pts	2 tsp	
Alfalfa, Clover, Sward	Aphids	2 pts at 10-15 day intervals	2 tsp	Do not use within 2 days of harvest.
Flowers, Ornamentals, Shrubs	Spider Mites, Aphids, White Fly, Thrips	1 1/2 pts 1 1/2 pts 1 1/2 pts	1 1/2 tsp 1 1/2 tsp 1 1/2 tsp	For mites, apply when needed. For aphids, apply when needed. For thrips, apply when needed.



50% MALATHION

Emulsifiable Concentrate

FOR CONTROL OF ANIMAL AND PLANT PESTS

ACTIVE INGREDIENTS:

Malathion* 50%
Solvent 45%

INERT INGREDIENTS

..... 5%
*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings

893-687

NET CONTENTS 1 QUART

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

DIRECTIONS

LIQUID FLY BAIT: 1 oz. per gallon of water. Apply around food disposals, etc. Mix with sugar syrup and add to 2 gallons of water. Apply to desired surfaces.

BEEF CATTLE: For control of ticks, apply 1 to 2 gallons of water. Apply a complete coverage spray at two week intervals if needed.

SWINE: For control of lice, apply a complete coverage spray. One treatment may be all that is needed.

BACK RUBBING DEVICES: Cattle, apply a 2% mixture of oil or kerosene to the back of the animal. This will be a reduction in horn flies one to 30 or 45 head of cattle a week. Do not use on dogs.

CAUTION: Do not use on dogs. Do not use on cats. Do not use on birds. Do not use on fish. Do not use on reptiles. Do not use on amphibians. Do not use on invertebrates.

WARNING: Do not use on dogs. Do not use on cats. Do not use on birds. Do not use on fish. Do not use on reptiles. Do not use on amphibians. Do not use on invertebrates.

May 26, 1971



DDT Concentrate Control of Insect Pests

50%
45%
5%
phosphate of diethyl

Reach of Children
for Additional Warnings

QUART

BY
CORPORATION
NEAS

DIRECTIONS FOR USE

LIQUID FLY BAIT: To control house flies, apply to barns, feed lots, poultry houses, etc., around food disposals. Mix 5 parts of concentrate with 95 parts of syrup and add to 2 gallons of water. Spray on desired surfaces.

BEEF CATTLE: For control of lice, apply 1 gallon of concentrate to 2 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed. For control of Ticks, apply 1 to 2 gallons per head of cattle. Apply a complete coverage spray. Repeat applications at two week intervals if needed.

SWINE: For control of lice, apply 1 gallon of concentrate to 2 gallons of water. Apply a complete coverage spray. One treatment may be sufficient. Repeat applications if needed.

BACK RUBBING DEVICES: For the control of lice on Cattle, apply a 2% mixture (1 part concentrate to 49 parts of oil or kerosene) to the back rubbing device. There will be a reduction in Horn Flies. Blanketing will result in one to 3 or 45 head of cattle. Repeat applications at 3 weeks. Do not use on dairy cattle.

CAUTION: Do not use on dairy cattle.
Do not use on poultry.
Do not use on fish.
Do not use on bees.
Do not use on honey bees.

WARNING: Do not use on dairy cattle.
Do not use on poultry.
Do not use on fish.
Do not use on bees.
Do not use on honey bees.

*Our Name Is Your
Guarantee*

Quality