




DAIRY and LIVESTOCK DUST



GORDON'S DAIRY AND LIVESTOCK DUST is a combination of two highly effective fly-control chemicals. To prevent the number of horn flies from increasing rapidly on dairy animals begin dusting animals in the spring when flies first appear.

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

DAIRY ANIMALS—HORN FLIES (on beef cattle also): Use 2 heaping tablespoonfuls (2 oz.) per animal for control of horn flies. Dust on poll neck, and back. Rub in lightly to insure that the material is worked down beneath the hair. If application is made by dust gun, apply against the grain of hair to force material into the hair. Do not repeat applications within a 3-week time period. Do not contaminate water troughs apply during lactation or within 2 weeks of freshening

BEEF CATTLE—LICE, TICKS. AND HORN FLIES: For control of lice. Licks and horn flies make application to all parts of the animal's body Rub into hair as application is made. Repeat tre-tment as necessary to control infestation. Repeat at 10 to 14 day intervals.

HOGS—LICE: Make application of dust to hogs, pens, and litter One treatment may be sufficient: if not, repeat in 10 days or thereafter as needed for control. May be dusted on baby pigs and farrowing-house litter. Apply ½ to ½ tablespoonful on pigs less than one month of age.

SHEEP—LICE AND KEDS: Make thorough application to entire animal Treatment just after shearing is preferred. Repeat as necessary for control.

CAUTION

Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed or foodstuffs, or feed or water troughs. Keep out of the reach of children.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product as specified in this label.

Do not reuse container. Destroy when empty.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

EPA REG. No. 2217-347-AA 516-1274

EPA FST No 2217-KS-1



DAIRY and LIVESTOCK DUST

Ready to Use

FOR HORN FLY CONTROL On Dairy and Beef Cattle and Sheep Also Controls Lice and Ticks

ACTIVE INGREDIENTS:	
Technical Methoxychlor*	5.0%
Malathion**	4.0%
INERT INGREDIENTS	91.0%
Total	
*Equivalent to 4.4% 2,2-bis (p-methoxyphenyl)-1,1,1-tric	chloro
ethane and 0.6% other isomers and related compound	is.

**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

(AUTION: Keep Out of Reach of Children

NET WEIGHT ONE POUND

GURDON'S DAIRY IND LALL JUNEAU from the state of the state of Commence of the Commence of th

BEIDIT OF WAS DIRECTORS

DAIRY ANIMALS AND AND THE REAL PRINCES ables, confus (1977) por an entre en COCK CONTROLL FOR A SECURITION OF THE SECURITION peneath the harmon transport grain of hair to be a second of the second withing Bankers 1. December 19 19 Comment

BEEF CATTLE And the seasoff of the second office of the service of the

HOGS THE TOWN the production of the second SHEEP

Avoid areathing of the control only a first and the washing a Agord contem now and to the contemporary timents koop out a die cost pagneta to a t

out conly where a m anditions to be deliced Samuel State of the Control of the C