

Best Available copy

ACCEPTED
MAY 10 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

2217-355

FOR USE ON	TO CONTROL	10 GAL.	100 GAL.	COMMENTS
Apples, Pear, Peach	European Red Mite 3-4 pts. Two Spotted Mite Woolly Apple Aphid 1 pt. Furrow Scale Green Apple Aphid 1 1/2 pts. Vasey Apple Aphid 1 1/2 pts.	2 lbs. 1 pt. 1 pt. 1 1/2 pts. 1 1/2 pts.	2 lbs. 1 pt. 1 pt. 1 1/2 pts. 1 1/2 pts.	Make two or more applications as needed for mites, one or more for aphids. At pre-bloom or peak bloom of blossoms or when aphids appear. Do not use within 7 days of harvest.
Timber	Sawyer, White, Ash-bark	1 1/2 pts. 1 pt.	1 1/2 lbs. 1 lb.	Make one or more applications as needed. Do not use within 3 days of harvest.
Beans	Mexican Bean Beetle, Solder Mites, Leafhopper	2 pts.	2 lbs.	
Peanut	Pea Aphid	2 pts.	2 lbs.	Make one or more applications. Do not use within 3 days of harvest. Do not use pea vines for forage.
Peas	Aphid	2 pts.	2 lbs.	
Cabbage, Broccoli, Kale, Mustard, Turnip	Aphid, Imported Cabbage Worm	2 pts.	2 lbs.	Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on Broccoli and Turnip. 2 days on others.
Alfalfa for Seed	Aphid	2 pts. per acre in at least 10 gal. water		Do not apply within 2 days of harvest, or when in bloom.
Cucumbers, Tomatoes, Squash	Cucumber Shoth Scale, Cucumber Scale Aphid, Spider Mite, White Fly, Thrips, Woolly Beg, Tobacco Root Aphid	1 pt. 1 1/2 pts.	1 lb. 1 1/2 lbs.	Apply in warm water when infestation occurs. Repeat in 10 to 12 days. For scales, apply when crawlers have settled on foliage.



50% MALATHION GARDEN SPRAY FOR CONTROL OF FLIES AND PLANT PESTS

ACTIVE INGREDIENTS:
Malathion* 50%
Xylene 45%
INERT INGREDIENTS 5%
*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings
E.P.A. Reg. No. 2217-355
NET CONTENTS 1 PINT

MANUFACTURED BY
GORDON CORPORATION
KANSAS CITY, KANSAS

DIRECTIONS FOR USE

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots, Poultry houses, Utility buildings and around food disposals mix 5 ounces with 5 ounces of corn syrup and add to 2 gallons of water. Paint or spray on desired surfaces.

WARNING: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist, avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before re-use. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

Malathion may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment. Do not spray when bees are active in area.

Our Name Is Your
Guarantee

Quality

2217-355

5-10-73

1081