
_
1
9
1
J
_ ^

FOR USE ON	TO CONTROL	IN 100 GAL.	IN 1 GAL	COMMENTS	
Apples, Pears,	European Red Mite Clover Mite Two Spotted Mite	1-2 pts.	2 tsp.	Make two or more ap- plications as needed for mites, one or more	
	Wooly Apple Aphid Forbes Scale	1 pt.	1 tsp.	for aphids. At pre-pink or pink stage of bloom	
	Green Apple Aphid Rosy Apple Aphid	142 pts.	14/2 tsp.	or when aphids appear. Do not use within 3 days of harvest.	
Tomatoes	Spider, Mites, Aphids	1½ pts. 1 pt.	1½ tsp. 1 tsp.	Make one or more ap- plications as needed.	
Beans	Mexican Bean Beet Spider Mites, Leafhopper	ie, 2 pts.			
Peas	Pea Aphid	2 pts.	2 tsp.	Make one or more applications. Do not use within 3 days of harvest. Do not use pea vines for forage.	
Potatoes	Aphids	2 pts.	2 tsp.		
Cabbage, Broccoli, Kale, Mustard, Turnip	Aphids Imported Cabbage Worm	2 pts.	2 tsp.	Make full coverage applications. Repeat an encessary. Do not use within 3 days of harvest on Broccoli and Turnips, 7 days on others.	
Alfalfa for Seed	Aphids,		er acre in 10 gal.	Do not apply within 7 days of harvest, or when in bloom.	
Flowers, Ornamentals, Shrubs	Oyster Shell Scale Euonymus Scale	1 pt.	1 tsp.	Apply in summer when infestation occurs. Re-	
	Aphids, Spider Mites, White Fly, Thrips, Mealy Bugs, Japanese Beetle Adi Four Lined Plant Bug Tarnished Plant Bug Rose Leafhopper	peat in 10 to 12 days. For scales, apply when crawlers have settled on foliage.			
	Potato Leathopper Bagworms, Juniper	2 pts.	2 tsp.		
	Scale				
	Soft Brown Scale	21/2 pts.	21/2 tsp.		
	Pine Needle Scale	4 pts.	4 tsp.		

CAUTION: Do not use on certain ferns, including Boston maidenhair and Pteris, as well as some species of Crassula, because of possible injury to these ornamentals.



50% MALATHION GARDEN SPRAY



DIRECTIONS FOR USE

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots. Poultry houses. Utility buildings and around food disposals, mix 5 ounces with 5 ounces of corn syrup and add to 2 gallons of water. Paint or spray on desired surfaces.

WARNING: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist, avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before re-use. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

Malathion may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish. Keep out of lakes, streams, or ponds, Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment. Do not spray when bees are active in area.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Do not re-use container. Destroy when empty. Apply this product only as specified on this label.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claim other than those stated on this label.

893-973

E.P.A. Reg. No. 2217-355-AA

ACCEPTED JAN 1 0 1974

UNDER THE FEDERAL INSECTICIDE FUNCTOIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER. ED UNDER NO.

