

DIRECTIONS FOR USE

APPLES & PEARS: To control European Red Mite, Clover Mite, and Two Spotted Mite, mix 2 pts. in 100 gals. water or 2 tsp. in 1 gal. water. To control Wooly Apple Aphid and Forbes Scale, mix 1 pt. in 100 gals. water or 1 tsp. in 1 gal. water. To control Green Apple Aphid and Rosy Apple Aphid, mix 1½ pts. in 100 gals. water or 1½ tsp. in 1 gal. water. Make two or more applications as needed for mites, one or more for aphids. At pre-pink or pink stage of bloom or when aphids appear. Do not use within 3 days of harvest. This product may injure foliage on some varieties of apples and pears.

TOMATOES: To control spiders and mites, mix 1½ pts. in 100 gals. water or 3 tsp. in 1 gal. water. To control aphids, mix 1 pt. in 100 gals. water or 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest.

BEANS: To control Mexican Bean Beetle, Spider Mites, Leafhopper, and aphids, mix 2 pts. in 100 gals. water or 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest.

PEAS & POTATOES: To control Pea Aphid on peas or aphids on potatoes, mix 2 pts. in 100 gals. water or 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest. Do not use pea vines for forage.

CABBAGE, BROCCOLI, KALE, MUSTARD, & TURNIP: To control aphids, Imported Cabbage Worm, mix 2 pts. in 100 gals. water or 3 tsp. in 1 gal. water. Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on broccoli and turnips, 7 days on others.

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots, Poultry Houses, Utility buildings and around food disposals, mix 5 ounces with 5 ounces of corn syrup and add to 2 gals. water. Paint or spray on desired surfaces.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application, equipment, will vary, Seller makes no claims other than those stated on this label.

DIRECTIONS FOR USE

FLOWERS, ORNAMENTALS, & SHRUBS: To control Oyster Shell Scale and Euonymus Scale mix 1 pt. in 100 gals. water or 1 tsp. in 1 gal. water. To control aphids, Spider Mites, White Fly, Thrips, Mealy bugs, Japanese Beetle Adult, Four Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, and Potato Leafhopper, mix 1½ pts. in 100 gals. water or 1½ tsp. in 1 gal. water. To control Bagworms and Juniper Scale, mix 2 pts. in 100 gals. water or 2 tsp. in 1 gal. water. To control Soft Brown Scale mix 2½ pts. in 100 gals. water or 2½ tsp. in 1 gal. water. To control Pine Needle Scale mix 4 pts. in 100 gals. water or 4 tsp. in 1 gal. water. Apply in summer when infestation occurs. Repeat in 10 to 12 days. For scales, apply when crawlers have settled on foliage.

NOTE THESE CAUTIONS—Do not use on certain ferns, including Boston, Maidenhair and Pteris, as well as some species of Crassula, because of possible injury to these ornamentals.

WARNING: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist, avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before re-use. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

Malathion may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

893-674

FOR USE

Red Mite, Clover Mite, and Two water or 2 tsp. in 1 gal. water. To Scale, mix 1 pt. in 100 gals. water Green Apple Aphid and Rosy Apple or 1½ tsp. in 1 gal. water. Make for mites, one or more for aphids. or when aphids appear. Do not use may injure foliage on some varieties

es, mix 1½ pts. in 100 gals. water aphids, mix 1 pt. in 100 gals. water or more applications as needed. Do

tle, Spider Mites, Leafhopper, and or 3 tsp. in 1 gal. water, Make one ot use within 3 days of harvest.

aid on peas or aphids on potatoes, in 1 gal water. Make one or more thin 3 days of harvest. Do not use

& TURNIP: To control aphids, im- 100 gals. water or 3 tsp. in 1 gal. . Repeat as necessary. Do not use nd turnips, 7 days on others.

d Stable Flies in Barns, Feed Lots, round food disposals, mix 5 ounces d to 2 gals. water. Paint or spray

product conforms to the ingredient ons of use, such as weather, com- conditions of application, equip- ns other than those stated on this

EPA EST. NO. 2217-KS-1



50% MALATHION GARDEN SPRAY

FOR CONTROL OF

MANUFACTURED BY
PBI - GORDON CORPORATION

KANSAS CITY, KANSAS 64118

DIRECTIONS FOR USE

On all vegetables listed below, mix 3 teaspoonfuls in one gallon water.

TOMATOES: To control spiders, mites, and aphids, spray as needed. Do not use within 3 days of harvest.

BEANS: To control Mexican Bean Beetle, Spider Mites, Leafhopper, and aphids, spray as needed. Do not use within 3 days of harvest.

PEAS & POTATOES: To control Pea Aphid on peas or aphids on potatoes, spray as needed. Do not use within 3 days of harvest.

CABBAGE, BROCCOLI, KALE, MUSTARD, & TURNIP: To control aphids and Imported Cabbage worms, spray as needed. Do not use within 3 days of harvest on broccoli and turnips, 7 days on others.

FLOWERS, ORNAMENTALS, & SHRUBS: Mix 3 teaspoonfuls in one gallon water. Make application when insects appear, and repeat as necessary to control aphids, mealy bugs, spider mites, white flies, and leafhoppers. To control scale on euonymus, juniper, and pine, spray in May. To control bagworm larvae, spray as soon as insects are seen and repeat in 10 to 14 days. Do not use on Boston or Maiden hair ferns, pteris, crassula, or canaerti juniper as injury may result.

EPA REG. NO. 2217-355-AA EPA EST. NO. 2217-KS-1



50% MALATHION GARDEN SPRAY

FOR CONTROL OF

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS 66118

WARNING: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist, avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before re-use. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

Malathion may cause cholinesterase inhibition. If poison occurs, get prompt medical aid. Atropine is antidotal.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment, will vary, Seller makes no claims other than those stated on this label.

893-674

DIRECTIONS FOR USE

On all vegetables, mix 3 teaspoonfuls in one gallon.

TOMATOES: To control mites, and aphids, spray as needed. Do not use within 3 days of harvest.

BEANS: To control Mexican Bean Beetle, Spider Mites, Leafhoppers, and aphids, spray as needed. Do not use within 3 days of harvest.

PEAS & POTATOES: To control Pea Aphid on peas or aphids on potatoes, spray as needed. Do not use within 3 days of harvest.

CABBAGE, BROCCOLI, KALE, MUSTARD, & TURNIP: To control aphids and Imported Cabbage worms, spray as needed. Do not use within 3 days of harvest on broccoli and turnips, 7 days on others.

FLOWERS, ORNAMENTALS, & SHRUBS: Mix 3 teaspoonfuls in one gallon water. Make application when insects appear and repeat as necessary to control aphids, mealy bugs, spider mites, white flies, and leafhoppers. To control scale on euonymus, juniper, and pine, spray in May. To control bagworm larvae, spray as soon as insects are seen and repeat in 10 to 14 days. Do not use on Boston or Maiden hair ferns, pteris, crassula, or canaerti juniper as injury may result.

EPA REG. NO. 2217-355-AA EPA EST. NO. 2217-KS-1



50% MALATHION GARDEN SPRAY

FOR CONTROL OF

MANUFACTURED BY
FBI - GORDON CORPORATION
KANSAS CITY, KANSAS 64108

WARNING: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist, avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before re-use. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

Malathion may cause cholinesterase inhibition. If poison occurs, get prompt medical aid. Atropine is antidotal.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment, will vary, Seller makes no claims other than those stated on this label.

893-674