

BACK RUBBER USE ON DAIRY AND BEEF CATTLE:
In back rubbers for horn flies, or face treaters for face flies, make a 1% solution by diluting 1 pint with 2 1/4 gallons of fuel oil, kerosene or mineral seal oil.

The face rubber mechanism must be so designed that one end is low enough (not more than 15 to 20 inches above the ground) to allow cattle to rub their faces. Sheet canvas type face treaters should be located at barn door openings, in loafing areas or around salt licks where animals may freely treat themselves as needed for face fly control. This product is only effective for face fly control on those cattle that rub the face and head on the face treater.

WARNING: Hazardous if swallowed, inhaled, or absorbed through skin. Wash thoroughly with soap and water after handling, and before eating or smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Do not contaminate feed or foodstuffs.

During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic



CIODRIN® 150



EMULSIFIABLE CONCENTRATE

CONTROLS FLIES ON BEEF AND DAIRY CATTLE

ACTIVE INGREDIENTS:

*Dimethyl Phosphate of Alpha-Methyl-benzyl 3-Hydroxy-Cis-Crotonate	13.1%
Aromatic Petroleum Hydrocarbons	76.9%
INERT INGREDIENTS	10.0%
Total	100.0%

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings

951-973

NET CONTENTS ONE PINT

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS



CIODRIN® 150



EMULSIFIABLE CONCENTRATE

CONTROLS FLIES ON BEEF AND DAIRY CATTLE

ACTIVE INGREDIENTS:

*Dimethyl Phosphate of Alpha-Methyl-
benzyl 3-Hydroxy-Cis-Crotonate

13.1%

Aromatic Petroleum Hydrocarbons

76.9%

INERT INGREDIENTS

10.0%

Total

100.0%

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings

951-973

NET CONTENTS ONE PINT

MANUFACTURED BY

PBI - GORDON CORPORATION

KANSAS CITY, KANSAS

DIRECTIONS

FLY CONTROL: For horn flies, face flies, house flies, stable flies, lice and the Lone Star tick, on dairy cattle and beef cattle, mix 1 pint with 6 to 12 gallons of water. Thoroughly apply from 2 to 3 quarts of diluted material per animal. Do not apply more often than once per week. Do not contaminate food or feed. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

PREMISE SPRAY FOR FLIES: As an aid in reducing populations of house flies and stable flies around barns, on walls, eaves, fences, in loafing sheds and corrals, mix 2 pints in 3 gallons of water. Use at the rate of one gallon of diluted material per 1,000 square feet of surface. Thoroughly treat. Repeat later as needed for control. Remove animals during treatment.

LICE AND TICKS: For application in fall and winter on dairy cattle, beef cattle, sheep, goats and hogs, mix 1 pint in 6 to 12 gallons of water and thoroughly

BACK RUBBER USE ON DAIRY AND BEEF CATTLE: In back rubbers for horn flies, or face treaters for face flies, make a 1% solution by diluting 1 pint with 2 1/4 gallons of fuel oil, kerosene or mineral seal oil.

The face rubber mechanism must be so designed that one end is low enough (not more than 15 to 20 inches above the ground) to allow cattle to rub their faces. Sheet canvas type face treaters should be located at barn door openings, in loafing areas or around salt licks where animals may freely treat themselves as needed for face fly control. This product is only effective for face fly control on those cattle that rub the face and head on the face treater.

WARNING: Hazardous if swallowed, inhaled, or absorbed through skin. Wash thoroughly with soap and water after handling, and before eating or smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Do not contaminate feed or feedstuffs.

During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type found suitable by the U.S. Bureau of Mines for protection against Clodrin insecticide. Use this material only for recommended purposes and at recommended dosages.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

WARNING! FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.

951-973

ClODRIN[®] 150



EMULSIFIABLE CONCENTRATE

CONTROLS FLIES ON BEEF AND DAIRY CATTLE

ACTIVE INGREDIENTS

1,1-Dichloro-2,2-bis(4-chlorophenyl)ethane

0.5% (0.5% of 150)

Other inert ingredients

99.5% (99.5% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

100% (100% of 150)

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings

951-973

NET CONTENTS ONE PINT

DIRECTIONS

FLY CONTROL: For horn flies, face flies, house flies, stable flies, lice and the Lone Star tick, on dairy cattle and beef cattle, mix 1 pint with 6 to 12 gallons of water. Thoroughly apply from 2 to 3 quarts of diluted material per animal. Do not apply more often than once per week. Do not contaminate food or feed. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

PREMISE SPRAY FOR FLIES: As an aid in reducing populations of house flies and stable flies around barns, on walls, eaves, fences, in loafing sheds and corrals, mix 2 pints in 3 gallons of water. Use at the rate of one gallon of diluted material per 1,000 square feet of surface. Thoroughly treat. Repeat later as needed for control. Remove animals during treatment.

LICE AND TICKS: For application in fall and winter on dairy cattle, beef cattle, sheep, goats and hogs, mix 1 pint in 6 to 12 gallons of water and thoroughly apply to cattle at the rate of one gallon of diluted spray per animal with similar reapplication 14 days later. Use approximately one-half gallon diluted spray per head on small animals and repeat this treatment also.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 2217-360-AA

951-973

CIDDRIN[®] 150



EMULSIFIABLE CONCENTRATE

CONTROLS FLIES ON BEEF AND DAIRY CATTLE

ACTIVE INGREDIENTS

1,1-Dichloro-2,2-bis(4-chlorophenyl)ethane 15.0%

INERT INGREDIENTS 85.0%

NET CONTENTS

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warnings

951-97

NET CONTENTS ONE PINT

DIRECTIONS

FLY CONTROL: For horn flies, face flies, house flies, stable flies, lice and the Lone Star tick, on dairy cattle and beef cattle, mix 1 pint with 6 to 12 gallons of water. Thoroughly apply from 2 to 3 quarts of diluted material per animal. Do not apply more often than once per week. Do not contaminate food or feed. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

PREMISE SPRAY FOR FLIES: As an aid in reducing populations of house flies and stable flies around barns, on walls, eaves, fences, in loafing sheds and corrals, mix 2 pints in 3 gallons of water. Use at the rate of one gallon of diluted material per 1,000 square feet of surface. Thoroughly treat. Repeat later as needed for control. Remove animals during treatment.

LICE AND TICKS: For application in fall and winter on dairy cattle, beef cattle, sheep, goats and hogs, mix 1 pint in 6 to 12 gallons of water and thoroughly apply to cattle at the rate of one gallon of diluted spray per animal with similar reapplication 14 days later. Use approximately one-half gallon diluted spray per head on small animals and repeat this treatment also.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 2217-360-AA

951-973

Use Name I