

ACCEPTED

Oct 12 - 1966

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 2217-351 SUBJECT TO ATTACHED COMMENTS.

SILVEX LAWN WEED KILLER



KILLS CHICKWEED, DANDELION, PLANTAIN AND MANY OTHER WEEDS.

8 Ounces Makes 8 Gallons of Dilute Spray
ACTIVE INGREDIENTS:
*Mixed ethyl ester of 2,4-Dichloro-phenoxyacetic acid 14.0%
**Mixed ethyl ester of Silvex 2-(2,4,5-Trichlorophenoxy)propionic acid) 4.3%
INERT INGREDIENTS 81.7%
Total 100.0%

*Equivalent to 9.2% 2,4-Dichlorophenoxyacetic acid
**Equivalent to 3.0% Silvex 2-(2,4,5-Trichlorophenoxy)propionic acid)

CAUTION: Keep Out of Reach of Children
See Side Panel for Additional Cautions
841-663 CONTENTS ONE PINT

Packed for
FARMERS UNION CENTRAL EXCHANGE, INC.
St. Paul, Minn.

DIRECTIONS FOR USE

MIXING DIRECTIONS—FOR HOSE SPRAYERS AND GARDEN PRESSURE SPRAYERS, MIX 2 TABLESPOONS PER GALLON OF WATER APPLIED. A SMALL AMOUNT OF LIQUID DETERGENT MAY BE MIXED WITH SPRAY TO HELP WET THE WEEDS.

WEEDS KILLED—Common and muscled chickweed, chicory, clover, dandelion, dock, henbit, plantain, ragweed, thistle, and wild onion. Do not use on dichondra or lippla lawns. Make two applications for plants such as thistle, bindweed, wild onion, spotted spurge and knotweed or as needed for control.

WHEN TO SPRAY—For chickweed, treat in the fall before seeds are produced (October). Repeat if needed for complete kill. For other susceptible weeds listed, spray in late spring when weeds are producing new growth or as needed through the season. Reseeding can be done within 3 to 4 weeks. Do not spray new lawns until after second cutting.

SPRAYING—Apply 1 gallon of dilute spray to 100 sq. feet of turf. To get best results, area treatment spray preferred over spot spraying. Spray again if rain occurs before it has dried on weeds. Do not use on bent grass, buffalo, carpet grass or St. Augustine lawns since damage will result.

841-663 U.S.D.A. Reg. No. 2217 351

DIRECTIONS FOR USE

WOODY PLANTS—For poison ivy in fence row and vacant lots, mix 8 ounces in 2 1/2 gallons of water. Use a garden pressure sprayer. Add 1 tablespoon of liquid detergent. Thoroughly wet ivy. Do not wet ivy climbing up tree trunks, spray sparingly and repeat as needed. Repeat 2 weeks later if needed. For stump resprouting, mix 2 ounces in 1 quart of kerosene. Soak cut surfaces, especially near bark and base of stump.

CAUTION—Avoid contact with skin, eyes and clothing. Harmful if swallowed. Do not use in greenhouses. Vapors from this product may injure susceptible plants in immediate vicinity. Do not contaminate water used for irrigation or domestic purposes. Do not store with or near seeds, fertilizers, insecticides or fungicides. Do not store in confined spaces with living plants. Do not allow spray to drift to nearby susceptible plants such as grapes, garden vegetables, roses or other flowers. Coarse sprays are less likely to drift. Do not use hose applicators when near susceptible plants.

NON-WARRANTY—Seller makes no warranty, expressed or implied regarding this product except the composition as set forth in the ingredient statement herein. Buyer and user assumes all risk of possession, handling or use, whether or not in accordance with any directions contained herein.
841-663

Side ->
pots

4)