

DIRECTIONS FOR USE

Use SEVIN 50W INSECTICIDE as a spray. Keep mixtures well agitated. Not effective if used with alkaline chemicals such as Bordeaux, but is compatible with most all fungicides and miticides. Spray thoroughly to cover upper and lower sides of foliage, fruits and vegetables. Do not spray open blossoms since SEVIN is toxic to honey bees. For continued insect control through the season, plan on a regular spray schedule every 7 to 10 days on all fruits.

APPLES AND PEARS—codling moth, plum curculio, apple aphids, green fruit worms, Japanese beetles, leaf rollers and San Jose scale—As insects appear, spray, using 2 tablespoonfuls per gallon of water, repeat as needed for control, up to one day before harvest. To avoid fruit thinning, delay use until at least 30 days after bloom (not before second cover).

PEACHES, APRICOTS, PLUMS AND CHERRIES—plum curculio, Oriental fruit moth, lesser peach tree borer, peach twig borer, cat facing bugs, and San Jose scale—Mix 2 tablespoonfuls per gallon of water. Allow one day between spraying and harvest; three days on apricots.

SWEET CORN—corn earworm, corn borer, flea beetles and grasshoppers—use 2 tablespoonfuls per gallon. Spray every 2 to 3 days for earworms until silks are dry. For other pests, thoroughly treat entire plant every 2 weeks, starting when corn is knee high.

GRAPES—Use 2 tablespoonfuls per gallon for grape leaf folder, grape leaf skeletonizer, grape berry moth, and June beetles. Apply as needed when insects appear. Use up until harvest time.

STRAWBERRIES—Use 2 tablespoonfuls per gallon for meadow spittlebug, strawberry leaf roller and strawberry weevil. Repeat at 7 to 10 day intervals, or as necessary until one day before harvest.

BEANS—green beans, lima and snap. Use 2 tablespoonfuls per gallon for Mexican bean beetle, bean leaf beetle, flea beetle and stink bugs. Use up until harvest time.

TOMATOES, EGG PLANTS, PEPPERS—For tomato hornworm and fruitworm, Fall armyworm, stink bugs and lace bugs, use 2 tablespoonfuls per gallon. Thoroughly treat. Use until day of picking.

EP-477



SEVIN* 50W INSECTICIDE

MANUFACTURED BY
PEI - GORDON CORPORATION
KANSAS CITY, KANSAS

DIRECTIONS FOR USE

CUCUMBER, MELONS AND SQUASH—For pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs, use 2 tablespoonfuls per gallon. Avoid excessive applications since leaf injury may occur on tender foliage when wet and humidity is high. Allow 3 days before harvest.

CABBAGE, BROCCOLI AND CAULIFLOWER—For control of small cabbage loopers, flea beetles, armyworms and corn earworms—use 2 tablespoonfuls per gallon on a 7 to 10 day spray schedule, allowing 3 days before picking.

ORNAMENTALS—On plants such as roses, gladiolus, lilacs, juniper, elm, boxelder and willow, use 2 tablespoonfuls per gallon and apply when insects appear. Controls bagworms, blister beetles, boxelder bugs, elm leaf beetle, lute beetles, leaf rollers, mimosa webworm, periodical cicada, scale insects and willow leaf beetles. Do not use on Boston ivy or Virginia creeper. Repeat as needed.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust or spray mist. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed or food stuffs. Keep out of reach of children.

Rinse empty container thoroughly with water and discard it. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not use in any manner not specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

EP-873

E.P.A. Reg. No. 2217-366 AA

DIRECTIONS FOR USE

Use SEVIN 50W INSECTICIDE as a spray. Keep mixtures well agitated. Not effective if used with alkaline chemicals such as Bordeaux, but is compatible with most all fungicides and miticides. Spray thoroughly to cover upper and lower sides of foliage, fruits and vegetables. Do not spray open blossoms since SEVIN is toxic to honey bees. For continued insect control through the season, plan on a regular spray schedule every 7 to 10 days on all fruits.

APPLES AND PEARS—codling moth, plum curculio, apple aphids, green fruit worms, Japanese beetles, leaf rollers and San Jose scale—As insects appear, spray, using 2 tablespoonfuls per gallon of water, repeat as needed for control, up to one day before harvest. To avoid fruit thinning, delay use until at least 30 days after bloom (not before second cover).

PEACHES, APRICOTS, PLUMS AND CHERRIES—plum curculio, Oriental fruit moth, lesser peach tree borer, peach twig borer, cat facing bugs, and San Jose scale—Mix 2 tablespoonfuls per gallon of water. Allow one day between spraying and harvest; three days on apricots.

SWEET CORN—corn earworm, corn borer, flea beetles and grasshoppers—Use 2 tablespoonfuls per gallon. Spray every 2 to 3 days for earworms until silks are dry. For other pests, thoroughly treat entire plant every 2 weeks, starting when corn is knee high.

GRAPES—Use 2 tablespoonfuls per gallon for grape leaf folder, grape leaf skeletonizer, grape berry moth, and June beetles. Apply as needed when insects appear. Use up until harvest time.

STRAWBERRIES—Use 2 tablespoonfuls per gallon for meadow spittlebug, strawberry leaf roller and strawberry weevil. Repeat at 7 to 10 day intervals, or as necessary until one day before harvest.

BEANS—green beans, lima and snap—Use 2 tablespoonfuls per gallon for Mexican bean beetle, bean leaf beetle, flea beetle and stink bugs. Use up until harvest time.

TOMATOES, EGG PLANTS, PEPPERS—For tomato hornworm and fruit worm, fall armyworm, stink bugs and lace bugs, use 2 tablespoonfuls per gallon. Thoroughly treat. Use up until time of picking.



SEVIN* 50W INSECTICIDE

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

DIRECTIONS FOR USE

CUCUMBER, MELONS AND SQUASH—For pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs, use 2 tablespoonfuls per gallon. Avoid excessive applications since leaf injury may occur on tender foliage when wet and humidity is high. Allow 3 days before harvest.

CABBAGE, BROCCOLI AND CAULIFLOWER—For control of small cabbage loopers, flea beetles, armyworms and corn earworms—use 2 tablespoonfuls per gallon on a 7 to 10 day spray schedule, allowing 3 days before picking.

ORNAMENTALS—On plants such as roses, gladiolus, lilacs, juniper, elm, boxelder and willow, use 2 tablespoonfuls per gallon and apply when insects appear. Controls bagworms, blister beetles, boxelder bugs, elm leaf beetle, June beetles, leaf rollers, mimosa webworm, periodical cicada, scale insects and willow leaf beetles. Do not use on Boston ivy or Virginia creeper. Repeat as needed.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust or spray mist. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed or food stuffs. Keep out of reach of children.

Rinse empty container thoroughly with water and discard it. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not use in any manner not specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

620-873

E.P.A. Reg. No. 2217-266 AA