

SEVIN® 50W INSECTICIDE

Controls Tomato Hornworm, Bean Beetle, Aphids,
Squash Bug, Scale, Fruit and Ornamental Pests.


ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)	50.0%
INERT INGREDIENTS	50.0%
TOTAL	100.0%

SEVIN® is the registered trademark of Rhone Poulenc for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN


CAUTION



See side panel for Precautionary Statements
and Statement of Practical Treatment

NET WEIGHT __ OUNCES

620 APXXXXXX
 EPA REG. NO. 2217-366
 EPA EST. NO. 2217-KS-1
 Mfd. by PBI/GORDON CORPORATION
 KANSAS CITY, MISSOURI 64101



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid breathing vapors [dust or spray mist]. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

- User Safety Requirements:**
- ◆ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
 - ◆ Wear a hat and eye protection when making overhead applications.
 - ◆ Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
 - ◆ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
 - ◆ Do not allow children or pets to contact treated area until sprays have dried.

Statement of Practical Treatment

IF SWALLOWED: Induce vomiting and seek medical attention immediately.

IF IN EYES OR ON SKIN: Flush eyes thoroughly with water, wash skin thoroughly with soap and water.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

Note to Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine sulfate is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS:

This pesticide is extremely toxic to aquatic and estuarine invertebrates. ~~For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.~~ Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or animals.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

Use Sevin® 50W Insecticide as a spray. Keep mixtures well agitated. Not effective if used with alkaline chemicals such as Bordeaux, but is compatible with most all fungicides and miticides. Spray thoroughly to cover upper and lower sides of foliage, fruits and vegetables.

Do not spray open blossoms since Sevin® is toxic to honey bees. For continued insect control through the season, plan on a regular spray schedule every 7 to 10 days on all fruits. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

APPLES AND PEARS: *codling moth, plum curculio, apple aphid, green fruitworms, Japanese beetle, leaf rollers, and San Jose scale* - As insects appear, spray, using ~~Mix~~ 2 tablespoons of product per gallon of water. repeat as needed for control, up to one day before harvest. To avoid fruit thinning, delay use until at least 30 days after bloom (not before second cover). For trees taller than 10 feet consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week. Allow one (1) day between spraying and harvests.

PEACHES, APRICOTS, PLUMS, AND CHERRIES: *plum curculio, Oriental fruit moth, lesser peach tree borer, peach twig borer, "cat-facing" bugs, and San Jose scale* - Mix 2 tablespoons of product per gallon of water. Allow one (1) day between spraying and harvest; three (3) days on Apricots. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

SWEET CORN: *corn earworm, corn borer, flea beetle, and grasshoppers* - Mix 2 tablespoons of product per gallon of water. Spray every 2 to 3 days for earworms until silks are dry. For other pests, thoroughly treat entire plant every 2 weeks, starting when corn is knee high.

GRAPES: Mix 2 tablespoons of product per gallon of water for *grape leaf folder, grape leaf skeletonizer, grape berry moth, and June beetle*. Apply as needed when insects appear. Use up until Allow 7 days before harvest. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

STRAWBERRIES: Mix 2 tablespoons of product per gallon of water for *meadow spittlebug, strawberry leaf roller and strawberry weevil*. Repeat at 7 to 10 day intervals, or as necessary until Allow one day before harvest. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

BEANS, green beans, lima and snap: Mix 2 tablespoons of product per gallon of water for *Mexican bean beetle, bean leaf beetle, flea beetle and stink bugs*. Use up until harvest. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

TOMATOES, EGGPLANTS, PEPPERS: For *tomato hornworm and fruitworm, fall armyworm, stink bugs and lace bug*, mix 2 tablespoons of product per gallon of water. Thoroughly treat. Use until day of picking. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

CUCUMBERS, MELONS AND SQUASH: For *pickleworm, melonworm, cucumber beetle, flea beetle, leafhoppers, and squash bug*, mix 2 tablespoons of product per gallon of water. Avoid excessive applications since leaf injury may occur on tender foliage when wet and humidity is high. Allow 3 days before harvest. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

CABBAGE, BROCCOLI AND CAULIFLOWER: For control of *small cabbage looper, flea beetle, armyworm and corn earworm* - mix 2 tablespoons of product per gallon of water on a 7 to 10 day spray schedule. Allowing 3 days before picking. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

ORNAMENTALS: On plants such as roses, gladiolus, lilacs, juniper, elm, boxelder and willow, mix 2 tablespoons of product per gallon of water. Apply when insects appear. *Controls bagworm, blister beetle, boxelder bug, elm leaf beetle, June beetle, leaf roller, mimosa webworm, periodical cicada, scale insects and willowleaf beetle.* Do not use on Boston ivy or Virginia creeper. Repeat as needed. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of products or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

