

DIRECTIONS FOR USE

GENERAL INFORMATION:

St. Augustine Broadleaf Weedkiller is specially formulated for maximum safety to 2,4-D sensitive grasses such as St. Augustine grass, bentgrass and buffalo grass. Sprays should be applied when broadleaf weeds are small and tender for best results. If weeds are older and hard to kill the higher rates recommended should be used. Avoid spraying St. Augustine grass under conditions of high temperature and moisture when the St. Augustine is growing rapidly. Under these conditions, or in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage might occur.


To ensure weeds are growing and able to absorb the herbicide irrigate the lawn 2 or 3 days prior to spraying. Do not water for 24 hours after treatment. Although spray works faster with temperatures above 70° F, sprays can be applied at any time plants are growing. Spray when air is calm to avoid drift. Spray will injure or kill clover, dichondra and most ornamental and vegetable crops.

Amount of water in spray is not important except to evenly distribute the chemical. If directions call for 1 oz. in 1 gallon on 400 square feet it may be more convenient to use 1 oz. in 2 gallons on 400 square feet depending on the sprayer. Generally large volumes of water result in more uniform coverage.

HOW TO MIX:

1. For use on St. Augustine grass, bentgrass and buffalo grass, mix 1 oz. in 1 gallon of water and uniformly spray 400 square feet, for example, an area 10 feet wide and 40 feet long. If necessary, repeat treatment in 2 to 3 weeks, if hard to kill weeds persist.
2. For use on all other grasses such as bermuda, bahia, bluegrass, centipede, fescue, rye, mix 1 1/2 oz. in 1 gallon of water and spray uniformly over 400 square feet. On hard to kill species, or under poor growing conditions, or on older weeds, mix 2 oz. in 1 gallon of water and spray on 400 square feet with special attention to thorough coverage of the hard to kill weeds.

880-672



ST. AUGUSTINE GRASS BROADLEAF HERBICIDE

Contains *TRIMEC® Herbicide
U. S. Patent No. 3,284,186

Kills: DANDELIONS KNOTWEED HENBIT
CHICKWEED PLANTAINS SPURGE
and most other broadleaf weeds

ONE QUART COVERS UP TO 12,800 SQUARE FEET

KEEP FROM FREEZING

ACTIVE INGREDIENTS:	
*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	3.23%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	10.59%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	1.28%
INERT INGREDIENTS	84.90%
TOTAL	100.00%
*2,4-dichlorophenoxyacetic acid equivalent 0.235 pounds per gallon or 2.69%	
**2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.764 pounds per gallon or 8.76%	
***3,6-dichloro-o-anisic acid equivalent 0.094 pounds per gallon or 1.07%	

CAUTION: Keep Out of Reach of Children

See Back Panel for Additional Cautions

*Trimec® is a registered trademark of P.B.I.—Gordon Corporation
U.S. Patent No. 3,284,186.

CONTENTS 1 GALLON

880-672

35

MANUFACTURED BY

PBI - GORDON CORPORATION

KANSAS CITY, KANSAS

66118

WEEDS CONTROLLED: St. Augustine Herbicide is especially effective on weeds such as SHEEP SORE, SPOTTED SPURGE, CHICKWEED AND HENBIT. In addition, it controls virtually all broadleaf weeds listed below.

Bedstraw	Henbit
Black Medic	Knotweed
Buckhorn	Lambquarters
Bull Thistle	Lespedeza
Burdock	Mallow
Buttercup	Morning Glory
Chicory	Mustard
Chickweed	Oxalis
Clover	Peppergrass
Cocklebur	Pigweed
Dandelion	Plantains
Dock	Poison Ivy
Galinsoga	Poison Oak
Ground Ivy	Purslane
Hawkweed	Ragweed
Heal-all	Sheep Sore

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with plenty of water.

Avoid drift of spray on flowers, ornamental plants, other desirable plants, domestic or irrigation wells. Do not contact with fertilizers, herbicides or fungicides. Use only for the purpose intended. Do not apply to roots of desirable trees.

When container is empty, wash thoroughly and destroy. Never reuse for anything else. Do not burn.

Under certain conditions, this herbicide may occur to bentgrass, St. Augustine grass.

NOTICE: Seller warrants that this herbicide conforms to the ingredients listed on the label. Since conditions of use, weather, compatibility and condition of application may vary, Seller makes no warranty other than those stated on this label.

EPA Reg. No. 2217-576



ACCEPTED
July 7, 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECOTOXICITY AND POISON REGISTER-
 ED AT 2217-576 SUBJECT
 TO ACTION COMMENTS.

ST. AUGUSTINE GRASS BROADLEAF HERBICIDE

Contains *TRIMEC® Herbicide
 U. S. Patent No. 3,284,186

Kills: DANDELIONS KNOTWEED HENBIT
 CHICKWEED PLANTAINS SPURGE
 and most other broadleaf weeds

ONE QUART COVERS UP TO 12,800 SQUARE FEET

KEEP FROM FREEZING

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichloro- phenoxyacetic acid	3.23%
**Dimethylamine salt of 2-(2-methyl-4- chlorophenoxy) propionic acid	10.59%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	1.28%
INERT INGREDIENTS	84.90%
TOTAL	100.00%
*2,4-dichlorophenoxyacetic acid equivalent 0.235 pounds per gallon or 2.69%	
**2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.764 pounds per gallon or 8.76%	
***3,6-dichloro-o-anisic acid equivalent 0.094 pounds per gallon or 1.07%	

CAUTION: Keep Out of Reach of Children
 See Back Panel for Additional Cautions

*Trimec® is a registered trademark of P.B.I.—Gordon Corporation
 U.S. Patent No. 3,284,186.

CONTENTS 1 GALLON

880-672

MANUFACTURED BY

PBI - GORDON CORPORATION

KANSAS CITY, KANSAS

66118

WEEDS CONTROLLED: St. Augustine Grass Herbicide is especially effective against weeds such as SHEEP SORREL, KNOT WEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Bedstraw	Henbit	Shepherd's
Black Medic	Knotweed	Purse
Buckhorn	Lamb-quarter	Speedwell
Bull Thistle	Lespedeza	Tansy Mustard
Burdock	Mallow	Thistle
Buttercup	Morning Glory	Spurge
Chicory	Mustard	Velvet Leaf
Chickweed	Oxalis	Wild Carrot
Clover	Peppergrass	Wild Garlic
Cocklebur	Pigweed	Wild Lettuce
Dandelion	Plaintains	Wild Onion
Dock	Poison Ivy	Yarrow
Galinsoga	Poison Oak	And many
Ground Ivy	Purslane	other broadleaf
Hawkweed	Ragweed	weeds
Heal-all	Sheep Sorrel	

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. If contact with eyes occurs flush with plenty of water.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not store contact with fertilizers, seeds, insectic. or fungicides. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs.

When container is empty, wash thoroughly and destroy. Never re-use. Destroy by perforating or crushing, and burying in a safe place. Do not burn.

Under certain conditions possible injury may occur to bentgrass, St. Augustine and centipede grass.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 2217-570

880-672