

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

PARATHION 10-G

ETHYL PARATHION GRANULES

FOR CONTROL OF CORN ROOTWORM ON CORN

STABILIZED 15-30 Mesh

ACTIVE INGREDIENTS:

Parathion* 10%

INERT INGREDIENTS: 90%

TOTAL 100%

*O,O-Diethyl O-P-Nitrophenyl Thiophosphate

ACCEPTED

APR 21 1978

Under the Federal Insecticide, Fungicide, and Fossiticide Act, as amended, for the pesticide registered under EPA Reg. No. 2217-377

Study safety directions on this label. Consult your physician about obtaining an adequate supply of 1/100 grain Atropine tablets for emergency use.

DANGER! Keep Out of Reach of Children

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.
RAPIDLY ABSORBED THROUGH SKIN

REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS PROGRESSIVELY INCREASE SUSCEPTIBILITY TO ETHYL PARATHION POISONING.



POISON



ATROPINE IS AN ANTIDOTE — CONSULT PHYSICIAN FOR EMERGENCY SUPPLY

MADE BY  **pbi/gordon**
corporation
KANSAS CITY, KANSAS 64108

BEST DOCUMENT AVAILABLE

NET WEIGHT 50 POUNDS

DO NOT GET EYES, ON SKIN, ON CLOTHING

Wear clean rubber gloves, goggles and clean water proof or freshly laundered protective clothing (coveralls, rubber boots, cap, etc.). Destroy and replace gloves frequently.

Wash thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

DO NOT BREATHE DUST OR VAPOR

Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Ethyl Parathion. If handled outdoors, provide mechanical exhaust ventilation.

DO NOT STORE NEAR FOOD OR FEED PRODUCTS

Bury spillage; wash area thoroughly with a solution containing 1 lb. soda ash per gallon of water to which a water-soluble wetting agent has been added.

Bags: Destroy empty bags by burning.

Stay away from smoke and fumes.

Remove unauthorized persons from operating area.

This product will kill fish and wildlife. Keep out of any body of water.

Apply this product only as specified on this label.

FIRST AID TREATMENT

Call a Physician at once in all cases of suspected Parathion Poisoning. If symptoms or signs of poisoning (See Physician's Note), include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1/100 grains) at once.

Remove patient immediately from the area where Staypar is present. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of Staypar. If swallowed, induce vomiting by giving warm salty or soapy water. Have patient lie down and keep quiet.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON!

Physicians' Note: Warning symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

Treatment: Give atropine, preferable by injection, grains 1/100 three or more tablets at once and every hour as required until pupils dilate. Thirty or more tablets may be required during the first 24 hours. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

— DIRECTIONS FOR USE —

READ CAUTION STATEMENT

CORN ROOTWORM—Treat at planting time. On field corn apply 10 to 13 pounds of Staypar to the soil per acre as a band application (4 to 6 inches wide). Granules must be distributed evenly with properly calibrated equipment in the top 1/2" to 1" of soil.

NOTE—This method of application will not give effective control of white grubs and soil cutworms. Staypar is not suited for application to the soil when incorporated with dry fertilizer.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

531-567

EPA Reg. No. 2217-377

EPA Est. No. 2217-KS-1