



SEVIN* DUST

5%

4-14-69
2217-303

MANUFACTURED BY

GORDON CORPORATION
KANSAS CITY, KANSAS

— DIRECTIONS —

Apply SEVIN DUST 5% when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 3-day intervals. For best results apply with a dust gun. Do not apply in excess.

VEGETABLE INSECT CONTROL

ASPARAGUS: To control Asparagus Beetle and Apache Cicada. Do not repeat applications within three days.

BEANS (Green Beans, Lima Beans, Snap Beans, Cowpeas, Black-eyed) To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvetbean Caterpillar, Western Bean Cutworm, Armyworms, Corn Earworm, Stink Bugs and Tarnish Plant Bugs

SWEET CORN: To control Corn Earworm, Corn Rootworm adults, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetles, Sap Beetles, and Leafhoppers. For larvae in Whorl (buitworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control.

CUCUMBER, MELONS, PUMPKIN, SQUASH: To control Pickleworm, Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers and Squash Bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

POTATO, TOMATO, EGGPLANT, PEPPER: To control Colorado Potato Beetle, Flea Beetles, Leafhoppers, European Corn Borer, Fall Armyworm, Lace Bugs, Tomato Fruit Worm, Tomato Hornworm, Tarnish Plant Bugs, Stink Bugs

OKRA: To control Corn Earworm and Stink Bugs.

HEAD LETTUCE AND CARROTS: To control Flea Beetles, Harlequin Bug, Leafhoppers, Armyworms, Cabbage Caterpillars, Corn Earworm, Spittlebugs, Tarnish Plant Bugs and Stink Bugs. To control small Cabbage Loopers use on a 5 to 7 day schedule.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: To control Flea Beetles, Harlequin Bug, Armyworms, Cabbage Caterpillars, Corn Earworm. To control small Cabbage Loopers, use on a 5 to 7 day schedule.

SMALL FRUIT INSECT CONTROL

GRAPES: To control Grape Berry Moth, Japanese Beetles, June Beetles, Red-banded Leaf Roller, Grape Leaf Folder, Grape Leafhopper and Grape Leaf Skeletonizer. Apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers.

STRAWBERRIES: To control Meadow Spittlebug, Strawberry Leaf Roller and Strawberry Weevil. Do not apply within one day of harvest.

GRASSHOPPERS ON THE ABOVE CROPS

Apply SEVIN DUST 5% as often as necessary on crops to control grasshoppers.

PET USE

DOGS AND CATS: For ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks of age. Do not treat puppies less than four weeks of age.

In and around dog and cat quarters. For ticks and fleas dust liberally in and around sleeping quarters, kennels and dog houses by means of a shaker can, hand or dust gun.

FLOWERING PLANTS

ROSES, CARNATIONS, GLADIOLUS, MUMS, AZALEA: For control of Flea Beetles, Winter Leafhoppers, June Beetles, Lace Bugs, Leafhoppers, Mealy Bugs, Plant Bugs, Rose Aphid, Rose Slug, Scale Insects, and Thrips. Apply dust when air is calm. Cover upper and lower surfaces of leaves as well as stems and buds. Repeat at 3-day intervals or as needed.

NOTE

Allow 1 day between dusting and harvest of Asparagus and strawberries. Three days on Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce and Kohlrabi. Do not use on Leaf Lettuce within 14 days of harvest. Injury to young tender foliage may occur if plants are wet or in the presence of high humidity of several days duration following treatment.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do not use internally. Avoid contact with clothing or dust. Skin contact may be harmful. Avoid unnecessary contact with eyes. Wash face and hands thoroughly after use. Wear regular protective work clothes. Change to clean clothing daily.

NOTE FOR PHYSICIANS: Carbaryl is a moderate to very fast acting cholinesterase inhibitor. Atropine is antidotal.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. No conditions of use, conditions of sale, or compatibility with other chemicals are intended. If application equipment is used, the user must follow the directions on the label. U.S.A. Reg. No. 1017-101-01

SEVIN DUST

5%

AN INSECTICIDE

**VEGETABLES—SMALL FRUITS
FLOWERING PLANTS**

For Home Garden Use

ACTIVE INGREDIENTS:	
Carbaryl (1-Naphthyl N-Methylcarbamate)	5%
INERT INGREDIENTS:	95%
	TOTAL <u>100%</u>

*Reg. T. M. Union Carbide Corp. for 1-Naphthyl
N-Methylcarbamate

CAUTION: Keep Out of Reach of Children
See Side Panel for Additional Cautions

CONTROLS TICKS AND FLEAS ON PETS

611-1067

Net Weight 1 Pound

UNION CARBIDE
CORPORATION
2909 NORTH ZEEB ROAD
PORTLAND, OREGON 97201
REGISTERED TRADEMARK
OF UNION CARBIDE
CORPORATION