

SEVIN 50W SPRAY

A WETTABLE POWDER

This 2 lb. Bag Treats 5,000 Sq. Ft. of Lawns

controls
SOD WEBWORM
ANTS - FLEAS - LEAFHOPPERS
VEGETABLE, FRUIT AND ORNAMENTAL PESTS

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)	50%
INERT INGREDIENTS	<u>50%</u>
TOTAL	100%

*Reg. T. M. Union Carbide Corporation for 1-Naphthyl N-methylcarbamate. U. S. Pat. No. 2,903,478.

CAUTION: Keep Out of Reach of Children
See Side Panel for Additional Cautions

520 273

NET WEIGHT 2 POUNDS

*TM Registered, P.B.I.



SEVIN[®] 50W ACCEPTED SPRAY

MAY 31 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 2217-389 SUBJECT
TO ATTACHED COMMENTS.

SEVIN 50W SPRAY DIRECTIONS

SOD WEBWORM CONTROL —

When to spray: In late June and July where numerous moths can be seen flying low over Bluegrass lawns during early evening, spraying anytime within the next week will prevent damage from hatching worms. If lawn damage was heavy the preceding year, an early spray (May 15th) will prevent Spring damage from overwintering worms.

How to spray: To get spray down to soil level, mow lawn and rake out any heavy grass. Water lawn. Spray using a garden pressure sprayer or hose sprayer. Add SEVIN[®] 50W SOD WEBWORM SPRAY to water using at least 1 pint for every 2 cups insecticide. Thoroughly mix, then fill spray container with water to spray level.

Use 2 cupfuls in 10 to 15 gallons of water per 1000 square feet. Sprinkle after spraying to wash insecticide into soil level of turf. This 2 lb. package will treat 5000 square feet, one time.

This treatment will control ants, fleas, leafhoppers, and chinch bugs.

ORNAMENTAL INSECTS: Use 2 bs. per 100 gallons or 2 tablespoonfuls per gallon. Spray all foliage thoroughly. Repeat as necessary to control bagworms, blister beetles, boxelder bugs, flea beetles, lace bugs, leafhoppers, rose slugs, and green canker worms. Do not spray Boston Ivy or Virginia Creeper.

HOME ORCHARDS AND VEGETABLES: For best insect control through season plan on a regular spray schedule every 7 to 10 days. Two (2) pounds to 100 gallons is equivalent to 2 tablespoonfuls per gallon.

APPLES AND PEARS — Codling Moth, Plum Curculio, Apple Aphids, Green Fruit Worms, Japanese Beetles, Leaf Rollers, and San Jose Scale — As insects appear, spray, using 2 tablespoonfuls per gallon of water, repeat as needed for control, up to one day before harvest. To avoid fruit thinning, delay use until at least 30 days after bloom (not before second cover).

PEACHES, APRICOTS, PLUMS, AND CHERRIES — Plum Curculio, Oriental Fruit Moth, Lesser Peach Tree Borer, Peach Twig Borer, Cat Facing Bugs, and San Jose Scale — Mix 2 tablespoonfuls per gallon of water. Allow one day between spraying and harvest; three days on Apricots.

CUCUMBERS, MELONS, AND SQUASH — For Pickleworm, Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers, and Squash Bugs, use 2 tablespoonfuls per gallon. Avoid excessive applications since leaf injury may occur on tender foliage when wet and humidity is high. Allow 3 days before harvest.

TOMATOES, EGG PLANTS, PEPPERS — For Tomato Hornworm, and Fruitworm, Fall Armyworm, Stink Bugs, and Lace Bugs, use 2 tablespoonfuls per gallon. Thoroughly treat. Use until day of picking.

CABBAGE, BROCCOLI, AND CAULIFLOWER — For control of Small Cabbage Loopers, Flea Beetles, Armyworms, and Corn Earworms — Use 2 tablespoonfuls per gallon on a 7 to 10 day spray schedule, allowing 3 days before picking.

SWEET CORN — Corn Earworm, European Corn Borer, Flea Beetles, and Grasshoppers — Use 2 tablespoonfuls per gallon. Spray for earworms until silks are dry. Apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals as infestation warrants. Re-apply immediately after rain. For other pests, thoroughly treat entire plant every 2 weeks, starting when corn is knee high. Do not apply within 7 days of harvest if fodder is to be fed to livestock, otherwise no limitation. **NOTE:** Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

HUMAN CAUTIONS: HARMFUL IF SWALLOWED OR INHALED! Do not breathe dust. Do not take internally. Avoid breathing of spray. Contact with skin may be harmful. Wash thoroughly after using. Wear regular long sleeved work clothing. Change to clean clothing daily. Avoid storage near feed and foodstuffs. Keep children and pets off treated areas until the insecticide is washed into the soil and grass is thoroughly dry.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL CAUTIONS: Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse container. Destroy when empty. Apply this product only as specified on this label.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary Seller makes no claims other than those stated on this label.

620-273

EPA Reg. No. 2217-389-4A

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS