in 19 32	17-389 Notebrotion 4	
(A) CERPA United States Environmental Protect Office of Pesticide Programs (H Washington, DC 20460 Application for Pesticide Programs)	Amendment	
Section I		
1. Company/Product Number	2. EPA Product Manager 3. Proposed Classification	
2217-389	Dennis Edwards	
4. Company/Product (Name) Gordon's Sevin 50W Spray, A Wettable Powder	(PM 19)	
	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)	
5. Name and Address of Applicant (Include ZIP Code)	(b)(i), my product is similar or identical in composition and labeling	
PBI/Gordon Corporation PO Box 4090	to:	
1217 West 12th Street	EPA Reg. No.	
Kans as C ity, MO 64101		
	Product Name	
Section 11		
Amendment - Explain below	Final printed labels in response to Agency letter dated	
Resubmission in response to Agency letter dated		
Notification - Explain below.	"Me Too" Application. NOTIFICATION LABEL NOT REVIEWED	
	Other - explain below. PER PR NOTICE 84-6	
Notice 93-3, "Labeling Statement Prohibiting Application to Water", March 9, 1993. This is to certify that this notification meets every requirement of PR-Notice 88-6. (2) Deletion of the chemigation statement defined in <u>PR Notice 87-1</u> , "Label Improvement Program for Pesticides Applied Through Irrigation Systems." This product is intended for residential use (such as indoor, yard, or garden). (3) Revision of the limited warranty and disclaimer statement. (4) Three copies of the revised labeling are enclosed.		
Section III		
Material This Product Will Be Packaged In: Child-Resistant Packaging Unit Packaging Water	Soluble Packaging 2. Type of Container	
	Yes Motal	
	No Plastic Glass	
If "Yes," No. per If "Yes	Paper	
	ge wgt. container Other (Specify)	
submitted. 4. Size(s) of Retain Contents	ontainer 5. Location of Label Directions	
Label Container Container On Labeling accompanying product Lithograph Other (
Paper glued Other ()		
1. Contact Point (Complete items directly below for identification of indiv	idual to be contacted, if necessary, to process this application.)	
Name Title	Telephone No. (Include Area Code)	
James A. Armbruster, Ph.D. M	gr. Regulatory Services (816)421-4070	
Certification 6. Date Application		
I certify that the statements I have made on this form and all attachments thereto are true, accurate and comp'3±. Received I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or (Stamped) both under applicable law.		
2. Signature 3. Title		
J. L. Sundmarter M	gr. Regulatory Services	
4 Typed Name Late	——————————————————————————————————————	
James A. Armbruster, Ph.D. May 20, 1993		
EPA Form 8570-1 (Rev. 12-90) Previous editions are obsolete. White EPA File Copy (orginal) Yellow Applicant copy		

SEVIN[®] 50W SPRAY

A WETTABLE POWDER

This 2 lb. bag Treats 5,000 Sq. Ft. of Lawn

Controls: SOD WEBWORM - ANTS - FLEAS - LEAFHOPPERS VEGETABLE, FRUIT AND ORNAMENTAL PESTS

ACTIVE INGREDIENT:	
Carbaryl (1-napthyl N-methylcarbamate)	. 50.0%
INERT INGREDIENTS.	
TOTAL	100.0%

SEVIN[®] is the registered trademark of Rhone Poulenc for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN

NOTIFICATION EL NOT REVIEWED ER PR NOTICE DAI

CAUTION


See side panel for Precautionary Statements and Statement of Practical Treatment

NET WEIGHT 2 POUNDS

620/ AP052093

EPA REG. NC 221 -389 E A L.ST. NO 2217-KS-1

Mfd. by PBI/GORDON CORPORATION KANSAS CITY, MISSOURI 64101



STOPI READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid breathing vapors [dust or spray mist]. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

4

Statement of Practical Treatment

IF SWALLOWED: Induce vomiting and seek medical attention immediately. IF IN EYES OR ON SKIN: Flush eyes thoroughly with water: wash skin thoroughly with soap and water. NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

Environmental Hazards

This pesticide is extremely toxic to equatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Store in original container in a locked storage area inaccessible to children or pets.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold and discard bag in trash.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

- SOD WEBWORM CONTROL -

<u>When to spray:</u> In late June and July where numerous moths can be seen flying low over bluegrass lawn during early evening, spraying anytime within the next week will prevent damage from hatching worms. If lawn damage was heavy the preceding year, an early spray (May 15th) will prevent spring damage from over-wintering worms.

<u>How to spray:</u> To get spray down to soil level, mow lawn and rake out any heavy grass. Water lawn. Spray using a garden pressure sprayer or hose sprayer. Add SEVIN[®] 50W SPRAY to water using at least 1 pint for every 2 cups insecticide. Thoroughly mix, then fill spray container with water to spray level.

Use 2 cupfuls in 10 to 15 gallons of water per 1000 square feet. Sprinkle after spraying to wash insecticide into soil level of turf. This 2 lb. package will treat 5000 square feet, one time.

This treatment will control ants, fleas, leafhoppers, and chinch bugs.

~ •**?**

Use 2 lbs. per 100 gallons or 2 tablespoonfuls per gallon. Spray all foliage thoroughly. Repeat as necessary to control bagworms, blister beetles, boxelder bugs, flea beetles, lace bugs, leafhoppers, rose slugs, and green canker worms. Do not spray Boston Ivy or Virginia Creeper.

- HOME ORCHARDS AND VEGETABLES -

For best insect control through season plan on a regular spray schedule every 7 to 10 days. Two (2) pounds to 100 gallons is equivalent to 2 tablespoonfuls per gallon.

<u>APPLES AND PEARS - Codling Moth, Plum Curculio, Apple Aphids, Green Fruit Worms, Japanese Beetles, Leaf Rollers, and San Jose Scale</u> - As insects appear, spray, using 2 tablespoonfuls per gallon of water, repeat as needed for control, up to one day before harvest. To avoid fruit thinning, delay use until at least 30 days after bloom (not before second cover).

PEACHES, APRICOTS, PLUMS, AND CHERRIES - Plum Curculio, Oriental Fruit Moth, Lesser Peach Tree Borer, Peach Twig Borer, Cat Facing Bugs, and San Jose Scale - Mix 2 lablespoonfuls per gallon of water. Allow one day between spraying and harvest; three days on Apricots.

<u>CUCUMBER, MELONS, AND SQUASH - For Pickleworm, Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers, and Squash Bugs, use 2 tablespoonfuls per gallen.</u> Avoid excessive applications since leaf injury may occur on tender foliage when wet and humidity is high. Allow 3 days before harvest.

TOMATOES, EGG PLANTS, PEPPERS - For Tomato Hornworm, and Fruitworm, Fall Armyworm, Stink Bugs, and Lace Bugs, use 2 tablespoonfuls per gallon. Thoroughly treat. Use until day of picking.

CABBAGE, BROCCOLI, AND CAULIFLOWER - For control of Small Cabbage Loopers, Flea Beetles, Armyworms, and Corn Earworms - Use 2 tablespoons per gallon on a 7 to 10 day spray schedule, allowing 3 days before picking.

<u>SWEET CORN - Corn Earworm, European Corn Borer, Flea Beetles, and Grasshoppers</u> - Use 2 tablespoonfuls per gallon. Spray for earworms until silks are dry. Apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals as infestation warrants. Reapply immediately after rain. For other pests, thoroughly treat entire plant every 2 weeks, starting when corn is knee high. Do not apply within 7 days of harvest if fodder is to be fed to livestock, otherwise no limitation.

<u>Note:</u> Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

LIMITED WARRANTY AND DISCLAIMER The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of products or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

>